



CORRECTION

Correction to: Neuraxial anesthesia and bladder dysfunction in the perioperative period: a systematic review

Stephen Choi, MD · Pdraig Mahon, MD · Imad T. Awad, MBChB

Published online: 18 December 2017
© Canadian Anesthesiologists' Society 2017

**Correction to: Can J Anesth/J Can Anesth (2012)
59:681–703
<https://doi.org/10.1007/s12630-012-9717-5>**

The following row on page 691 of Table 3, as highlighted in bold under Incidence of Retention (%) should read “0” instead of “N/A”.

Table 3 Intrathecal anesthesia studies

Author/ Year	Study Design	Jadad Score	N	Surgical Class	Intrathecal Drug/Dose	Incidence of Retention (%)	PVR (mL)	Catheterization (%)	Infection (%)	Comments
Linares-Gil 2009 ⁴²	OBS	N/A	406	GEN (n = 219) ORTHO (n = 187)	3% Hyperbaric Lidocaine 0.9 mg·kg ⁻¹	0	N/A	0/406 (0)	N/A	Criteria for catherization not indicated

The original article can be found online at <https://doi.org/10.1007/s12630-012-9717-5>.

S. Choi, MD · I. T. Awad, MBChB (✉)
Department of Anesthesia, Sunnybrook Health Sciences
Centre, University of Toronto, 2075 Bayview Avenue,
Toronto M4N 3M5, Canada
e-mail: imad.awad@sunnybrook.ca

P. Mahon, MD
Department of Anaesthesia and Intensive Care Medicine, Cork
University Hospital, University College Cork, Cork, Ireland