CORRESPONDENCE

Uncertainty in using chest computed tomography in early coronavirus disease (COVID-19)

Rujittika Mungmunpantipantip, PhD · Viroj Wiwanitkit, MD

Received: 14 March 2020 / Revised: 15 March 2020 / Accepted: 15 March 2020 / Published online: 2 April 2020
© Canadian Anesthesiologists' Society 2020

To the Editor,

The title of a recent *Images* article in the *Journal* by Chen *et al.*¹ describes chest computed tomography (CT) findings in a pregnant woman with coronavirus disease (COVID-19) as “early coronavirus disease.” We argue that these findings are inconsistent with “early” COVID-19. Cases of COVID-19 were identified in Thailand shortly after its first appearance in China²; as of 14 March 2020, there have been 82 confirmed diagnoses of COVID-19 reported in Thailand. All Thai cases are confirmed by reverse transcriptase polymerase chain reaction (RT-PCR) in two reference labs, and all patients receive chest CT scans. Thirty-eight of the 82 cases were diagnosed during active screening for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in persons who had been in close contact to patients known to have COVID-19. Most of those screened were asymptomatic or reported only mild symptoms; none showed abnormal findings on chest CT scans. One asymptomatic Thai patient underwent general anesthesia for orthopedic surgery before being diagnosed with COVID-19 in the postoperative period. This case also had normal CT findings at the time of diagnosis.

Hu *et al.* reported that 29.2% of asymptomatic patients from China who were infected with SARS-CoV-2 and who

presented with no or mild symptoms at the time of diagnosis by RT-PCR test had a normal chest CT scan.³ Asymptomatic SARS-CoV-2 infection is possible and has been previously reported in Thailand and Croatia.⁴ COVID-19 may be asymptomatic in its early stages even though viral RNA can be detected by RT-PCR screening.⁴ We therefore discourage the consideration of CT findings when ruling out cases of COVID-19.

Conflicts of interest None.

Funding statement None.

Editorial responsibility This submission was handled by Dr. Hilary P. Grocott, Editor-in-Chief, *Canadian Journal of Anesthesia*.

References

1. Chen R, Chen J, Meng Q. Chest computed tomography images of early coronavirus disease (COVID-19). *Can J Anesth* 2020. DOI: <https://doi.org/10.1007/s12630-020-01625-4>.
2. Yasri S, Wiwanitkit V. Wuhan coronavirus outbreak and imported case. *Adv Trop Med Pub Health Int* 2020; 10: 1-2.
3. Hu Z, Song C, Xu C, *et al.* Clinical characteristics of 24 asymptomatic infections with COVID-19 screened among close contacts in Nanjing, China. *Sci China Life Sci* 2020. DOI: <https://doi.org/10.1007/s11427-020-1661-4>.
4. Joob S, Ronald M. Asymptomatic COVID-19: an important clinical consideration. *Adv Lab Med Int* 2020; 10: 4-5.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This letter is accompanied by a reply. Please see *Can J Anesth* 2020; 67: this issue.

R. Mungmunpantipantip, PhD (✉) ·
Medical Center, Bangkok, Thailand
e-mail: rujittika@gmail.com

V. Wiwanitkit, MD
Dr DY Patil University, Pune, India

Hainan Medical University, Haikou, China