CORRESPONDENCE

In reply: Uncertainty in using chest computed tomography in early coronavirus disease (COVID-19)

Rong Chen, MD · Qing-tao Meng, MD, PhD

Received: 19 March 2020 / Revised: 19 March 2020 / Accepted: 19 March 2020 / Published online: 2 April 2020
© Canadian Anesthesiologists' Society 2020

To the Editor,

The recent letter by Drs Mungmunpuntipantip and Wiwanitkit¹ argues that our report about chest computed tomography (CT) findings in a pregnant woman with coronavirus disease (COVID-19)² are inconsistent with “early” COVID-19. In doing so, they discourage the consideration of CT findings when ruling out cases of COVID-19. Nevertheless, we think that their concern may have been guided by an incomplete understanding of the COVID-19 epidemic. As of 17 March 2020, more than 50,000 people have been confirmed with COVID-19 in Wuhan. During the period of the most severe increases in COVID-19 cases, the supply of nucleic acid detection kits could not meet the demands of the sharp increase in COVID-19 patients. So the diagnosis of COVID-19 followed the New Coronavirus Pneumonia Prevention

and Control Program (fifth edition)^A issued by the National Health Commission of China that took “suspected cases with pneumonia imaging features” as the clinical diagnosis case standard in Hubei Province. This allowed more COVID-19 patients to receive more timely treatment.

In earlier stages of the disease progression, the changes of chest CT scan were mainly congestion, edema, and exudation. For example, Shi *et al.* reported that “COVID-19 manifests with chest CT imaging abnormalities, even in asymptomatic patients, with rapid evolution from focal unilateral to diffuse bilateral ground-glass opacities that progressed to or co-existed with consolidations”.³ It is worth noting that only a single ground-glass opacity can be seen in some patients at an early stage, and the range significantly increases in the short-term re-examination.⁴ When a patient’s condition improves, a few fibrous streaks may appear. On the contrary, when the condition is more advanced, diffuse lesions appear in the lungs, and the density of both lungs increases, showing a “white lung”, which seriously affects the patient’s lung function.⁴

Importantly, all diagnostic methods have disadvantages. In China, 29.2% of asymptomatic patients who were found to be infected with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and presented with either no or mild symptoms at the time of diagnosis by real-time reverse-transcription polymerase chain reaction (RT-PCR) test had a normal chest CT scan.⁵ In addition, there were also some patients with positive chest CT findings who presented with a negative RT-PCR for SARS-CoV-2.⁶ Ai

R. Chen, MD · Q. Meng, MD, PhD (✉)
Department of Anesthesiology, Renmin Hospital of Wuhan University, Wuhan, China
e-mail: mengqingtao2018@126.com

Department of Anesthesiology, East Hospital, Renmin Hospital of Wuhan University, Wuhan, China

^A National Health Commission of China. New Coronavirus Pneumonia Prevention and Control Program (5th ed.). Available (in Chinese) from URL: <http://www.nhc.gov.cn/yzygj/s7653p/202002/d4b895337e19445f8d728fcfa1e3e13a/files/ab6bec7f93e64e7f998d802991203cd6.pdf> (accessed March 2020).

et al. showed that 60–93% of cases had an initial positive CT consistent with COVID-19 prior to (or parallel with) the initial positive RT-PCR results. Forty-two percent (24/57) of cases showed improvement in follow-up chest CT scans before the RT-PCR results turned negative.⁷ Thus, chest CT may still be considered as one of the primary tools for detecting COVID-19 in epidemic areas.

We therefore still suggest that the combined assessment of CT imaging features with clinical and laboratory findings could facilitate diagnosis of COVID-19 earlier and more accurately. This may also help curb the spread of the COVID-19 pandemic.

Conflicts of interest None.

Funding statement None.

Editorial responsibility This submission was handled by Dr. Hilary P. Grocott, Editor-in-Chief, *Canadian Journal of Anesthesia*.

References

1. Mungmunpuntipantip R, Wiwanitkit V. Uncertainty in using chest computed tomography in early coronavirus disease (COVID-19). *Can J Anesth* 2020; <https://doi.org/10.1007/s12630-020-01639-y>.
2. Chen R, Chen J, Meng Q. Chest computed tomography images of early coronavirus disease (COVID-19). *Can J Anesth* 2020; <https://doi.org/10.1007/s12630-020-01625-4>.
3. Shi H, Han X, Jiang N, *et al.* Radiological findings from 81 patients with COVID-19 pneumonia in Wuhan, China: a descriptive study. *Lancet Infect Dis* 2020; [https://doi.org/10.1016/S1473-3099\(20\)30086-4](https://doi.org/10.1016/S1473-3099(20)30086-4).
4. Pan Y, Guan H, Zhou S, *et al.* Initial CT findings and temporal changes in patients with the novel coronavirus pneumonia (2019-nCoV): a study of 63 patients in Wuhan, China. *EurRadiol* 2020; <https://doi.org/10.1007/s00330-020-06731-x>.
5. Hu Z, Song C, Xu C, *et al.* Clinical characteristics of 24 asymptomatic infections with COVID-19 screened among close contacts in Nanjing, China. *Sci China Life Sci* 2020; <https://doi.org/10.1007/s11427-020-1661-4>.
6. Xie X, Zhong Z, Zhao W, Zeng C, Wang F, Liu J. Chest CT for typical 2019-nCoV pneumonia: relationship to negative RT-PCR testing. *Radiology* 2020; <https://doi.org/10.1148/radiol.2020200343>.
7. Ai T, Yang Z, Hou H, *et al.* Correlation of chest CT and RT-PCR testing in coronavirus disease, (COVID-19) in China: a report of 1014 cases. *Radiology* 2019; <https://doi.org/10.1148/radiol.2020200642>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.