CORRESPONDENCE

In reply: Use of a second ventilatory circuit when using a double-lumen endotracheal tube

Gerard Bruin, MD, FRCPC

Received: 18 March 2020 / Revised: 18 March 2020 / Accepted: 18 March 2020 / Published online: 30 March 2020
© Canadian Anesthesiologists' Society 2020

To the Editor,

In their comments on my recently published letter¹ in the *Journal*, Bussires *et al.*² discuss other monitoring options to detect a leak around the bronchial cuff of a double-lumen endotracheal tube (DL-ETT). A simple, though not specific, test for the leak—and one not mentioned by them—is to use fresh gas flows of 500 mL·min^{−1} or less. Failure of the ventilator bellows to refill while using this low fresh gas flow can indicate a possible leak around the bronchial cuff. Nevertheless, none of these methods, including spirometry, will be reliably sensitive if the bronchial cuff is over-filled or if the DL-ETT is too large. That is, too much air in the bronchial cuff can prevent detection of an early leak around the cuff. It can also cause an outward (from the carina) avulsion of the DL-ETT. This is why we recommended routine testing for the correct amount of air in the bronchial cuff at the time of initial placement (as indicated in the electronic supplementary material [ESM] video that accompanied the letter; <https://vimeo.com/370217402>; time stamp: 5 min 26 sec).

We have encountered situations where a size 35Fr DL-ETT passes easily into the bronchus, yet no air is needed in the bronchial cuff for an effective seal when we do our leak test. When we do the pull out test while observing the carina using the flexible bronchoscope (ESM video; time stamp: 5 min 51 sec), we have found that the technique is just as sensitive for detecting outward movement. In those cases, the filling of the bag is slower than shown in the video. If there is any resistance to advancing the DLT into

the bronchus, we consider another technique for lung isolation.

Dr. Bussires suggests using a separate oxygen blender towards the end of thoracic surgical procedures just prior to lung re-inflation. I agree that this would be helpful to minimize reabsorption atelectasis. Another option is to switch to two-lung ventilation and decrease the fraction of inspired oxygen to the lowest tolerable level. I look forward to other methods becoming available to continuously monitor DL-ETT position. Even with a more direct method of monitoring DL-ETT position, monitoring for a leak is simple and will likely continue to be a useful adjunct.

Conflicts of interest None.

Funding statement None.

Editorial responsibility This submission was handled by Dr. Hilary P. Grocott, Editor-in-Chief, *Canadian Journal of Anesthesia*.

References

1. Bruin G. Advantages of a second ventilation circuit when using a double-lumen endotracheal tube. *Can J Anesth* 2020; DOI: <https://doi.org/10.1007/s12630-019-01553-y>.
2. Bussires J, Cournoyer C, Couture EJ. Use of a second ventilatory circuit when using a double-lumen endotracheal tube. *Can J Anesth* 2020; DOI: <https://doi.org/10.1007/s12630-020-01641-4>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

G. Bruin, MD, FRCPC (✉) ·
Trillium Health Partners, Credit Valley Site, Mississauga, ON,
Canada
e-mail: gerardbruinabc@gmail.com