



Correction to: Silicon (Si) Application Improved the Antioxidant Response and Grain Yield Formation in rice Under High Temperature Conditions

Huizi Deng¹ · Xinyi Wang¹ · Siying Deng¹ · Long Zhang¹ · Umair Ashraf² · Muhammad Imran¹ · Hua Tian¹ · Xiangru Tang¹ · Zhaowen Mo¹

Published online: 14 March 2023
© The Author(s), under exclusive licence to Springer Nature B.V. 2023

Correction to: Silicon

<https://doi.org/10.1007/s12633-023-02354-1>

For errors in the article, we have made the following detailed notes:

Description of the stage in Figure 1(b), heading stage should be corrected to maturity stage; Variety names in Figures 1, 2, 3 and 4, NR-OR should be corrected to NO-OR.

The correct figure is shown below.

The original article has been corrected.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s12633-023-02354-1>.

✉ Xiangru Tang
tangxr@scau.edu.cn

✉ Zhaowen Mo
scaumozhw@126.com

Huizi Deng
deng_hz@126.com

Xinyi Wang
1010068440@qq.com

Siying Deng
dengsy729@163.com

Long Zhang
2918977869@qq.com

Umair Ashraf
umair.ashraf@ue.edu.pk

Muhammad Imran
Imran_m1303@yahoo.com

Hua Tian
tianhua@scau.edu.cn

¹ State Key Laboratory for Conservation and Utilization of Subtropical Agro-Bioresources, College of Agriculture, South China Agricultural University, Guangzhou 510642, People's Republic of China

² Department of Botany, Division of Science and Technology, University of Education, Lahore, Punjab 54770, Pakistan




Figure 1 Effects of Si application on SOD activity of different rice genotypes at (a) heading stage, and (b) at maturity stage under high temperature conditions. Si0 represents the silicon application as 0 g/pot, Si12 represents the silicon application as 12 g/pot. The capped bars represent the SDs. Columns with different lowercase letter(s) differ statistically at $P < 0.05$




Figure 2 Effects of Si application on POD activity of different rice genotypes at (a) heading stage, and (b) at maturity stage under high temperature conditions. Si0 represents the silicon application as 0 g/pot, Si12 represents the silicon application as 12 g/pot. The capped bars represent the SDs. Columns with different lowercase letter(s) differ statistically at $P < 0.05$




Figure 3 Effects of Si application on CAT activity of different rice genotypes at (a) heading stage, and (b) at maturity stage under high temperature conditions. Si0 represents the silicon application as 0 g/pot, Si12 represents the silicon application as 12 g/pot. The capped bars represent the SDs. Columns with different lowercase letter(s) differ statistically at $P<0.05$




Figure 4 Effects of Si application on MDA contents of different rice genotypes at (a) heading stage, and (b) at maturity stage under high temperature conditions. Si0 represents the silicon application as 0 g/pot, Si12 represents the silicon application as 12 g/pot. The capped bars represent the SDs. Columns with different lowercase letter(s) differ statistically at $P<0.05$