



Check for updates

Correction to: An in vitro and clinical dose-finding study of antifoaming effects of simethicone during colonoscopy

Marguerite J. Kutyla^{1,2} · Ayesha Shah^{1,3} · Jerome Elson¹ · Vera Meeusen¹ · Sam O'Connor^{1,3} · Luke F. Hourigan^{1,3} · Gerald J. Holtmann^{1,3}

Published online: 7 February 2020

© Indian Society of Gastroenterology 2020

Correction to: Indian Journal of Gastroenterology 38(3): 268–272

<https://doi.org/10.1007/s12664-019-00966-3>

In the above article, due to probable typo error with the picture and legend, the correct Fig. 1 and the Legend to the Fig. 1 are printed here:




Fig. 1 Side view of standard detergent solution (left) with a 200 mg, b 20 mg, and c 2 mg simethicone per 100 mL (right)

The online version of the original article can be found at <https://doi.org/10.1007/s12664-019-00966-3>

✉ Gerald J. Holtmann
gerald.holtmann@health.qld.gov.au

¹ Department of Gastroenterology and Hepatology, Princess Alexandra Hospital, Brisbane, QLD, Australia

² School of Pharmacy, University of Queensland, Brisbane, QLD, Australia

³ Faculty of Medicine, University of Queensland, Brisbane, QLD, Australia

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.