

# Androgen Receptor Splice Variants in the Era of Enzalutamide and Abiraterone

Mary Nakazawa · Emmanuel S. Antonarakis · Jun Luo

Received: 29 April 2014 / Accepted: 10 July 2014 / Published online: 22 July 2014  
© Springer Science+Business Media New York 2014

**Abstract** The FDA approvals of enzalutamide and abiraterone have rapidly changed the clinical landscape of prostate cancer treatment. Both drugs were designed to further suppress androgen receptor (AR) signaling, which is restored following first-line androgen deprivation therapies. Resistance to enzalutamide and abiraterone, however, is again marked by a return of AR signaling, indicating a remarkable “addiction” of prostate cancer cells to the AR pathway. Several mechanisms of castration resistance have been uncovered in the past decades, featuring a wide spectrum of molecular alterations that may explain sustained AR signaling in castration-resistant prostate cancers (CRPC). Among these, the androgen receptor splice variants (AR-Vs), particularly variant 7 (AR-V7), have been implicated in resistance to enzalutamide and abiraterone in preclinical studies, and they cannot be targeted by currently available AR-directed drugs. Drug development for AR-V-associated CRPC may therefore be necessary to augment the preexisting treatment repertoire. In this mini-review, we will discuss general mechanisms of resistance to AR-directed therapies, with a focus on the role of androgen receptor splice variants in the new era of treating advanced prostate cancer with enzalutamide and abiraterone.

## Introduction

Prostate cancer (PCa) is the most common non-skin cancer in men, responsible for approximately 258,000 deaths annually

worldwide [43]. Clinical management of advanced PCa hinges on the use of surgical or medical castration, an approach known as androgen deprivation therapy (ADT) [81]. ADT exploits the cancer’s dependence on androgen signaling, first revealed by Huggins in 1941 [42]. Today, ADT is not only used for those who are diagnosed with (or develop) advanced disease, but also in a subset of patients as an adjuvant to local therapy. The main goals of ADT are to decrease tumor burden, enhance quality of life, and improve survival [8].

While the vast majority of patients with advanced prostate cancer respond to initial ADT, given time, virtually all patients progress to a disease state known as castration-resistant PCa (CRPC) [45, 75]. CRPC describes the most advanced state of PCa, when patients demonstrate serial rising prostate-specific antigen (PSA) values or radiographic/clinical progression despite castrate levels of androgen (usually defined as serum testosterone <50 ng/dL) [34]. In the CRPC state, androgen signaling remains active and is mediated through the androgen receptor (AR) [30, 80]. A whole spectrum of molecular alterations, both primary and acquired, has been identified that results in sustained AR signaling in the castrate setting [37, 21].

Until 2004, the only Food and Drug Administration (FDA)-approved therapy for CRPC was the chemotherapeutic agent, docetaxel [27]. As such, until recently, the treatment paradigm for PCa was to use luteinizing hormone-releasing hormone (LHRH) agonists/antagonists and/or antiandrogens (AR antagonists), followed by docetaxel after development of metastatic CRPC. In the last decade, a variety of other therapies has been approved or is in development; the most notable of which are abiraterone and enzalutamide, two FDA-approved AR-targeted therapies demonstrating significant improvement in survival in the CRPC setting (both before and after docetaxel use) [77, 24, 7, 70]. Enzalutamide is a second-generation AR antagonist with

M. Nakazawa · J. Luo (✉)  
The James Buchanan Brady Urological Institute and Department of Urology, The Johns Hopkins School of Medicine, 600 N Wolfe St, 411 Marburg Bldg, Baltimore, MD 21287, USA  
e-mail: jluo1@jhmi.edu

E. S. Antonarakis  
Department of Oncology, The Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins, 1650 Orleans Street, CRB1-1M45, Baltimore, MD 21287, USA

properties superior to its first-generation counterparts such as flutamide, bicalutamide, and nilutamide. Enzalutamide was developed to bind to the AR ligand-binding domain (LBD) with high affinity, to reduce AR nuclear translocation efficiency, to disrupt AR binding to androgen response elements (ARE) on DNA, and to impair recruitment of coactivators to the AR transcriptional complex [90, 74]. Abiraterone is a CYP17A1 inhibitor designed to target steroid biosynthesis in CRPC and acts to inhibit androgen synthesis from alternative sources, such as from within the tumor or the adrenal gland [5]. The efficacy of these next-generation endocrine treatments confirms that the AR-signaling axis is still a viable target after progression to CRPC. While the current CRPC drugs were designed for and approved for late-stage patients, the prospect of using enzalutamide and abiraterone for hormone-sensitive patients is gaining attention. Data from a phase II study involving the administration of abiraterone in combination with androgen deprivation therapy prior to radical prostatectomy showed robust PSA responses and a high rate of pathologic complete (or near-complete) responses in high-risk localized PCa patients [67, 87]. In another phase II trial, enzalutamide was used (without androgen deprivation therapy) in men with hormone-sensitive disease, resulting in a very high rate of PSA responses [89]. The use of these next-generation agents could possibly provide for better AR ablation in the early stages of prostate cancer and stall progression to the lethal CRPC phenotype.

The new paradigm of treating CRPC with enzalutamide and abiraterone provides opportunities to dissect the drug resistance mechanisms to these agents. It is now established that a proportion of patients are primary refractory (i.e., show no response) to therapy or develops resistance despite showing an initial response. Because resistance is often marked by a return of AR signaling, it is critical to dissect mechanisms of sustained AR signaling in the presence of potent AR pathway suppressors. The extent to which the spectrum of molecular alterations acquired after first-line therapy contributes to resistance in this new treatment setting remains largely uncharacterized. The constitutively active AR splice variants (AR-Vs) represent a conceptually simple and biologically plausible mechanism of resistance to abiraterone and enzalutamide. These AR-Vs contain an intact N-terminal domain (NTD) and DNA-binding domain (DBD) but lack the c-terminal ligand-binding domain (LBD), the intended therapeutic target of abiraterone and enzalutamide (Fig. 1). In addition to AR-Vs, a number of novel mechanisms of resistance have been recently investigated in the specific setting of treatment with enzalutamide and abiraterone (summarized in Fig. 2), including drug-specific molecular alterations acquired during treatment with the two drugs. In this review, we will provide a succinct summary of these resistance mechanisms, with a particular focus on the androgen receptor splice variants.

## General Mechanisms of Resistance and Progression to CRPC

### AR Alterations


In the normal prostate gland, dihydrotestosterone (DHT) is the primary androgen mediating AR signaling. Androgen binding to the ligand-binding domain (LBD) of the AR causes receptor dimerization, translocation to the nucleus, and subsequent transcription of androgen-responsive genes; this signaling in PCa cells sustains tumor growth and proliferation. Upon initiation of ADT (and withdrawal of these ligands), PCa cells undergo apoptosis or are held in G1 cell cycle arrest. Cells with primary resistance to androgen ablation, or surviving cells held in cell cycle arrest that eventually adapt to low-androgen conditions, later emerge to resume proliferation [76]. The transition from hormone-sensitive to castration-resistant PCa is also consistently coupled with an increase in AR messenger (m)RNA expression, as demonstrated by microarray profiling [20]. Multiple mechanisms involving the AR-signaling axis have been proposed to account for the emergence of CRPC [21, 65, 79]. Among these, alterations involving the AR gene and its products (including AR amplification [92, 58, 14], gain-of-function mutation [86, 82], and overexpression [20]) are the most specific to CRPC.

### Sustained Androgen Levels

Incomplete ablation of circulating androgens can cause direct activation of the AR, resulting in sustained tumor growth. Studies have shown that surgical and medical castrations often do not eliminate androgens from the tumor tissue [76, 88]. In fact, residual levels of both testosterone and DHT in CRPC patients were found to be adequate to sustain androgen signaling [59, 88]. The adrenal gland is a contributor to levels of circulating androgens in the CRPC state [103]. In addition, testosterone and DHT synthesis may also occur via de novo pathways, possibly involving the conversion of acetic acid, a cholesterol precursor to steroids including DHT [57]. Thus, the continued presence of androgens (at lower but sufficient levels), coupled with more abundant androgen receptor levels, appear to be a prevailing explanation for CRPC.

### Growth Factors and AR Coregulators

The AR can also be activated via alternative pathways and molecules in a low-androgen state. Several growth factors, namely insulin-like growth factor 1 (IGF-1), keratinocyte growth factor (KGF), and epidermal growth factor (EGF), are able to activate the AR directly [23]. In addition, coactivators that enhance AR function may be involved in increasing the sensitivity of the AR to low levels of androgens or alternative ligands [80]. For example, a coactivator protein known as ARA70 was shown to increase AR responsiveness



**Fig. 1** Transcript structures for the full-length AR and major splice variants AR-V7 and AR<sup>v567ES</sup>. The human AR gene has eight canonical exons, color coded in relation to the AR protein domains they encode, including the N-terminal domain (NTD) (*purple*), DNA-binding domain (DBD) (*green*), the hinge region (*blue*), and the ligand-binding domain (LBD) (*red*). AR-V7 retains the first three canonical exons, followed by variant-

specific cryptic exon 3 within intron 3, leading to premature translation termination after 16 variant-specific amino acids (*gray*). AR<sup>v567ES</sup> retains the first four canonical exons, followed by exon 8. Skipping of exons 5–7 leads to disruption of the open reading frame and a variant-specific peptide of ten amino acids (*gray*). Peptide positions are marked according to GRCh36/hg18 human genome sequences (not drawn to scale)

to estradiol, suggesting that ARA70 can induce AR activity using an alternative ligand [97].

### Novel Mechanisms of Enzalutamide/Abiraterone Resistance

#### Reciprocal Regulation of AR and PI3K

Studies have also shown that AR inhibition could lead to upregulation of other oncogenic pathways to sustain PCa growth. Along with AR, the phosphatidylinositol 3-kinase (PI3K) pathway is another signaling pathway that promotes prostate carcinogenesis. The PI3K enzymes play a role in mediating signal transduction across the cellular membrane [73]. The AR and PI3K pathways are known to be cross-regulated by reciprocal feedback. That is, the inhibition of AR signaling through androgen deprivation and other manipulations results in a heightened activation of the PI3K pathway, and vice versa [17].

#### Alternative Steroid Receptors

Recently, there has been increasing interest in the role of the glucocorticoid receptor (GR) in PCa. GR activation has been shown to drive the expression of AR target genes, suggesting that AR-signaling inhibition through antiandrogens can be bypassed via glucocorticoid signaling [3]. Studies from clinical samples showed that GR expression is only present in


approximately 38 % of untreated PCa, but exists in higher proportion (78 %) in androgen-deprived PCa [85]. Indeed, in enzalutamide-resistant patients, GR expression is heightened [3]. Earlier studies also identified an AR mutation, L701H (or L701H coupled with T877A) which allows the AR to respond to glucocorticoids to sustain tumor growth [102]. Further, it is known that both the AR and GR bind to a common response element on DNA and regulate the same genes. This could imply that both the GR and its ligands may sustain AR signaling in the CRPC state [72], depicted in Fig. 2, pathway 4 and 5.

The progesterone receptor (PR), another member of the steroid receptor family, was shown to be increased in CRPC [13] and may be yet another resistance mechanism due to continued progesterone production. Two isoforms of the PR (isoforms A and B) are expressed in stromal fibroblast and smooth muscle cells of the prostate and are known to regulate cell proliferation [98, 32]. While evidence supporting a role of PR in resistance to abiraterone and enzalutamide is currently weak, further studies could implicate the role of PR signaling in CRPC in the absence of circulating androgens, depicted in Fig. 2, pathway 6.

### Therapy-Specific Mechanisms of Resistance

#### Resistance to Enzalutamide

An AR gain-of-function mutation known to confer resistance to enzalutamide has been identified in PCa cell lines.



**Fig. 2** A summary of candidate molecular alterations leading to the return of AR signaling in spite of treatment with enzalutamide and abiraterone. These alterations include 1 androgen receptor splice variants, 2 AR F876L mutation, 3 steroidogenic enzymes, 4 glucocorticoid receptor overexpression, 5 AR L701H mutation, and 6 progesterone receptor

activation. Diverse mechanisms may operate independently or cooperatively. The list is not intended to be comprehensive and emphasis was on those under investigation in the specific setting of the two new drugs. Dissection of the major drivers of resistance to enzalutamide and abiraterone will direct future prostate cancer drug development

Specifically, the missense mutation F876L (depicted in Fig. 2, pathway 2) in the LBD of the AR confers agonist properties to enzalutamide, sustaining AR signaling in the presence of the drug [6]. This has been verified as an acquired mechanism of resistance in patients receiving enzalutamide [44]. Further, ARN-509 (a newer AR antagonist with a higher therapeutic index compared to enzalutamide [22]) also demonstrates agonist properties to the F876L mutation [46]. The clinical impact of these findings will need to be determined in larger collections of patient samples, but the F876L mutation will likely account for a small proportion of cases of acquired resistance to enzalutamide (and ARN-509).

### Resistance to Abiraterone

Studies with CRPC xenografts have shown that various genes in the androgen biosynthesis pathway are upregulated as a result of treatment with abiraterone, notably the target gene CYP17A1 itself [60]. Another study has shown that DHT can be synthesized from 5 $\alpha$ -androstane-3-one instead of testosterone, the usual precursor to DHT [18, 19]. Further, recently, a gain-of-function mutation (N367T) was shown to produce 3 $\beta$ -hydroxysteroid dehydrogenase type 1 (3 $\beta$ HSD1) resistant to ubiquitination, which results in the accumulation of this enzyme responsible for DHT synthesis [18]. Depicted in Fig. 2, pathway 3, all of these represent molecular adaptations

of the AR-signaling pathway to evade the effects of abiraterone-mediated downregulation of androgen synthesis. Similarly, the clinical impact of these findings needs to be confirmed in large-scale studies.

## Androgen Receptor Splice Variants

### Overview

Since 2008, multiple AR splice variants (AR-V) lacking the AR ligand-binding domain have been identified and characterized [25, 44]. Some AR-Vs, notably AR-V7 and AR<sup>v567es</sup>, are known to be constitutively active, allowing for the activation of AR signaling in the absence of natural ligand binding [38, 84, 69], depicted in Fig. 2, pathway 1. Levels of AR-V are known to correlate with PCa progression and CRPC [49, 38, 31]. Additionally, expression of AR-V7 increases markedly upon androgen deprivation [99, 40, 95]. AR-V expression is found to be common in tumor metastases; furthermore, levels of variant expression directly correlate with accelerated disease progression and shorter cancer-specific survival [36, 101]. Experiments on mice have shown that AR<sup>v567es</sup> can induce autonomous tumor formation and proliferation [54], and AR-V7 expression leads to the expression of tumor promoting growth factors such as TGFβ2 and IGF1 [83]. These findings suggest that AR-Vs are biologically significant and may contribute to clinical progression of prostate cancer.

### Diversity of AR-Vs

The transcript structures of 18 androgen receptor splice variants have been fully characterized [44]. The majority of these splice variants arise through splicing of intronic sequences (i.e., cryptic exons). In addition to the sequence features that distinguish the different variants, other features critical for clinical translation have been characterized [69, 26, 38, 31, 95, 39], including their relative abundance, functional activity, and evidence for the corresponding protein product. On the basis of these studies, AR-V7 was determined to be the most important AR splice variant because it is the most abundant AR-V, its expression increases by ~20-fold in CRPC specimens, it is constitutively active in a ligand-independent manner, and it is detectable as a protein using variant-specific antibodies.

### Genomic Functions of AR-V7

Relative to the full-length AR (AR-FL), AR-V7 was initially described as a low-abundance transcript, at ~1–5 % of the full-length AR in the vast majority of CRPC specimens (as measured by quantitative RT-PCR) [38]. This observation has

been confirmed by independent investigators [94]. Because AR-V7 coexists with the full-length AR, it is critical to dissect the differential functions of AR-V7 and AR-FL, respectively. To determine the transcriptional output of AR signaling mediated by AR-V7 in the context of suppressed AR-FL signaling, transcriptional changes driven by forced expression of AR-V7 in the presence or absence of AR-FL signaling identified a set of cell cycle genes enriched for upregulation independent of the presence or absence of ligand-mediated AR-FL signaling [40]. On the other hand, top gene sets enriched for upregulation by ligand-dependent AR-FL signaling are dominated by those related to biosynthesis, metabolism, and secretion [40]. These findings suggest that although AR-V7 and AR-FL both regulate canonical AR target genes, they are associated with distinct transcriptional programs in CRPC.

In addition, it is known that AR-V7 can be acutely induced following suppression of AR-FL in two cell lines, VCaP and LNCaP95 [40]. Corroborating the functional distinctions between AR-FL and AR-Vs, endogenous induction of AR-V7 was accompanied by genome-wide changes of gene expression consistent with a shift of AR signaling mediated by AR-Vs [40]. Therefore, constitutively active AR-Vs appear to mediate a broader function than simply another mechanism to “rescue” canonical AR signaling. To this end, the emergence of AR-Vs in CRPC may drive a lethal mitotic phenotype while maintaining certain components of AR signaling. This notion is supported by other correlative studies in clinical specimens as well as animal xenografts [40, 101, 36]. Collectively, these studies show that although AR-V7 is expressed at levels that are substantially lower than those of AR-FL, it is sufficient to mediate a shift toward a transcriptional program mediated by AR-V7 when AR-FL is suppressed. Therefore, targeting the AR-V and its transcriptome is an area of priority for discovery and development of novel approaches to overcome resistance to abiraterone and enzalutamide.

### Biphasic and Context-Specific Functions of AR-Vs

In a recent study, AR-V-dependent genes were found to be induced at low receptor levels but repressed at high receptor levels [49]. This observation mirrors the canonical biphasic androgen-stimulated (i.e., AR-FL-mediated) growth response observed in cell line models. Study findings also suggest that AR-Vs mediate enzalutamide resistance by re-activating the AR-FL transcriptional programs, rather than by targeting a unique set of AR-V-associated genes [49]. These study findings are in contrast to those from our previous study [40], in which we showed distinct transcriptional programs mediated by AR-FL and AR-Vs, using two different cell line models in which AR-V7 can be induced following suppression of AR-FL. AR-regulated genes are known to be cell-context specific [12]. Because the two studies were performed in different cell

lines, cell-context differences may explain the seemingly opposing findings, although different methodologies utilized in the studies may also be a factor. Nevertheless, we posit that the relative lower AR-V levels (relative to AR-FL) may not fully “rescue” the androgen-regulated genes suppressed by potent AR-FL inhibition. Instead, AR-Vs may mediate genomic functions distinct from those mediated by the AR-FL, although the underlying mechanism remains unclear. It is possible that a distinct set of coregulators, target genes, and other DNA regulatory elements may collectively mediate AR-V signaling.

#### Molecular Origin of AR-Vs

Multiple mechanisms may account for elevated expression of AR-Vs in CRPC. Complex CRPC-specific genomic alterations involving the *AR* locus have been characterized by genome-wide copy number analysis [56]. However, the relationship between these alterations and the genesis of AR-V is presently not fully characterized. In the CWR22Rv1 cell line [48] as well as the LuCaP86.2 xenograft [50], unusually high levels of the constitutively active AR-V7 and AR<sup>V567es</sup> have been linked to genomic alterations including duplication and deletion of segments of the *AR* DNA. DNA rearrangements have been modeled using genome engineering [66]. A recent study has also shown that ADT can induce AR gene transcription and recruitment of RNA splicing factors at the specific 3′ splice site, which together contribute to the formation of AR-V7 [4, 55]. However, these specific alterations are yet to be demonstrated in clinical CRPC specimens. Alternatively, generation of AR-Vs may be coupled to the transcriptional output from AR locus, possibly regulated by negative feedback mediated by AR-FL [15]. Indeed, in CRPC specimens as well as in cell lines with induced AR-V7, AR-V7 expression is often (but not always) associated with elevated AR-FL levels, though the AR-V7/AR-FL ratios trend up in CRPC specimens and cells with induced AR-V7 [38, 94, 99]. These studies also suggest that the adaptive shift to AR-V expression is regulated at the mRNA level and that the magnitude of AR-V mRNA change is greater than that of AR-FL. Therefore, in the setting of AR-directed therapies targeting the AR-LBD, AR-V expression is associated with that of AR-FL but may not strictly parallel to that of AR-FL, and AR-V levels may be negatively regulated by AR-FL signaling.

#### Clinical and Therapeutic Implications of the AR-V Concept

Investigations encompassing the full spectrum of molecular characterization and clinical validation of the AR-V concept are still at a nascent stage, requiring carefully designed, multi-pronged approaches to dissect the biological and clinical significance of AR-Vs in the context of the coexisting (and often more abundant) AR-FL as well as other competing

mechanisms of resistance in the setting of men treated with abiraterone and enzalutamide. In the *in vitro* setting, AR-Vs have been implicated in resistance to enzalutamide and abiraterone [49, 60, 40, 16, 63]. This concept, even in the absence of clinical validation, is beginning to fuel the development of novel agents that target all AR molecules to overcome resistance [1, 71, 78, 51, 47, 96, 64, 11, 33, 52, 53, 61, 63] (also, see most recent reviews on this topic [29, 93]). We envision that conceptual advances in the field will provide the sustained impetus to drive and guide clinical development of novel agents designed to overcome resistance to AR-LBD-targeting agents, much like the compelling and rational evidence establishing the functional mechanism of sustained AR signaling in CRPC that marked the successful clinical development of abiraterone and enzalutamide. Development of methodologies for detecting these variants in the relevant patient population has significant implications for the future of drug development and clinical decision-making.

#### AR-Independent Mechanisms

As PCa is being aggressively managed by different androgen deprivation strategies, it is important to note the effects of therapy to the disease itself. As novel, more potent strategies are developed to target the AR, we could anticipate the rise of tumors that show new mechanisms of resistance, or even become AR-independent [100, 65]. Neuroendocrine prostate cancer (NEPC) is an aggressive manifestation of PCa that arises due to neuroendocrine differentiation of prostate adenocarcinoma [68, 28]. It is associated with very poor prognosis, characterized by low PSA (despite high disease burden) and the development of nonosseous (visceral) metastases [62]. The median survival is only 6–12 months [2]. NEPC does not express AR and is thus unresponsive to AR-directed therapies [9], but does show transient responses to chemotherapy [91]. The development of NEPC correlates with progression of disease [10], as well as the use of ADT [35]. A study has shown that NEPC cells express interleukin-8, a molecule with demonstrated involvement in androgen-independent growth [41]. While NEPC is only found in about 5–10 % of CRPC patients [2], its emergence after androgen deprivation strategies is crucial to consider in the clinical setting.

#### Conclusions

With multiple emerging therapies for CRPC, the treatment paradigm for PCa is ever evolving. The new range of treatment options has also given rise to the need to assess mechanisms of resistance to these therapies; an understanding of their limitations will allow for better use of these options. The

various resistance mechanisms suggest molecular heterogeneity and individual differences among patients, both of which need to be accounted for in the clinical setting. It is evident that the availability of drugs like enzalutamide and abiraterone has facilitated studies aimed at understanding the general mechanism of sustained AR signaling under castrate conditions. Future biomarker-driven clinical trials may help to achieve the dual goal of drug development and overcoming drug resistance mechanisms. It is worth noting that the successful clinical development of enzalutamide, a potent AR antagonist designed to target the LBD (a target which is missing in AR-Vs), is a direct result of laboratory mechanistic studies establishing AR protein overexpression as the key molecular determinant of castration resistance [21]. Along these lines, the recent appreciation that AR-Vs may drive resistance to enzalutamide (and abiraterone) in a subset of patients may fuel a new wave of AR-pathway-directed drugs focusing on inhibiting or eliminating AR splice variants or their transcriptional programs. Further understanding of additional molecular mechanisms of resistance should greatly facilitate drug development for CRPC moving forward.

## References

- Andersen RJ, Mawji NR, Wang J et al (2010) Regression of castrate-recurrent prostate cancer by a small-molecule inhibitor of the amino-terminus domain of the androgen receptor. *Cancer Cell* 17(6):535–546
- Aparicio A, Tzelepi V, Araujo JC et al (2011) Neuroendocrine prostate cancer xenografts with large-cell and small-cell features derived from a single patient's tumor: Morphological, immunohistochemical, and gene expression profiles. *Prostate* 71(8):846–856
- Arora VK, Schenkein E, Murali R et al (2013) Glucocorticoid receptor confers resistance to antiandrogens by bypassing androgen receptor blockade. *Cell* 155(6):1309–1322
- Aryee MJ, Liu W, Engelmann JC et al (2013) DNA methylation alterations exhibit intraindividual stability and interindividual heterogeneity in prostate cancer metastases. *Sci Transl Med* 5(169):169ra110
- Attard G, Reid AH, Yap TA et al (2008) Phase I clinical trial of a selective inhibitor of CYP17, abiraterone acetate, confirms that castration-resistant prostate cancer commonly remains hormone driven. *J Clin Oncol* 26(28):4563–4571
- Balbas MD, Evans MJ, Hosfield DJ et al (2013) Overcoming mutation-based resistance to antiandrogens with rational drug design. *Elife* 2:e00499
- Beer TM, Armstrong AJ, Rathkopf DE, et al (2014) Enzalutamide in metastatic prostate cancer before chemotherapy. *N Engl J Med*
- Beltran H, Beer TM, Carducci MA et al (2011) New therapies for castration-resistant prostate cancer: efficacy and safety. *Eur Urol* 60(2):279–290
- Beltran H, Tagawa ST, Park K et al (2012) Challenges in recognizing treatment-related neuroendocrine prostate cancer. *J Clin Oncol* 30(36):e386–e389
- Berruti A, Mosca A, Porpiglia F et al (2007) Chromogranin A expression in patients with hormone naive prostate cancer predicts the development of hormone refractory disease. *J Urol* 178(3 Pt 1): 838–843, quiz 1129
- Bohrer LR, Liu P, Zhong J et al (2013) FOXO1 binds to the TAU5 motif and inhibits constitutively active androgen receptor splice variants. *Prostate* 73(10):1017–1027
- Bolton EC, So AY, Chaivorapol C et al (2007) Cell- and gene-specific regulation of primary target genes by the androgen receptor. *Genes Dev* 21(16):2005–2017
- Bonkhoff H, Fixemer T, Hunsicker I et al (2001) Progesterone receptor expression in human prostate cancer: correlation with tumor progression. *Prostate* 48(4):285–291
- Bubendorf L, Kononen J, Koivisto P et al (1999) Survey of gene amplifications during prostate cancer progression by high-throughout fluorescence in situ hybridization on tissue microarrays. *Cancer Res* 59(4):803–806
- Cai C, He HH, Chen S, et al. Androgen receptor gene expression in prostate cancer is directly suppressed by the androgen receptor through recruitment of lysine-specific demethylase 1. *Cancer Cell* 20(4):457–471
- Cao B, Qi Y, Zhang G et al (2014) Androgen receptor splice variants activating the full-length receptor in mediating resistance to androgen-directed therapy. *Oncotarget* 5(6):1646–1656
- Carver BS, Chapinski C, Wongvipat J et al (2011) Reciprocal feedback regulation of PI3K and androgen receptor signaling in PTEN-deficient prostate cancer. *Cancer Cell* 19(5):575–586
- Chang K-H, Li R, Kuri B et al (2013) A gain-of-function mutation in DHT synthesis in castration-resistant prostate Cancer. *Cell* 154(5):1074–1084
- Chang K-H, Li R, Papari-Zareei M et al (2011) Dihydrotestosterone synthesis bypasses testosterone to drive castration-resistant prostate cancer. *Proc Natl Acad Sci* 108(33):13728–13733
- Chen CD, Welsbie DS, Tran C et al (2004) Molecular determinants of resistance to antiandrogen therapy. *Nat Med* 10(1):33–39
- Chen Y, Sawyers CL, Scher HI (2008) Targeting the androgen receptor pathway in prostate cancer. *Curr Opin Pharmacol* 8(4): 440–448
- Clegg NJ, Wongvipat J, Tran C, et al (2012) ARN-509: a novel anti-androgen for prostate cancer treatment. *Cancer Res*
- Culig Z, Hobisch A, Cronauer MV et al (1994) Androgen receptor activation in prostatic tumor cell lines by insulin-like growth factor-I, keratinocyte growth factor, and epidermal growth factor. *Cancer Res* 54(20):5474–5478
- de Bono JS, Logothetis CJ, Molina A et al (2011) Abiraterone and increased survival in metastatic prostate cancer. *N Engl J Med* 364(21):1995–2005
- Dehm SM, Tindall DJ (2011) Alternatively spliced androgen receptor variants. *Endocr Relat Cancer* 18(5):R183–R196
- Dehm SM, Schmidt LJ, Heemers HV et al (2008) Splicing of a novel androgen receptor exon generates a constitutively active androgen receptor that mediates prostate cancer therapy resistance. *Cancer Res* 68(13):5469–5477
- Di Lorenzo G, Buonerba C, Autorino R et al (2010) Castration-resistant prostate cancer. *Drugs* 70(8):983–1000
- Epstein JI, Amin MB, Beltran H et al (2014) Proposed morphologic classification of prostate cancer with neuroendocrine differentiation. *Am J Surg Pathol* 38(6):756–767
- Ferraldeschi R, Welti J, Luo J, et al (2014) Targeting the androgen receptor pathway in castration-resistant prostate cancer: progresses and prospects. *Oncogene*
- Gregory CW, He B, Johnson RT et al (2001) A mechanism for androgen receptor-mediated prostate cancer recurrence after androgen deprivation therapy. *Cancer Res* 61(11):4315–4319
- Guo Z, Yang X, Sun F et al (2009) A novel androgen receptor splice variant is up-regulated during prostate cancer progression and promotes androgen depletion-resistant growth. *Cancer Res* 69(6): 2305–2313

32. Hagan CR, Lange CA (2014) Molecular determinants of context-dependent progesterone receptor action in breast cancer. *BMC Med* 12:32
33. He S, Zhang C, Shafi AA et al (2013) Potent activity of the Hsp90 inhibitor ganetespib in prostate cancer cells irrespective of androgen receptor status or variant receptor expression. *Int J Oncol* 42(1):35–43
34. Higano CS, Crawford ED (2011) New and emerging agents for the treatment of castration-resistant prostate cancer. *Urol Oncol Semin Orig Investig* 29(6, Supplement):1–8
35. Hirano D, Okada Y, Minei S et al (2004) Neuroendocrine differentiation in hormone refractory prostate cancer following androgen deprivation therapy. *Eur Urol* 45(5):586–592, discussion 592
36. Hörnberg E, Ylitalo EB, Crnalic S et al (2011) Expression of androgen receptor splice variants in prostate cancer bone metastases is associated with castration-resistance and short survival. *PLoS ONE* 6(4):e19059
37. Hu R, Denmeade SR, Luo J (2010) Molecular processes leading to aberrant androgen receptor signaling and castration resistance in prostate cancer. *Expert Rev Endocrinol Metab* 5(5):753–764
38. Hu R, Dunn TA, Wei S et al (2009) Ligand-independent androgen receptor variants derived from splicing of cryptic exons signify hormone-refractory prostate cancer. *Cancer Res* 69(1):16–22
39. Hu R, Isaacs WB, Luo J (2011) A snapshot of the expression signature of androgen receptor splicing variants and their distinctive transcriptional activities. *Prostate* 71(15):1656–1667
40. Hu R, Lu C, Mostaghel EA, et al (2012) Distinct transcriptional programs mediated by the ligand-dependent full-length androgen receptor and its splice variants in castration-resistant prostate cancer. *Cancer Res* accepted
41. Huang J, Yao JL, Zhang L et al (2005) Differential expression of interleukin-8 and its receptors in the neuroendocrine and non-neuroendocrine compartments of prostate cancer. *Am J Pathol* 166(6):1807–1815
42. Huggins C, Hodges CV (2002) Studies on prostatic cancer. I. The effect of castration, of estrogen and of androgen injection on serum phosphatases in metastatic carcinoma of the prostate. 1941. *J Urol* 167(2 Pt 2):948–951, discussion 952
43. Jemal A, Bray F, Center MM et al (2011) Global cancer statistics. *CA Cancer J Clin* 61(2):69–90
44. Joseph JD, Lu N, Jing Q, et al (2013) A clinically relevant androgen receptor mutation confers resistance to 2nd generation anti-androgens enzalutamide and ARN-509. *Cancer Discovery*
45. Knudsen KE, Scher HI (2009) Starving the addiction: new opportunities for durable suppression of AR signaling in prostate cancer. *Clin Cancer Res* 15(15):4792–4798
46. Korpala M, Korn JM, Gao X et al (2013) An F876L mutation in androgen receptor confers genetic and phenotypic resistance to MDV3100 (enzalutamide). *Cancer Discov* 3(9):1030–1043
47. Levine PM, Imberg K, Garabedian MJ et al (2012) Multivalent peptidomimetic conjugates: a versatile platform for modulating androgen receptor activity. *J Am Chem Soc* 134(16):6912–6915
48. Li Y, Alsagabi M, Fan D et al (2011) Intragenic rearrangement and altered RNA splicing of the androgen receptor in a cell-based model of prostate cancer progression. *Cancer Res* 71(6):2108–2117
49. Li Y, Chan SC, Brand LJ et al (2013) Androgen receptor splice variants mediate enzalutamide resistance in castration-resistant prostate cancer cell lines. *Cancer Res* 73(2):483–489
50. Li Y, Hwang TH, Oseth L, et al (2012) AR intragenic deletions linked to androgen receptor splice variant expression and activity in models of prostate cancer progression. *Oncogene In Press*
51. Lim AC, Attard G (2013) Improved therapeutic targeting of the androgen receptor: rational drug design improves survival in castration-resistant prostate cancer. *Curr Drug Targets* 14(4):408–419
52. Lin TH, Izumi K, Lee SO et al (2013) Anti-androgen receptor ASC-J9 versus anti-androgens MDV3100 (enzalutamide) or Casodex (bicalutamide) leads to opposite effects on prostate cancer metastasis via differential modulation of macrophage infiltration and STAT3-CCL2 signaling. *Cell Death Dis* 4:e764
53. Liu C, Lou W, Zhu Y et al (2014) Niclosamide inhibits androgen receptor variants expression and overcomes enzalutamide resistance in castration-resistant prostate cancer. *Clin Cancer Res* 20(12):3198–3210
54. Liu G, Cynthia S, Shihua S et al (2013) AR variant ARv567es induces carcinogenesis in a novel transgenic mouse model of prostate cancer. *Neoplasia* 15(9):1009, New York, NY
55. Liu LL, Xie N, Sun S, et al (2013) Mechanisms of the androgen receptor splicing in prostate cancer cells. *Oncogene*
56. Liu W, Laitinen S, Khan S et al (2009) Copy number analysis indicates monoclonal origin of lethal metastatic prostate cancer. *Nat Med* 15(5):559–565
57. Locke JA, Guns ES, Lubik AA et al (2008) Androgen levels increase by intratumoral de novo steroidogenesis during progression of castration-resistant prostate cancer. *Cancer Res* 68(15):6407–6415
58. Merson S, Yang ZH, Brewer D et al (2014) Focal amplification of the androgen receptor gene in hormone-naive human prostate cancer. *Br J Cancer* 110(6):1655–1662
59. Mohler JL, Gregory CW, Ford OH 3rd et al (2004) The androgen axis in recurrent prostate cancer. *Clin Cancer Res* 10(2):440–448
60. Mostaghel EA, Marck BT, Plymate SR et al (2011) Resistance to CYP17A1 inhibition with abiraterone in castration-resistant prostate cancer: induction of steroidogenesis and androgen receptor splice variants. *Clin Cancer Res* 17(18):5913–5925
61. Myung JK, Banuelos CA, Fernandez JG et al (2013) An androgen receptor N-terminal domain antagonist for treating prostate cancer. *J Clin Invest* 123(7):2948–2960
62. Nadal R, Schweizer M, Kryvenko ON et al (2014) Small cell carcinoma of the prostate. *Nat Rev Urol* 11(4):213–219
63. Nadiminty N, Tummala R, Liu C et al (2013) NF-kappaB2/p52 induces resistance to enzalutamide in prostate cancer: role of androgen receptor and its variants. *Mol Cancer Ther* 12(8):1629–1637
64. Nakka M, Agoulnik IU, Weigel NL (2013) Targeted disruption of the p160 coactivator interface of androgen receptor (AR) selectively inhibits AR activity in both androgen-dependent and castration-resistant AR-expressing prostate cancer cells. *Int J Biochem Cell Biol* 45(4):763–772
65. Nelson PS (2012) Molecular states underlying androgen receptor activation: a framework for therapeutics targeting androgen signaling in prostate cancer. *J Clin Oncol* 30(6):644–646
66. Nyquist MD, Li Y, Hwang TH et al (2013) TALEN-engineered AR gene rearrangements reveal endocrine uncoupling of androgen receptor in prostate cancer. *Proc Natl Acad Sci U S A* 110(43):17492–17497
67. Palmos PL, Hussain M (2013) Non-castrate metastatic prostate cancer: have the treatment options changed? *Semin Oncol* 40(3):337–346
68. Palmgren JS, Karavadia SS, Wakefield MR (2007) Unusual and underappreciated: small cell carcinoma of the prostate. *Semin Oncol* 34(1):22–29
69. Plymate SR, Luo J (2013) The expression signature of androgen receptor splice variants and their distinctive transcriptional activities in castration-resistant prostate cancer. In *Androgen-Responsive Genes in Prostate Cancer*, 201–213. Springer
70. Ryan CJ, Smith MR, de Bono JS et al (2013) Abiraterone in metastatic prostate cancer without previous chemotherapy. *N Engl J Med* 368(2):138–148
71. Sadar MD (2011) Small molecule inhibitors targeting the “Achilles’ heel” of androgen receptor activity. *Cancer Res* 71(4):1208–1213



72. Sahu B, Laakso M, Pihlajamaa P et al (2013) FoxA1 specifies unique androgen and glucocorticoid receptor binding events in prostate cancer cells. *Cancer Res* 73(5):1570–1580
73. Sarker D, Reid AHM, Yap TA et al (2009) Targeting the PI3K/AKT Pathway for the treatment of prostate cancer. *Clin Cancer Res* 15(15):4799–4805
74. Scher HI, Beer TM, Higano CS et al (2010) Antitumour activity of MDV3100 in castration-resistant prostate cancer: a phase 1–2 study. *Lancet* 375(9724):1437–1446
75. Scher HI, Buchanan G, Gerald W et al (2004) Targeting the androgen receptor: improving outcomes for castration-resistant prostate cancer. *Endocr Relat Cancer* 11(3):459–476
76. Scher HI, Sawyers CL (2005) Biology of progressive, castration-resistant prostate cancer: directed therapies targeting the androgen-receptor signaling axis. *J Clin Oncol* 23(32):8253–8261
77. Scher HI, Fizazi K, Saad F et al (2012) Increased survival with enzalutamide in prostate cancer after chemotherapy. *N Engl J Med* 367(13):1187–1197
78. Schmidt LJ, Tindall DJ (2013) Androgen receptor: past, present and future. *Curr Drug Targets* 14(4):401–407
79. Schweizer MT, Antonarakis ES (2012) Abiraterone and other novel androgen-directed strategies for the treatment of prostate cancer: a new era of hormonal therapies is born. *Ther Adv Urol* 4(4):167–178
80. Shafi AA, Yen AE, Weigel NL (2013) Androgen receptors in hormone-dependent and castration-resistant prostate cancer. *Pharmacol Ther* 140(3):223–238
81. Sharifi N, Gulley JL, Dahut WL (2005) Androgen deprivation therapy for prostate cancer. *JAMA* 294(2):238–244
82. Steinkamp MP, O'Mahony OA, Brogley M et al (2009) Treatment-dependent androgen receptor mutations in prostate cancer exploit multiple mechanisms to evade therapy. *Cancer Res* 69(10):4434–4442
83. Sun F, Chen H-g, Li W et al (2014) Androgen receptor splice variant AR3 promotes prostate cancer via modulating expression of autocrine/paracrine factors. *J Biol Chem* 289(3):1529–1539
84. Sun S, Sprenger CC, Vessella RL et al (2010) Castration resistance in human prostate cancer is conferred by a frequently occurring androgen receptor splice variant. *J Clin Invest* 120(8):2715–2730
85. Szmulewitz RZ, Chung E, Al-Ahmadie H et al (2012) Serum/glucocorticoid-regulated kinase 1 expression in primary human prostate cancers. *Prostate* 72(2):157–164
86. Taplin ME, Bubley GJ, Shuster TD et al (1995) Mutation of the androgen-receptor gene in metastatic androgen-independent prostate cancer. *N Engl J Med* 332(21):1393–1398
87. Taplin M-E, Montgomery RB, Logothetis C, et al (2012) Effect of neoadjuvant abiraterone acetate (AA) plus leuprolide acetate (LHRHa) on PSA, pathological complete response (pCR), and near pCR in localized high-risk prostate cancer (LHRPC): results of a randomized phase II study. In ASCO Meeting Abstracts
88. Titus MA, Schell MJ, Lih FB et al (2005) Testosterone and dihydrotestosterone tissue levels in recurrent prostate cancer. *Clin Cancer Res* 11(13):4653–4657
89. Tombal B, Borre M, Rathenborg P, et al (2014) Enzalutamide monotherapy in hormone-naive prostate cancer: primary analysis of an open-label, single-arm, phase 2 study. *Lancet Oncol*
90. Tran C, Ouk S, Clegg NJ et al (2009) Development of a second-generation antiandrogen for treatment of advanced prostate cancer. *Science* 324(5928):787–790
91. Tzelepi V, Zhang J, Jing-Fang L et al (2012) Modeling a lethal prostate cancer variant with small-cell carcinoma features. *Clin Cancer Res* 18(3):666–677
92. Visakorpi T, Hyytinen E, Koivisto P et al (1995) In vivo amplification of the androgen receptor gene and progression of human prostate cancer. *Nat Genet* 9(4):401–406
93. Ware KE, Garcia-Blanco MA, Armstrong AJ, et al (2014) Significance of androgen receptor variants in castration resistant prostate cancer. *Endocr Relat Cancer*
94. Watson PA, Chen YF, Balbas MD et al (2010) Constitutively active androgen receptor splice variants expressed in castration-resistant prostate cancer require full-length androgen receptor. *Proc Natl Acad Sci U S A* 107(39):16759–16765
95. Watson PA, Chen YF, Balbas MD, et al (2010) Constitutively active androgen receptor splice variants expressed in castration-resistant prostate cancer require full-length androgen receptor. *Proc Natl Acad Sci*
96. Yamashita S, Lai KP, Chuang KL et al (2012) ASC-J9 suppresses castration-resistant prostate cancer growth through degradation of full-length and splice variant androgen receptors. *Neoplasia* 14(1):74–83
97. Yeh S, Miyamoto H, Shima H et al (1998) From estrogen to androgen receptor: a new pathway for sex hormones in prostate. *Proc Natl Acad Sci U S A* 95(10):5527–5532
98. Yu Y, Liu L, Xie N et al (2013) Expression and function of the progesterone receptor in human prostate stroma provide novel insights to cell proliferation control. *J Clin Endocrinol Metab* 98(7):2887–2896
99. Yu Z, Chen S, Sowalsky AG, et al (2014) Rapid induction of androgen receptor splice variants by androgen deprivation in prostate cancer. *Clin Cancer Res: clincanres*. 1863.2013
100. Yuan X, Cai C, Chen S, et al (2013) Androgen receptor functions in castration-resistant prostate cancer and mechanisms of resistance to new agents targeting the androgen axis. *Oncogene*
101. Zhang X, Morrissey C, Sun S et al (2011) Androgen receptor variants occur frequently in castration resistant prostate cancer metastases. *PLoS ONE* 6(11):e27970
102. Zhao X-Y, Malloy PJ, Krishnan AV et al (2000) Glucocorticoids can promote androgen-independent growth of prostate cancer cells through a mutated androgen receptor. *Nat Med* 6(6):703–706
103. Zhu H, Garcia JA (2013) Targeting the adrenal gland in castration-resistant prostate cancer: a case for orteronel, a selective CYP-17 17, 20-lyase inhibitor. *Curr Oncol Rep* 15(2):105–112