Correction to: Late lumen enlargement after drug-coated balloon angioplasty for denovo coronary artery disease

Takayuki Onishi¹ · Yuko Onishi¹ · Isshi Kobayashi¹ · Yasuhiro Sato¹

Published online: 17 September 2020

© Japanese Association of Cardiovascular Intervention and Therapeutics 2020

Correction to:

Cardiovascular Intervention and Therapeutics (2020)

<https://doi.org/10.1007/s12928-020-00690-2>

In the original publication of the article, Table 3 was published incorrectly. The correct Table 3 is given in this Correction [1].

The original article has been corrected.

Reference

1. Onishi T, Onishi Y, Kobayashi I, Sato Y. Late lumen enlargement after drug-coated balloon angioplasty for denovo coronary artery disease. Cardiovascular Intervention and Therapeutics. 2020. <https://doi.org/10.1007/s12928-020-00690-2>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s12928-020-00690-2>.

✉ Takayuki Onishi
takayuki.onishi@yahoo.com

¹ Department of Cardiology, Hiratsuka Kyosai Hospital, 9-11 Oiwake, Hiratsuka, Kanagawa, Japan

Table 3 Results from angiography

	Total		LLE		No LLE		<i>p</i>	
Lesion, <i>n</i>	196		109		87			
Preprocedure								
RVD, mm	2.07	±0.66	1.96	±0.55	2.20	±0.76	0.090	
MLD, mm	0.86	±0.44	0.82	±0.44	0.91	±0.44	0.212	
DS, %	58.45	±19.31	58.48	±20.05	58.42	±18.46	0.408	
Lesion length, mm	11.23	±6.07	11.03	±5.80	11.46	±6.39	0.721	
Postprocedure								
RVD, mm	2.16	±0.57	2.03	±0.49	2.32	±0.64	0.002	
MLD, mm	1.49	±0.42	1.37	±0.34	1.64	±0.47	<0.001	
DS, %	29.75	±14.59	31.18	±14.04	27.96	±15.14	0.065	
Acute lumen gain, mm	0.64	±0.45	0.56	±0.43	0.74	±0.45	0.001	
TIMI flow grade								
3	189	(96%)	106	(97%)	83	(95%)	0.476	
2	5	(3%)	3	(3%)	2	(2%)		
1	2	(1%)	0	(0%)	2	(2%)		
Residual dissection								
NHLBI classification								
A	28	(14%)	16	(15%)	12	(14%)	0.099	
B	24	(12%)	18	(17%)	6	(7%)		
C	15	(8%)	9	(8%)	6	(7%)		
D	3	(2%)	1	(1%)	2	(2%)		
E	1	(1%)	1	(1%)	0	(0%)		
F	0	(0%)	0	(0%)	0	(0%)		
Angiographic follow-up								
Duration, months	7.2	±2.5	7.3	±2.6	7.2	±2.3	0.908	
RVD, mm	2.20	±0.65	2.19	±0.50	2.23	±0.82	0.110	
MLD, mm	1.46	±0.48	1.63	±0.37	1.25	±0.51	<0.001	
DS, %	31.97	±19.10	24.64	±11.36	41.15	±22.62	<0.001	
Late lumen loss, mm	0.03	±0.43	−0.26	±0.20	0.39	±0.37	<0.001	
Binary restenosis	42	(21%)	6	(6%)	36	(41%)	<0.001	
Binary restenosis according to the ACC/AHA lesion type								
A	(<i>n</i> =22)	1	(5%)	0	(0%)	1	(17%)	0.273
B1	(<i>n</i> =59)	10	(17%)	2	(5%)	8	(38%)	0.002
B2	(<i>n</i> =80)	19	(24%)	3	(7%)	16	(41%)	<0.001
C	(<i>n</i> =35)	12	(34%)	1	(7%)	11	(52%)	0.006

Data are presented as numbers (%) or means ± standard deviations

ACC American College of Cardiology, AHA American Heart Association, DS diameter stenosis, LLE late lumen enlargement, NHLBI National Heart, Lung, and Blood Institute, MLD minimal lumen diameter, RVD reference vessel diameter, TIMI thrombolysis in myocardial infarction