Correction to: Intracerebral Hemorrhage–Induced Brain Injury in Rats: the Role of Extracellular Peroxiredoxin 2

Liheng Bian^{1,2} · Jingwei Zhang¹ · Ming Wang¹ · Richard F. Keep¹ · Guohua Xi¹ · Ya Hua¹

Published online: 9 August 2019

© Springer Science+Business Media, LLC, part of Springer Nature 2019

Correction to: Transl. Stroke Res.

<https://doi.org/10.1007/s12975-019-00714-x>

In Fig. 1, molecular weight of peroxiredoxin 2 (PRX-2) is 22 kDa rather than 32 kDa.

The online version of the original article can be found at <https://doi.org/10.1007/s12975-019-00714-x>

✉ Ya Hua
yahua@umich.edu

¹ Department of Neurosurgery, University of Michigan, R5018 BSRB, 109 Zina Pitcher Place, Ann Arbor, MI 48109-2200, USA

² Department of Neurology, Beijing Tiantan Hospital, Capital Medical University, Beijing, China

Fig. 1 Western blot assessing PRX-2 levels in the ipsilateral basal ganglia at 1 h after lysed RBCs or saline injection (15 μ l). Values are means $\pm$ SD; $n = 5$ for both groups; # $P < 0.01$ vs. saline group by the Student t test

The authors regret the errors.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.