CORRECTION

Correction to: Spatiotemporal Analysis of Tropical Cyclone Landfalls in Northern Bay of Bengal, India and Bangladesh

Sunando Bandyopadhyay¹ · Susmita Dasgupta² · Zahirul Huque Khan³ · David Wheeler⁴

Published online: 28 May 2021

© Korean Meteorological Society and Springer Nature B.V. 2021

Correction to: Asia-Pacific Journal of Atmospheric Sciences.

<https://doi.org/10.1007/s13143-021-00227-4>

Due to an unfortunate oversight during the correction process the images have not been updated. They should be:

The online version of the original article can be found at <https://doi.org/10.1007/s13143-021-00227-4>

✉ Sunando Bandyopadhyay
sunando@live.com

Susmita Dasgupta
sdasgupta@worldbank.org

Zahirul Huque Khan
zhk@iwmbd.org

David Wheeler
wheelrdr@gmail.com

¹ Department of Geography, University of Calcutta, Kolkata 700019, India

² Development Research Group, World Bank, Washington, DC 20433, USA

³ Institute of Water Modelling, Dhaka 1206, Bangladesh

⁴ World Resources Institute, Washington, DC 20002, USA

Figure 1 Cyclonic storms (wind speed ≥ 34 kn or 62 km/h) occurring in (a) the Northern Indian Ocean basin and landfalling in (b) the Odisha–West Bengal–Bangladesh (Northern Bay of Bengal) coast during 1877–2016. The coastline used for storm landfall analysis is shown in orange. Storm track data from: IBTrACS, IMD

Figure 2 Storm occurrence / landfall and maximum wind speed characteristics of the cyclones occurring in the hierarchically smaller basins of Northern Indian Ocean (**a–c**), Bay of Bengal (**d–f**), and Northern Bay of Bengal, comprising Odisha, West Bengal and Bangladesh coasts (**g–i**),

during 1877–2016. Periods (1–7) are defined in lower axes annotations of the left panel charts. Note region-wise differences in vertical scales in the figures of the left and central panels. Data source: IBTrACS, IMD

Figure 3 Regional and seasonal variation in occurrence of cyclonic storms in Northern Indian Ocean (NIO), Bay of Bengal (BoB), and landfall in Northern Bay of Bengal (NBoB) during 1877–2016. (a–

g) show the percentage of all storm crossings for each of the seven time periods; (h) considers storms of the entire span of study. Data source: IBTrACS, IMD

Figure 4 Storm landfalls in the Northern Bay of Bengal coast: Median locations of 168 storms by period, composite for 1877–2016. Data source: IBTrACS, IMD

Figure 5 Storm landfalls in the Northern Bay of Bengal coast: Distribution by seven periods between 1877 and 2016. Q1 and Q3 stand for first quartile and third quartile, respectively. Data Source: IBTrACS, IMD

Figure 6 Cyclone landfall intensities (strikes/year) in Northern Bay of Bengal coasts: shown by periods (a–g). Data source: IBTrACS, IMD

Figure 7 Cyclone landfalls in the Northern Bay of Bengal coast, composite for 1877–2016. Data source: IBTrACS, IMD

Figure 8 Mean maximum wind speed of cyclones landfalling in the Northern Bay of Bengal coast, shown by periods (a–g), 1877–2016. Data Source: IBTrACS, IMD

Figure 9 Mean maximum wind speeds of cyclones landfalling in the Northern Bay of Bengal coast, composite for 1877–2016. Data source: IBTrACS, IMD

Figure 10 Cyclone impact quintiles, Northern Bay of Bengal coast, shown by periods (a–g). Data Source: IBTrACS, IMD

Figure 11 Cyclone impact percentiles, Northern Bay of Bengal coast, composite for 1877–2016. Data source: IBTrACS, IMD

The original article has been corrected.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.