



## Correction to: Indigenous siliciclastic and extraneous polygenetic carbonate facies in the Albian-Turonian Karai Shale, Cauvery Basin, India

Nivedita Chakraborty<sup>1</sup> · Anudeb Mandal<sup>2</sup> · Adrita Choudhuri<sup>3</sup> · Sunipa Mandal<sup>3</sup> · Subir Sarkar<sup>3</sup>

Published online: 5 March 2018  
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

**Correction to: Carbonates and Evaporites**  
<https://doi.org/10.1007/s13146-018-0419-0>

Unfortunately, Fig. 1 was incorrectly published in the original publication and the correct version is updated here. The original article has been corrected.

---

The original article can be found online at <https://doi.org/10.1007/s13146-018-0419-0>.


---

✉ Nivedita Chakraborty  
nivedita.jugeo@gmail.com

<sup>1</sup> Department of Geology, K.J.R Govt. General Degree College, Bankura 722143, India

<sup>2</sup> Department of Geology, Presidency University, Kolkata 700073, India

<sup>3</sup> Department of Geological Sciences, Jadavpur University, Kolkata 700032, India



**Fig. 1** Lithological and location map of the studied area (number showing studied locations of Karai Shale), (map of India within inset) (modified after Sarkar et al. 2014) (a). Stratigraphic successions of the area on the right-hand side (Ba. Barremian; Cen. Cenomanian; Co. Coniacian; San. Santonian; Fm. Formation) (b). Spatial and

temporal distribution of the Karai Shale in different studied localities. Note at location no. 7 the Karai Shale directly overlies basement rocks (c). The structural definition of Cauvery basin (outline of the basin including onshore and offshore within inset) (d)