



Monitoring Impact of Salt-Marsh Vegetation Characteristics on Sedimentation: an Outlook for Nature-Based Flood Protection

B. Martina Baaij^{1,2}  · Jeroen Kooijman² · Juul Limpens¹  · Richard J. C. Marijnissen²  · Jantsje M. van Loon-Steensma² 

Received: 10 December 2020 / Accepted: 27 May 2021 / Published online: 1 July 2021
© The Author(s) 2021

Abstract

Salt marshes can protect coastlines against flooding by attenuating wave energy and enhancing shoreline stabilization. However, salt-marsh functioning is threatened by human influences and sea level rise. Although it is known that protection services are mediated by vegetation, little is known about the role of vegetation structure in salt-marsh accretion. We investigated the role of vegetation presence, vegetation type and structural vegetation characteristics in sedimentation and sediment grain size. We established 56 plots on a salt marsh on the Dutch Wadden island of Texel. Plots were divided over four vegetation types contrasting in vegetation structure and varied in elevation and distance to creeks. Vegetation presence was controlled by clipping in subplots. Within each plot, we measured seven vegetation characteristics, sedimentation and the sediment grain size distribution. Furthermore, we explored the effect of the natural variation in vegetation structure on wave attenuation with a simple model approach. For this, we developed vegetation scenarios based on the field measurements of stem height, diameter and density. We found that vegetation presence increased sedimentation on average by 42%. Sedimentation was highest in *Salicornia* vegetation and increased with stem height and branching level. Grain size also seemed to increase with branching level. Modelled wave attenuation was 7.5 times higher with natural vegetation compared to topography only, was strongest for *Spartina* vegetation and most sensitive to the natural variance in stem density. Our results can be used to improve predictions of salt-marsh accretion and the implementation of salt marshes in nature-based flood defences.

Keywords Vegetation type · Vegetation structure · Biomass · Sediment deposition · Wave damping · Grain size

Introduction

Salt marshes provide important ecosystem services, such as protection against coastal erosion and flooding (Gedan et al. 2009; Barbier et al. 2011). However, salt marshes are threatened worldwide by changes in land use and climate, leading to salt-marsh degradation and the loss of ecosystem services (Adam 2002; Gedan et al. 2009). Salt marshes are expected

to drown from sea level rise when landward migration is blocked by steep slopes or barriers (Torio and Chmura 2013), or when relative sea level rise exceeds accretion rates (Redfield 1965). Crosby et al. (2016) predicted that 60–91% of their studied salt marshes are threatened by sea level rise, based on the sea-level rise scenarios predicted by the Intergovernmental Panel on Climate Change (IPCC) for 2100 (Stocker et al. 2013). However, salt-marsh vulnerability to sea level rise may be exaggerated without consideration of biophysical feedbacks between plant growth and accretion (Kirwan and Megonigal 2013; Crosby et al. 2016; Kirwan et al. 2016).

The expected sea level rise and increased storm activity highlights the role of salt marshes in coastal protection (Shepard et al. 2011; Leonardi et al. 2018). A meta-analysis by Shepard et al. (2011) demonstrates positive contributions of salt-marsh vegetation to shoreline stabilization (i.e. sediment deposition, prevention of erosion and increase in elevation) and wave attenuation. Wave attenuation can be observed

✉ Juul Limpens
juul.limpens@wur.nl

✉ Jantsje M. van Loon-Steensma
jantsje.vanloon@wur.nl

¹ Plant Ecology and Nature Conservation Group, Wageningen University & Research, P.O. Box 47, Wageningen 6700 AA, The Netherlands

² Water Systems and Global Change Group, Wageningen University & Research, P.O. Box 47, Wageningen 6700 AA, The Netherlands

even during storm conditions (Möller et al. 2014; Vuik et al. 2016; Garzon et al. 2019a). Temmerman et al. (2013) therefore argue that coastal ecosystems, like salt marshes, are a preferable long-term flood protection measure over conventional hard infrastructure because of their natural adaptive capacity to sea level rise. However, the implementation of salt marshes in nature-based solutions requires an in-depth understanding of the role of vegetation in their adaptive capacity and flood protection function.

Vegetation enhances the settling of sediment and prevents erosion by reducing the flow velocity and turbulence between stems and by wave attenuation (Möller et al. 1999; Christiansen et al. 2000; Shi et al. 2012; Möller et al. 2014). Furthermore, vegetation itself can trap suspended sediment by adhesion of the sediment particles (Li and Yang 2009). Sedimentation and wave attenuation are influenced by vegetation characteristics like height, density, biomass and stiffness of the vegetation (e.g., Bouma et al. 2010; Shepard et al. 2011; Anderson and Smith 2014; Reef et al. 2018).

When studying the role of vegetation in the adaptive capacity and flood protection function of salt marshes, geomorphological factors must be taken into account, as they influence sediment deposition, wave attenuation and vegetation growth (e.g., Möller et al. 1999; Wood and Hine 2007; Townend et al. 2011). Sedimentation decreases with a larger distance to the source of sediment, such as the sea or a creek, due to the progressive settlement of sediment out of suspension (e.g., Reed et al. 1999; Wood and Hine 2007). More elevated locations experience less inundation and thus accumulate less sediment (e.g., Townend et al. 2011; Reef et al. 2018). Also the sediment grain size generally decreases with elevation and distance to the source of sediment, due to a reduction of wave action, flow velocity and turbulence (Yang 1999a; Yang et al. 2000; Yang et al. 2008; Gibeault et al. 2016). Wave energy diminishes with elevation due to wave breaking in shallow water and friction with the foreshore (e.g., Möller et al. 1999; Möller 2006; Ysebaert et al. 2011). Furthermore, elevation, inundation frequency and distance to the creek also influence plant species occurrence, resulting in vegetation zonation (Adam 2002; Townend et al. 2011).

Studies investigating the impact of vegetation on sedimentation or wave attenuation have focussed on a limited number of plant species (e.g., often *Spartina* species) (for an overview of studies see Shepard et al. (2011) or Vuik et al. (2016)). Elevation differences between vegetated and unvegetated sites complicate studying the effect of vegetation presence (Möller et al. 1999; Shepard et al. 2011). A limited number of field studies have investigated the effect of vegetation characteristics on sediment deposition and sediment grain size (Morris et al. 2002; Yang et al. 2008; Gibeault et al. 2016; Reef et al. 2018). These studies primarily focus on the density, height or biomass of the vegetation (Shepard et al. 2011). Studies investigating the effect of vegetation characteristics on wave

attenuation mainly focus on the density, height or stiffness of the vegetation (Augustin et al. 2009; Shepard et al. 2011; Anderson and Smith 2014; Peruzzo et al. 2018). This effect is often studied in flume experiments with artificial or real vegetation (Augustin et al. 2009; Anderson and Smith 2014; Peruzzo et al. 2018). However, the relative importance of each vegetation characteristic to wave attenuation (Anderson and Smith 2014) and their variation in the field require further investigation.

In this paper, we investigated the effect of vegetation presence, vegetation type and natural variation in vegetation structure characteristics on sedimentation and sediment grain size on a salt marsh in the Netherlands. Additionally, we developed field-based vegetation scenarios to explore the potential effect of natural variations in vegetation characteristics on wave attenuation with a simple model approach. In the field, a total of 56 plots were established in four vegetation types and over a range of elevations and distances to the creek. Vegetation presence was controlled and seven vegetation characteristics (total cover, vegetation height, stem height, stem diameter, branching level, stem density and biomass), sedimentation and the sediment grain size distribution were measured. For this field experiment, we expected (1) vegetation presence to increase sedimentation, (2) to observe differences in sedimentation and sediment grain size between vegetation types and (3) effects of vegetation characteristics on sedimentation and sediment grain size. Subsequently, vegetation scenarios were developed based on the field measurements of three vegetation characteristics (stem height, stem diameter and stem density) to model wave height with the XBeach 1D model over a transect. With this established model we aimed to explore (1) how modelled wave attenuation by vegetation zonation (i.e. the field situation) compares to the absence of vegetation, (2) how modelled wave attenuation differs between the four vegetation types and (3) how modelled wave attenuation is affected by natural variance of vegetation characteristics.

Methods

Study Area

All field research was conducted on a salt marsh in nature reserve the Slufter on the barrier island of Texel, the Netherlands (53°08'18.4"N, 4°48'31.1"E) (Fig. 1a, b). The Slufter is managed by the State Forest Service and part of the Natura 2000 network, a European network of protected nature areas. The nature reserve of about 700 ha consists of a dune valley that is surrounded by two sand dikes and connected to the North Sea by a channel (Reitsma and de Jong 2019). The tides are semi-diurnal with a tidal range of 1.7 m on




Fig. 1 Study area and design. **a** Location of study area the Slufter on the barrier island of Texel, the Netherlands. **b** Aerial view of the Slufter with locations of the 56 plots used for the field experiment, arranged in four vegetation types and seven blocks, and the transect (5,4 km) used to model wave attenuation. **c** Overview of the four vegetation types. **d**

Schematic representation on the left of a plot with two subplots where vegetation is present (grey) or removed (white) and on the right of a subplot with four sediment traps (circles). Distances are indicated with thin lines

average (between -0.93 and 0.74 m + NAP; Dutch ordnance level, close to mean sea level), 1.9 m during spring tides (between -1.08 and 0.82 m + NAP) and 1.3 m during neap tides (between -0.74 and 0.57 m + NAP).

The vegetation in the dune valley of the Slufter consists of a relatively young salt marsh with only a small area of high salt marsh (Reitsma and de Jong 2019). Pioneer vegetation growing along the creeks and in low lying areas is dominated by

Salicornia species and *Suaeda maritima*, and some small patches of *Spartina anglica* are present. Vegetation of the low marsh is dominated by *Limonium vulgare* and *Atriplex portulacoides*. The medium salt marsh mainly consists of *Festuca rubra*, *Juncus gerardii*, *L. vulgare* and *Plantago maritima*. The high salt marsh is dominated by *Elytrigia atherica* (Reitsma and de Jong 2019).

Study Design of Field Experiment

The field experiment was conducted between August and October in 2018. 56 plots (2.5 × 2.5 m) were established in the Slufter according to a factorial design (Fig. 1b). Plots were made in four vegetation types contrasting in vegetation structure and elevation: vegetation dominated by either *Salicornia* species (*Salicornia* type), *S. anglica* (*Spartina* type), *A. portulacoides* (*Atriplex* type) or *L. vulgare* (*Limonium* type) (Fig. 1c; for a schematic representation of the dominating plant species, see Fig. 3a in Möller et al. 1999). Because distance to the source of sediment is known to influence sedimentation (Reed et al. 1999; Temmerman et al. 2003; Wood and Hine 2007), pairs of plots were made near a feeder creek within the same vegetation type. One of the plots within a pair was established at the shortest possible distance to the creek, while the other plot was established at the longest possible distance to the creek. The position relative to the main channel was also expected to influence sedimentation. Therefore, plots were arranged in seven experimental blocks, each block consisting of one pair of plots per vegetation type (Fig. 1b).

In each plot, two subplots (50 × 50 cm) were made 1 m apart and with the same distance to the creek (Fig. 1d). One subplot consisted of intact vegetation, whereas the vegetation in the other subplot was clipped to the soil surface to investigate the effect of vegetation presence on sedimentation. Locations with the same geomorphological factors (e.g., distance to the creek, elevation) but different vegetation could be compared in this setup. The subplots were relatively small to minimize damage to the vegetation of the nature reserve.

The location and elevation were recorded in the centre of the subplots by a Real-Time Kinematic (RTK-)GNSS receiver. The elevation of the plot was determined as the average of the two subplots. The distance to the creek was measured from the centre of the plot to the edge of the creek (transition between vegetation and unvegetated creek).

Vegetation Composition and Characteristics

The total vegetation cover as well as the cover per plant species were estimated in percentages per plot. Plant species were identified according to van der Meijden (2005). The plant communities in the plots were determined with SynBioSys (Hennekens et al. 2018). This Dutch program contains

information on plant species, plant communities and landscapes and has classification software. Vegetation height, defined as the estimated average height of the plant community, was based on measurements in one to three representative locations per plot.

Aside from total cover and vegetation height, five other vegetation characteristics were measured per plot, using the clipped plant material of a representative, smaller area (10 × 10 cm) within the subplot. Only for six plots of the *Salicornia* type, that had a total cover below 50%, plant material of the whole subplot was used. The number of stems was counted per species to calculate the stem density per square meter. Also the stem height of straightened stems, the stem diameter at around 1/3 of the stem (following van Loon-Steensma et al. 2016) and the branching level of five representative individuals per species were measured. The branching level consisted of the following categories: 1(=stem without branches), 2(=stem with branches), 3(=stem with branched branches), and sometimes 4(=stem that is branched three times). When less than five individuals of a species with a cover above 1% were present, measurements on this species were usually also done on other plant material of the plot. To obtain one value per plot for the stem height, stem diameter and branching level, weighted means were calculated by correcting the average per species for the stem density per species by the following equation:

Weighted mean

$$= \frac{\sum(\text{average per species} * \text{stem density per species})}{\text{stem density per plot}}$$

This equation is derived from the frequently used calculation of the community weighted mean of plant traits (CWM) (Lavorel et al. 2008). The clipped plant material was dried in a stove at 70 °C for 48 h and weighed to calculate the biomass per square meter.

Sedimentation

Sedimentation was measured over a period of two weeks covering one spring tide in the middle of the study period. Four sediment traps (Reed 1989; Nolte et al. 2013) were placed per subplot in September and secured with iron pegs (Fig. 1d). A sediment trap consisted of an open plastic petri dish (diameter of 9 cm) and a filter paper in it. The filter paper was kept in place by wetting it and a small marble on top. During collection of the sediment traps, remaining water in the traps was filtered through new filter paper. Sediment traps and filter papers with sediment were dried in a stove at 60 °C for 66 h and weighed (Leonard et al. 1995; Reed et al. 1997). Sedimentation over a period of two weeks was expressed per square meter for each subplot. Storm surge led to a study

period of eleven and seventeen days for two blocks. This deviation from the original fourteen days was accounted for statistically in the block effect.

Sediment Grain Size

Samples were taken of the upper 1 cm of the soil for analysis of the grain size distribution. Two samples were taken from each of the four corners of a plot and mixed. A pre-treatment was performed to a spoon of soil from each plot following the description by Konert and Vandenberghe (1997). Organic material and calcium carbonate were removed with 30% hydrogen-peroxide (H_2O_2) and 10% hydrochloric acid (HCl), respectively. Remaining coarse organic particles and shells were removed by carefully sieving (sieve size of 200 μm). A Sympatec HELOS/KR Laser Diffraction Machine (range of 0.1–2000 μm) was used to determine the grain size distribution of the pre-treated samples. The median grain size was used from the output for the data analysis.

Data Analysis

The R software package, version 3.5.1, was used to analyse the data. The data was obtained either at the plot level (distance to the creek, elevation, seven vegetation characteristics, median grain size) or at the subplot level (sedimentation). Differences between the four vegetation types (*Salicornia*, *Spartina*, *Atriplex*, *Limonium* type) were tested with a one-way ANOVA for elevation and with Kruskal-Wallis rank sum tests for distance to the creek (separately for the short and long distance category) and the seven vegetation characteristics. The tests were followed by a Tuckey HSD post hoc test and multiple comparisons with the package *pgirmess* (Giraudeau 2018), respectively.

Linear mixed models (LMM) were made with the package *nlme* (Pinheiro et al. 2018) to test the effect of vegetation on sedimentation and median grain size. The response variables sedimentation and median grain size were always log transformed to meet the model assumptions. First, we tested as fixed effects vegetation presence (present, removed) (only for sedimentation), vegetation type, distance to the creek (short, long) and interactions. The random structure of the models only consisted of random intercepts and was based on the study design: distance to the creek (only for sedimentation) within vegetation type within the random factor block. Final mixed models were obtained after backward model simplification using Likelihood Ratio tests. Multiple comparisons were made for vegetation type using the package *multcomp* (Hothorn et al. 2008).

Second, we tested the effect of one or two vegetation characteristics on sedimentation and median grain size at a time, while correcting for the most important geomorphological factor and vegetation presence (only for sedimentation). For

median grain size, two groups of plots existed with either a low (42 plots, $18.28 \pm 0.83 \mu\text{m}$, mean \pm standard error) or a high median grain size (14 plots, $175.91 \pm 14.24 \mu\text{m}$), leading to violation of model assumptions. Therefore, only plots with a low median grain size were used for the response variable median grain size. Elevation was included as fixed effect for sedimentation as it was stronger correlated with sedimentation (Spearman rank correlation, $\rho = -0.49$) than distance to the creek (m) ($\rho = -0.27$), while distance to the creek was included for median grain size as it was stronger correlated with median grain size ($\rho = -0.24$) than elevation ($\rho = 0.15$, Table 1). Interactions were included between two vegetation characteristics and for sedimentation between vegetation presence and the vegetation characteristics. Variables displaying high co-linearity (Spearman rank correlation, $-0.6 > \rho > 0.6$) were not included together in a model (Table 1). The above resulted into six mixed models with one vegetation characteristic and ten mixed models with two vegetation characteristics as fixed effects for sedimentation. For median grain size, seven mixed models with one vegetation characteristic and sixteen mixed models with two vegetation characteristics were made. Stem density was log transformed because of some extreme high values. The random structure of the models for sedimentation was defined as plot within block. Only the random factor block was included for median grain size. The models were ranked based on the Second-order Akaike Information Criterion (AICc) with the package *MuMIn* (Barton 2018). Backward model simplification was applied to all models with one vegetation characteristic. For the models with two vegetation characteristics, simplification was only applied to the model(s) with the lowest AICc (or similar with a difference < 2). Marginal and conditional pseudo R^2 were calculated for the linear mixed models with the package *MuMIn* (Barton 2018). Statistics are reported for the final models.

Modelling the Potential Influence of Real Vegetation Characteristics on Wave Height with XBeach

Subsequently, we explored how the observed variation of vegetation characteristics in the field would translate to wave attenuation in XBeach. This numerical model has been developed by Roelvink et al. (2009) and extended by van Rooijen et al. (2016) with vegetation modelled as an energy sink for propagating waves (see Appendix for details). Individual plants are represented by cylinders, and the size, density and drag properties of these cylinders can be adapted to the stem height, diameter, density and stiffness of the plant species studied. After calibration of the bulk drag coefficient, the extended XBeach model can adequately predict wave attenuation (van Rooijen et al. 2016; Garzon et al. 2019a, 2019b).

For this simple exploration, a drag coefficient of 1 was used, although the drag force exerted on the vegetation can

Table 1 Spearman rank correlations between field measurements on sedimentation or median grain size and elevation, distance to the creek and seven vegetation characteristics. Correlation coefficients smaller than -0.6 or larger than 0.6 are indicated in bold

	Elevation (m+NAP)	Distance to creek (m)	Total cover (%)	Vegetation height (cm)	Stem height (cm)	Stem diameter (mm)	Branching level	Stem density (#/m ²)	Biomass (g/m ²)
Sedimentation (g/m ²) ^a	-0.49	-0.27	-0.25	0.27	0.25	0.36	0.12	-0.36	0.09
Median grain size (μm) ^b	0.15	-0.24	0.28	0.09	0.17	0.04	0.26	-0.19	-0.01
Elevation (m+NAP) ^c	-	0.20	0.70	-0.05	0.05	-0.23	0.27	0.22	0.15
Distance to creek (m) ^c	0.20	-	-0.06	-0.05	-0.11	0.05	0.06	-0.03	0.02
Total cover (%) ^c	0.70	-0.06	-	0.28	0.42	0.07	0.46	0.15	0.53
Vegetation height (cm) ^c	-0.05	-0.05	0.28	-	0.80	0.75	0.23	-0.14	0.74
Stem height (cm) ^c	0.05	-0.11	0.42	0.80	-	0.80	0.46	-0.38	0.74
Stem diameter (mm) ^c	-0.23	0.05	0.07	0.75	0.80	-	0.40	-0.59	0.59
Branching level ^c	0.27	0.06	0.46	0.23	0.46	0.40	-	-0.51	0.39
Stem density (#/m ²) ^c	0.22	-0.03	0.15	-0.14	-0.38	-0.59	-0.51	-	0.03
Biomass (g/m ²) ^c	0.15	0.02	0.53	0.74	0.74	0.59	0.39	0.03	-

^a $n = 112$ ^b $n = 42$ (only plots with low median grain size)^c $n = 56$

vary considerably (Defina and Bixio 2005; Anderson and Smith 2014; Möller et al. 2014; Vuik et al. 2016). While dissipation by vegetation is the main interest of this model exploration, other important dissipation terms across the foreshore were included as well: dissipation by wave breaking (Roelvink 1993), friction, and dissipation by roller energy. Parameters for these processes were kept at the default values of XBeach representing a sandy environment like the Slufter. Other processes of wave energy dissipation or wave energy generation were not included.

We simulated a small storm event within the Slufter comparable to the sheltered environment of the Wadden sea (water level of 2.8 m + NAP; wave height of 0.9 m, wave period of 5 s). The significant wave height (H_{m0}) was modelled along a 1D transect in the Slufter based on the digital elevation map of the Netherlands (AHN3) (Fig. 1b; Fig. S1, Online Resource). The transect was made through the opening of the Slufter perpendicular to the sand dike eastward of the salt marsh with a length of 5.4 km, of which the last 2.5 km over the salt-marsh vegetation.

Vegetation Scenarios

Wave attenuation was modelled for six vegetation scenarios: vegetation absent, *Salicornia* type, *Spartina* type, *Atriplex* type, *Limonium* type and zonation of the four vegetation types. The first scenario, with only topological data without

vegetation, served as a reference to compare vegetation effects. The next four scenarios with one vegetation type were developed to compare the effect of different vegetation types on wave attenuation. The last scenario represents the vegetation in the Slufter.

In each scenario, no vegetation was present below 0.61 m + NAP, based on the minimum recorded elevation of the plots. In the scenario with four vegetation types, plants representing the *Salicornia* or *Spartina* type (pioneer salt-marsh vegetation) both occupied the zones between 0.61 m and 0.99 m + NAP, while plants representing the *Atriplex* or *Limonium* type (low salt-marsh vegetation) both occupied elevations above 0.99 m + NAP. This distribution was based on plot elevations recorded in the field. Data on vegetation characteristics of the medium and high salt marsh was not available as plots were not established here during our field experiment.

The values for the stem height, stem diameter and stem density of the plants in the model were based on the mean per vegetation type measured in the field experiment (Fig. 2d–g; Table 2). A high variation, as indicated by the large standard error, was observed for the stem density of the *Salicornia* and *Spartina* types, because some plots contained a large number of small *Puccinellia maritima* (Fig. 2g). These extreme densities, that were more than two times larger than the mean, were excluded from calculations of the mean for the model input.




Fig. 2 Means with standard error of the **a** elevation, **b** total cover, **c** vegetation height, **d** stem height, **e** stem diameter, **f** branching level, **g** stem density and **h** biomass for the vegetation types *Salicornia* (sal), *Spartina* (spa), *Atriplex* (atr) and *Limonium* (lim). The capital letters above the bars indicate significant groups based on a Tuckey HSD post hoc test for the elevation (**a**) and multiple comparisons after a Kruskal-Wallis rank sum test for the vegetation characteristics (**b-h**)

Table 2 Overview of the vegetation characteristics used in the vegetation and sensitivity scenarios with the XBeach model and the resulting wave attenuation percentages for the first 100 m of the salt-marsh vegetation and over the total length of the salt-marsh vegetation

(2.5 km) for a water level of 2.8 m + NAP, a wave height of 0.9 m and a wave period of 5 s. The salt-marsh vegetation occurs in the last 2.5 km of the transect (5.4 km)

	Stem height (cm)	Stem diameter (mm)	Stem density (#/m ²)	Wave attenuation (%)	
				first 100 m	2.5 km
Vegetation scenarios					
Vegetation absent	–	–	–	–3.03	12.72
<i>Salicornia</i> type	13.19	1.19	2128	29.30	94.92
<i>Spartina</i> type	30.57	2.78	2330	71.68	99.12
<i>Atriplex</i> type	25.35	1.86	2222	45.41	97.35
<i>Limonium</i> type	15.89	0.88	4680	57.27	98.34
Zonation of vegetation types	a	a	a	59.66	98.00
Sensitivity scenarios					
Average scenario	21.25	1.68	2887	56.84	98.31
Min. stem height	6.85	1.68	2887	29.17	94.89
Max. stem height	60.06	1.68	2887	79.01	99.41
Min. stem diameter	21.25	0.60	2887	31.50	95.38
Max. stem diameter	21.25	4.55	2887	78.01	99.37
Min. stem density	21.25	1.68	96	1.64	65.74
Max. stem density	21.25	1.68	9288	80.79	99.47

^aThe vegetation characteristics of all four vegetation types are applied

Sensitivity Scenarios

To explore the influence of natural variance in key vegetation characteristics on modelled wave attenuation, a sensitivity analysis was performed based on the observed variation of vegetation characteristics in the field. Wave attenuation was modelled for seven scenarios, under the same conditions as described before (Table 2). For the baseline scenario, the mean of the stem height, stem diameter and stem density based on all plots was applied to the whole transect above 0.61 m + NAP. For the other scenarios, one vegetation characteristic was changed at a time, either to the minimum or maximum value of all plots. This analysis will give an indication of the relative influence of natural variations in vegetation characteristics on wave attenuation. The same extreme stem density values as described before were excluded when calculating the mean and maximum stem density. Wave attenuation percentages were calculated over the salt-marsh vegetation for all scenarios based on the significant wave height.

Results

Plant Species and Communities

A total of seventeen plant species were identified in the plots (Table S1, Online Resource). Based on the cover percentage of these species, the plots were classified into four different plant communities. All plots of the *Salicornia* type belonged to the plant community of *Salicornietum dolichostachyae*. Most of the *Spartina* type plots were classified as the *Spartinetum townsendii* plant community. The plots of the *Limonium* type all belonged to the *Plantagini-Limonietum* plant community and all the *Atriplex* type plots belonged to the *Halimionetum portulacoides* plant community. However, also five plots of the *Spartina* type were classified as the latter plant community because both *S. anglica* and *A. portulacoides* had a high cover percentage in these plots.

Comparison of Vegetation Types

The elevation of the plots differed significantly between the vegetation types (one-way ANOVA: $F = 26.19$, $df = 3$, $p < 0.001$) and increased from the *Salicornia* type, to the *Spartina*, *Atriplex* and *Limonium* type (Fig. 2a). The distance to the creek was 3.54 ± 0.56 m (mean \pm standard error) for plots of the short distance category and 8.96 ± 0.78 m for plots of the long distance category without significant differences between the vegetation types (Kruskal-Wallis rank sum test: short, $\chi^2 = 0.39$, $df = 3$, $p = 0.94$; long, $\chi^2 = 1.09$, $df = 3$, $p = 0.78$). It should be noted that the similar distance of the vegetation types to the creek was important for our study design and characteristic of our study area.

All vegetation structure characteristics differed significantly between the vegetation types (Table 3; Fig. 2b-h). The *Salicornia* and *Limonium* types were both characterized by low plant biomass, low vegetation height and short, mostly unbranched stems with a small diameter. However, total cover was on average lower for the *Salicornia* type and varied more, compared to the other vegetation types where total cover was almost 100%. The *Spartina* type was characterized by high plant biomass, tall vegetation and long, thick and mostly unbranched stems. The *Atriplex* type also had high plant biomass, but a lower vegetation height and long, highly branched stems with an intermediate stem diameter. The high mean and large variation in stem density for the *Salicornia* and *Spartina* types was caused by some plots with extreme stem densities due to a large number of *P. maritima*.

Sedimentation

Vegetation presence (LMM: $df = 55$, $F = 16.63$, $p < 0.001$), vegetation type (LMM: $df = 18$, $F = 9.50$, $p < 0.001$) and distance to the creek (LMM: $df = 27$, $F = 11.52$, $p = 0.002$) all had a significant effect on sedimentation (marginal $R^2 = 0.346$, conditional $R^2 = 0.905$; note that sedimentation was always log transformed) (Fig. 3). Sedimentation was on average 42% higher in the presence of vegetation compared to locations where vegetation was removed. This positive effect of vegetation on sedimentation did not significantly differ between vegetation types or between distances to the creek, as illustrated by the absence of significant interaction effects. No obvious signs of erosion of the soil surface were observed in the subplots where vegetation was removed. Only the *Limonium* type was significantly different from the other vegetation types. Sedimentation was on average more than five times higher for the *Atriplex* and *Spartina* types and more than eleven times higher for the *Salicornia* type compared to the *Limonium* type. On average 60% more sedimentation took place in plots at a short distance to the creek compared to

Table 3 Chi-squared, degrees of freedom and p values for Kruskal-Wallis rank sum tests to compare seven vegetation characteristics between the vegetation types *Salicornia*, *Spartina*, *Atriplex* and *Limonium*

Vegetation characteristic	χ^2	df	p value
Total cover	31.89	3	<0.001
Vegetation height	38.29	3	<0.001
Stem height	26.39	3	<0.001
Stem diameter	32.78	3	<0.001
Branching level	26.80	3	<0.001
Stem density	10.57	3	0.014
Biomass	32.23	3	<0.001




Fig. 3 Means with standard error of sedimentation over a period of two weeks in four vegetation types. In each vegetation type, plots are located at a short or long distance to the creek. In each plot, vegetation is removed in one of the two subplots. The capital letters above the bars of each vegetation type indicate significant groups based on multiple comparisons after a linear mixed model

the long distance category. No interaction effect was found between distance to the creek and vegetation type on sedimentation.

Although sedimentation was mostly correlated with stem diameter and stem density from the vegetation characteristics based on Spearman rank correlations (Table 1), the linear mixed model with branching level was the best from the six models with one vegetation characteristic as the AICc was lowest and the marginal pseudo R^2 was highest (Table 4). Stem height (LMM: $df = 47$, $F = 3.93$, $p = 0.053$; marginal $R^2 = 0.320$, conditional $R^2 = 0.901$) and branching level (LMM: $df = 47$, $F = 10.02$, $p = 0.003$; marginal $R^2 = 0.363$, conditional $R^2 = 0.901$) were significant in the models, while other vegetation characteristics were not significant. Sedimentation increased exponentially with both stem height and branching level (Fig. 4a, b; note the logarithmic scale of the y-axes).

Branching level was also important based on the analysis of the ten models with two vegetation characteristics. In this case, the model with the combination branching level and stem diameter was the best based on the AICc and marginal pseudo R^2 (Table 4). The interaction between branching level and stem diameter was significant (LMM: $df = 45$, $F = 4.43$, $p = 0.041$; marginal $R^2 = 0.398$, conditional $R^2 = 0.904$). Branching level significantly affected sedimentation in the final model (LMM: $df = 45$, $F = 10.69$, $p = 0.002$), while stem diameter was not significant (LMM: $df = 45$, $F = 1.08$, $p = 0.30$). The interaction between stem diameter and branching level suggests that branching level only has a positive effect on sedimentation when the stem is thick enough (above 1.3 cm). The AICc of the model with the combination branching level and stem height was similar, but only branching level was significant in this model (LMM: $df = 47$, $F = 10.02$, $p = 0.003$; marginal $R^2 = 0.363$, conditional $R^2 = 0.901$).

Elevation (LMM: $df = 45-47$, $F = 37.59-45.28$, $p < 0.001$; df and F depend on the model considered) and vegetation presence (LMM: $df = 55$, $F = 16.63$, $p < 0.001$) were always significant in the models with one or two vegetation characteristics. Sedimentation decreased exponentially with elevation (Fig. 4c; note the logarithmic scale of the y-axes) and was positively affected by vegetation presence as before (Fig. 4). Interactions between vegetation presence and vegetation characteristics were never significant, suggesting that stem height and branching level of the surrounding vegetation still affected sedimentation in the subplots where vegetation was removed.

Sediment Grain Size

A significant effect was found of vegetation type on median grain size (LMM: $df = 18$, $F = 4.87$, $p = 0.012$), while distance to the creek or an interaction was not significant (marginal $R^2 = 0.320$; conditional $R^2 = 0.859$; note that median grain size was always log transformed) (Fig. 5). Median grain size was on average more than two times higher for the *Atriplex* and *Spartina* types and almost seven times higher for the *Salicornia* type compared to the *Limonium* type. However, only the median grain size of the *Salicornia* type was significantly different from the other vegetation types. This higher grain size for the *Salicornia* type was reflected in a high sand percentage of on average 58% compared to 18–27% for the other vegetation types.

The effect of vegetation characteristics on median grain size was studied only for plots with a low median grain size. Total cover and branching level had the highest Spearman rank correlation coefficients with median grain size from the seven vegetation characteristics (Table 1). The model with branching level also had the lowest AICc and highest marginal pseudo R^2 , compared to the other models for median grain size with one vegetation characteristic (Table 4). However, none of the vegetation characteristics were significant. From the sixteen mixed models with two vegetation characteristics, the model with the combination branching level and stem diameter was the best model based on their AICc and marginal pseudo R^2 (Table 4). The models with the combinations biomass and branching level or vegetation height and branching level were similar (AICc difference < 2). However, the vegetation characteristics or interactions were not significant. Branching level and distance to the creek (m) were both just not significant ($0.1 > p > 0.05$) in the models for median grain size, but a positive and negative exponential trend can be observed, respectively (Fig. S2, Online Resource; note the logarithmic scale of the y-axes). Very little variance is explained in the models for median grain size with vegetation characteristics as the marginal and conditional pseudo R^2 were very low (Table 4).

Table 4 Second-order Akaike Information Criterion (AICc) and marginal (m) and conditional (c) pseudo R^2 of several linear mixed models for sedimentation or median grain size with one or two vegetation characteristics. Note that sedimentation, median grain size and stem density were always log transformed and that backward model simplification is not yet applied. Lowest or similar AICc (difference < 2) and highest marginal pseudo R^2 are indicated in bold

Vegetation characteristics	Sedimentation			Median grain size		
	AICc	R^2 (m)	R^2 (c)	AICc	R^2 (m)	R^2 (c)
–	–	0.296	0.900	–	0.083	0.083
Total cover	–	–	–	–52.8	0.089	0.089
Vegetation height	67.1	0.305	0.900	–52.9	0.093	0.093
Stem height	61.9	0.343	0.900	–52.5	0.083	0.083
Stem diameter	64.4	0.326	0.899	–52.6	0.085	0.085
Branching level	58.7	0.384	0.897	–56.0	0.158	0.158
Stem density	66.1	0.320	0.897	–53.6	0.108	0.108
Biomass	65.7	0.324	0.899	–52.5	0.084	0.084
Total cover*vegetation height	–	–	–	–47.9	0.106	0.106
Total cover*stem height	–	–	–	–47.3	0.093	0.093
Total cover*stem diameter	–	–	–	–47.5	0.098	0.098
Total cover*branching level	–	–	–	–51.4	0.179	0.179
Total cover*stem density	–	–	–	–48.4	0.115	0.115
Total cover*biomass	–	–	–	–49.4	0.138	0.138
Vegetation height*branching level	64.1	0.417	0.898	–53.3	0.216	0.216
Vegetation height*stem density	72.1	0.355	0.895	–48.7	0.122	0.122
Stem height*branching level	63.0	0.407	0.903	–51.2	0.175	0.175
Stem height*stem density	64.8	0.394	0.903	–48.4	0.117	0.117
Stem diameter*branching level	61.1	0.428	0.902	–53.9	0.226	0.226
Stem diameter*stem density	69.2	0.344	0.903	–48.9	0.127	0.127
Stem diameter*biomass	68.9	0.335	0.908	–47.1	0.088	0.088
Branching level*stem density	67.3	0.385	0.899	–50.6	0.162	0.162
Branching level*biomass	64.7	0.394	0.903	–53.7	0.223	0.223
Stem density*biomass	72.2	0.353	0.898	–48.5	0.120	0.120

Modelled Wave Attenuation

The significant wave height that was modelled with XBeach along a transect in the Slufter increased on the beach and then slowly declined when vegetation was absent or declined exponentially when vegetation was present until the dike was reached (Fig. 6; Table 2). The modelled total wave attenuation over the salt marsh was clearly higher for the scenario with zonation of the vegetation types (wave attenuation of 98%) compared to the scenario where vegetation is absent (13%). Differences between the vegetation types were most pronounced for the first 100 m of salt-marsh vegetation. The *Salicornia* type with the shortest stems had the weakest modelled wave attenuation (29%). Wave attenuation was higher for the *Atriplex* type (45%), which had taller and thicker stems with a similar density, and for the *Limonium* type (57%), which had similar stem heights and stem diameters as the *Salicornia* type but twice the stem density. Wave attenuation was strongest for the *Spartina* type (72%), which had the tallest and thickest stems.

The sensitivity analysis of vegetation characteristics based on field measurements indicated that modelled wave

attenuation was most sensitive to the natural variance in stem density (Fig. 7; Table 2). As expected, stem height, stem diameter and stem density all positively influenced wave attenuation in the XBeach model. The maximum scenarios for the stem height, stem diameter and stem density resulted in equally more wave attenuation (78–81% wave attenuation for the first 100 m) compared to the average scenario (57%). The minimum scenarios resulted in an equally weaker wave attenuation for stem height and stem diameter (29 and 32%), but to a very strong reduction in the wave attenuation capacity for stem density (1.6%). The total wave attenuation over the salt-marsh vegetation was still low for the minimum stem density scenario (66%) compared to the other scenarios (95–99%).

Discussion

Vegetation Presence

The presence of vegetation increased sedimentation on average by 42% relative to small-scale vegetation removal in our study area, illustrating the importance of vegetation presence




Fig. 4 The vegetation characteristics **a** stem height and **b** branching level significantly influenced sedimentation in two separate linear mixed models, together with vegetation presence and **c** elevation. Measurements are indicated per vegetation type, although the effect of vegetation type on sedimentation is tested in another model. Note that the y-axes have a logarithmic scale

for the adaptive capacity of salt marshes to sea level rise. Due to our study design, we were able to rule out geomorphological effects such as elevation and distance to the source of sediment. Although we were not able to fully exclude that plant material from the surrounding vegetation in subplots where vegetation was present contributed to the dry weight in the sediment traps, we expect this to have a limited contribution to the vegetation effect. Our results are consistent with Culberson et al. (2004), who measured on average almost two times more sedimentation in vegetated locations compared to unvegetated locations of 1 m² in two estuarine marshes. Field and modelling studies showed that large-scale vegetation removal can result in spatially redistributed sedimentation (Temmerman et al. 2005; Schepers et al. 2020), sheet flow and increased flow velocities over the marsh platform (Temmerman et al. 2005; Temmerman et al. 2012; Schepers et al. 2020). Whether the reduction in sediment deposition we observed after small-scale removal of marsh vegetation is due to a locally enhanced water flow velocity needs further study. Our results confirm the major contribution of salt-marsh vegetation to sedimentation, which is important for the vertical growth of salt marshes to keep up with sea level rise. This importance of vegetation is also illustrated by studies that found decreased accretion rates, erosion and loss of elevation after vegetation dieback (Baustian et al. 2012; Coleman and Kirwan 2019).

We investigated the vegetation effect on sedimentation in four vegetation types. No statistically significant interaction was found between vegetation presence and vegetation type on sedimentation. However, the presence of vegetation increased sedimentation on average by 14, 29, 28 and 60% relative to vegetation removal for the vegetation types *Limonium*, *Atriplex*, *Spartina* and *Salicornia* respectively. This suggests the strongest impact for vegetation types where elevation is low and sediment availability is high, stressing the importance of the pioneer *Salicornia* species as key initiators




Fig. 5 Means with standard error of median grain size of the sediment in four vegetation types. In each vegetation type, plots are located at a short or long distance to the creek. The capital letters above the bars of each vegetation type indicate significant groups based on multiple comparisons after a linear mixed model

of the biophysical feedback between salt-marsh vegetation and sedimentation.

The wave attenuation of 98% modelled in XBeach over the 2.5 km transect representing the salt marsh in the Slufter suggests that the significant wave height is almost zero in front of the sand dike under modelling conditions. This is mostly attributed to the presence of vegetation, not topography, in our study area, as modelled wave attenuation was only 13% when vegetation was absent. Our modelled wave attenuation (76% over the first 180 m) is in line with the measured wave attenuation by Möller et al. (1999) for low marsh vegetation with a similar species composition (on average 61% for a 180 m transect). Wave attenuation over the salt marsh was around 50% higher than over a bare sand flat seaward of the salt marsh for similar water depths (Möller et al. 1999). While our 1D approach is sufficient to explore potential wave attenuation by vegetation, a 2D model that also includes processes like diffraction or wave-current interactions will give more accurate predictions of the wave height in the Slufter. Given that we modelled the peak of the storm at high tide (when tidal currents are minimal) and did not include wave diffraction, wave heights can be expected to reduce further in a 2D approach. The exponential decrease of modelled wave height over vegetated salt marshes we observed is also found by field studies (Möller et al. 1999; Yang et al. 2012) and indicates that wave height is mainly reduced in the first part of the salt marsh. This suggests that proper management of the most

seaward part of a salt marsh is important with respect to flood protection.

Vegetation Type

Sedimentation was highest in the *Salicornia* type, intermediate for the *Spartina* and *Atriplex* types and lowest for the *Limonium* type. The same pattern was observed for sediment grain size. Sedimentation differences between vegetation types were also observed by Silva et al. (2009) in Portugal and by French and Spencer (1993) in the UK as a result of differences in distance to the creek (French and Spencer 1993; Silva et al. 2009) and elevation (French and Spencer 1993). Yang (1999a) and Yang et al. (2008) observed larger grain sizes for low marsh vegetation compared to high marsh vegetation in the Yangtze River Delta, China. This is likely due to a landward decrease of the flow velocity and wave energy (Yang 1999a; Yang et al. 2008). The vegetation types in our study area were similar in their distance to the creek, controlling for the distance to the sediment source, but did differ in elevation and vegetation characteristics. As sedimentation and grain size is generally found to decrease with elevation (e.g., Yang 1999b; Yang et al. 2008; Townend et al. 2011; Reef et al. 2018), sedimentation and grain size differences between the vegetation types in our study are most likely explained by a combination of elevation and vegetation characteristics. Our results confirm that large sediment particles are rapidly deposited in the *Salicornia* vegetation as water enters the salt marsh and thus this pioneer vegetation will enhance accretion more than in the higher marsh.




Fig. 6 Significant wave height (H_{m0}) along the transect (5.4 km) modelled with XBeach for six vegetation scenarios and for a water level of 2.8 m + NAP, a wave height of 0.9 m and a wave period of 5 s. The vegetation scenarios are based on field measurements of vegetation characteristics. The first scenario (vegetation absent) served as a reference. One of the four vegetation types is present in the next four scenarios (*Salicornia* type, *Spartina* type, *Atriplex* type, *Limonium* type). The last scenario (zonation of the four vegetation types) represents the field situation. The salt-marsh vegetation occurs in the last 2.5 km of the transect




Fig. 7 Significant wave height (H_{m0}) along the transect (5.4 km) modelled with XBeach for seven sensitivity scenarios and for a water level of 2.8 m + NAP, a wave height of 0.9 m and a wave period of 5 s. The sensitivity scenarios for three vegetation characteristics are based on field measurements. Only the last 2.5 km of the transect is shown, where salt-marsh vegetation occurs. The scenario where vegetation is absent is shown for comparison. Note that the significant wave height of the three maximum scenarios is overlapping

Modelled wave attenuation was strongest for the *Spartina* type (72% for the first 100 m) and weakest for the *Salicornia* type (29%). In contrast to sedimentation, the differences in modelled wave attenuation between the vegetation types can only be attributed to differences in values for the height, diameter and density of the stems, as the elevation profile and drag coefficient were the same in all scenarios. Stems of the *Spartina* type were more than two times taller and thicker than the stems of the *Salicornia* type. In the field, however, water depth and incoming wave height can differ between vegetation types and will reduce or increase wave attenuation, respectively (e.g., Möller et al. 1999; Möller 2006; Maza et al. 2015; Vuik et al. 2016). For simplicity, the drag coefficient was the same in all scenarios, although it is likely to vary between vegetation types as vegetation characteristics can influence the drag coefficient (Nepf 2011). More flexible vegetation, for example, can have a lower drag coefficient than rigid vegetation (van Veelen et al. 2020). The stems of the dominating species in the vegetation types we studied are relatively stiff, although some other species are more flexible like *Puccinellia maritima* and *Suaeda maritima* (Bouma et al. 2013; Möller et al. 1999). Vuik et al. (2016) observed more wave attenuation by a *Scirpus maritimus* marsh compared to a *S. anglica* marsh for the same wave height and water depth, likely due to a higher biomass density of the *S. maritimus* marsh. We found that modelled wave attenuation was stronger for the scenario with a zonation of the four vegetation types (60% for the first 100 m) compared to the average wave damping of the individual vegetation types weighted to their occurrence (51%), indicating the importance of including vegetation zonation in the model. Field measurements of wave attenuation in various vegetation types are required to validate our modelling results.

Vegetation Characteristics

While positive effects of biomass (field experiment, Morris et al. 2002) and stem density (laboratory experiment, Gleason et al. 1979) on sedimentation are reported for *Spartina alterniflora*, we found that especially stem height and branching level positively affected sedimentation in our study area. Our study is distinguished by the relatively large number of vegetation characteristics and species considered in the field. No effects of vegetation height and biomass were observed for *Spartina alterniflora* in a field flume experiment during calm summer conditions by sequential removal of plant material (Reef et al. 2018). These results suggest that effects of vegetation characteristics can depend on the season, the species and on the importance of vegetation relative to geomorphological and hydrodynamic factors (e.g., Davidson-Arnott et al. 2002; Wood and Hine 2007; Townend et al. 2011). Based on our study, tall (e.g., *Spartina* species) and highly branched (e.g.,

A. portulacoides) salt marsh vegetation will best enhance sedimentation. However, it should be noted that the elevation range where species grow determine sedimentation as well. We encourage to further investigate the role of vegetation characteristics in sedimentation for multiple sites, species, seasons and weather conditions and for a longer period. Artificial manipulation of vegetation characteristics could be investigated with the method used by Reef et al. (2018) with treatments at locations subjected to the same geomorphological effects (e.g., elevation, distance to the sea) as in our experiment for vegetation presence.

Although we observed a positive effect of branching level on median grain size for plots with a low grain size, we found no significant effects of vegetation characteristics on sediment grain size. Gibeault et al. (2016) also found no significant effects of vegetation height and cover on grain size in a subarctic salt marsh. However, Yang (1999a) observed that sediment was usually finer in denser and higher marsh vegetation in the Yangtze Delta, China. A negative relation was found between grain size and the product of plant height and cover (Yang et al. 2008). Possibly, hydrodynamic factors were more dominant in our study system, resulting in plots with either a large or small median grain size. An alternative explanation could be that current vegetation characteristics are not representative for the period of sedimentation that is represented by the top 1 cm of the sampled soil used for measuring the grain size distribution.

Unsurprisingly, an increase in stem height, stem diameter and stem density all increased modelled wave attenuation. These results are consistent with flume experiments, which showed positive effects on wave attenuation by the stem height to water depth ratio and stem density (e.g., Augustin et al. 2009; Bouma et al. 2010; Anderson and Smith 2014; Peruzzo et al. 2018). Based on a sensitivity analysis using field measurements of stem height, stem diameter and stem density, modelled wave attenuation in our study was most sensitive to the natural variation in stem density, dropping strongly for the minimum stem density of 96 stems/m². The relative importance of stem density is also indicated by other modelling studies (van Loon-Steensma et al. 2014; Kalra et al. 2017; Garzon et al. 2019b). This suggests that salt marshes with low density vegetation are less suited for flood protection by attenuating waves, which is relevant to consider for constructing or restoring salt marshes. Besides, our results imply that maximizing vegetation characteristics like the density could increase wave attenuation. However, field experiments testing the relative importance of vegetation characteristics on wave attenuation are required to validate our modelling results.

Vegetation characteristics, such as stem height, diameter and density, can decrease during winter, potentially reducing the positive vegetation effect on sedimentation and wave attenuation (Silinski et al. 2016; Garzon et al. 2019b). This reduction will likely be larger for the *Salicornia* type relative

to the *Spartina*, *Atriplex* and *Limonium* types, because annual *Salicornia* species can completely disappear during winter (Bouma et al. 2014), while the perennial plants *S. anglica*, *A. portulacoides* and *L. vulgare* will only lose part of their biomass. As a result, the capacity of the salt marsh to aid in flood protection can be severely reduced depending on the plant species present.

Implications for Salt-Marsh Functioning and Future Challenges

Our results underscore the importance of plant species composition in salt marshes for sedimentation and expected wave attenuation. Therefore, the composition of the salt marsh vegetation is crucial to understand the biophysical feedbacks between inundation, plant growth and sediment deposition and predict the marsh's response to sea level rise (Kirwan et al. 2016). Because of the natural adaptive capacity of salt marshes and potential wave attenuation by vegetation, salt marshes can be considered in nature-based solutions for coastal protection (Temmerman et al. 2013; Vuik et al. 2016). Our study indicates that these services are influenced by vegetation type and vegetation characteristics. We found that sedimentation increased with stem height and branching level. Our model exploration suggests that natural variation in vegetation characteristics, especially stem density, has a significant influence on modelled wave attenuation rates. This implies that changes in vegetation characteristics and species composition in the future, due to for example climate change and invasive species, can affect salt-marsh functioning (e.g., Wang et al. 2006; Kirwan et al. 2009; Ysebaert et al. 2011; Watson et al. 2014; Ratliff et al. 2015). Although further field research is required to gain more insights in the relative importance of vegetation characteristics to geomorphological and hydrodynamic factors, management can be adapted to this by stimulating plant growth or promoting specific species to maximize vegetation characteristics. Furthermore, our results suggest that model predictions of salt-marsh accretion and wave attenuation can be improved by incorporating differences in vegetation characteristics between vegetation types.

Conclusion

The purpose of this study was to investigate the role of salt-marsh vegetation in sedimentation, sediment grain size and modelled wave attenuation in the Slufter, the Netherlands, with an outlook for implementing the natural adaptive capacity of salt marshes to sea level rise in nature-based flood defences. Vegetation presence increased sedimentation on average by 42% and positively affected modelled wave attenuation, which is consistent with the results of a meta-analysis by Shepard et al. (2011). Sedimentation, sediment grain size and

modelled wave attenuation were all different for the four vegetation types. Stem height and branching level were found to affect sedimentation, while a non-significant effect of branching level was found on sediment grain size. Modelled wave attenuation was affected most by the natural variability in stem density. Our results suggest that the natural adaptive capacity and flood protection role of salt marshes are affected by the type and characteristics of the vegetation and thereby also by vegetation management.

Appendix: Implementation of vegetation in the XBeach model

In the surf-beat mode used for this study, XBeach propagates waves within the 1D domain following the wave action balance (Roelvink et al. 2009; van Rooijen et al. 2016):

$$\frac{\partial A}{\partial t} + \frac{\partial c_x A}{\partial x} + \frac{\partial c_\theta A}{\partial \theta} = \frac{\sum D}{\sigma}, \quad (1)$$

where: A = Wave action $\frac{J}{m}$,

c = Wave celerity $\frac{m}{s}$,

θ = Wave direction rad ,

D = Wave energy dissipation $\frac{J}{m^2 s^2}$,

σ = Intrinsic frequency $\frac{1}{s}$.

In XBeach, dissipation by a layer of vegetation is modelled with the approach by Suzuki et al. (2012) based on the formulation by Mendez and Losada (2004):

$$D_{v,i} = A_v * \frac{\rho \tilde{C}_{D,i} b_{v,i} N_{v,i}}{2\sqrt{\pi}} \left(\frac{kg}{2\sigma} \right)^3 H_{rms}^3, \quad (2)$$

with $A_v = \frac{\sinh^3 k\alpha_i h - \sinh^3 k\alpha_{i-1} h + 3(\sinh^3 k\alpha_i h - \sinh^3 k\alpha_{i-1} h)}{3k \cosh^3 kh}$,

where: D_v = Dissipation by vegetation $\frac{J}{m^2 s^2}$,

\tilde{C}_D = Bulk drag coefficient –,

b_v = The vegetation stem diameter m ,

N_v = Vegetation density $\frac{stems}{m^2}$,

α = Factor of vegetation height (h_v) as: h_v/h –,

and: ρ = Water density $\frac{kg}{m^3}$,

H_{rms} = Root mean square wave height m ,

k = Wave number $\frac{1}{m}$,

h = Water depth m .

Vegetation in XBeach is modelled as an energy sink for waves. The magnitude of the energy loss is calculated with the vegetation characteristics like stem diameter, density, and height. Results are calibrated with the bulk drag coefficient to account for attributes not included in the model like the shape and deformation of the stems during wave loads.

The XBeach model can adequately predict wave attenuation after calibration of the bulk drag coefficient (van Rooijen

et al. 2016; Garzon et al. 2019a, 2019b). The formulation by Mendez and Losada (2004) is also implemented in the SWAN model by Suzuki et al. (2012) and several studies (e.g., Vuik et al. 2016; Baron-Hyppolite et al. 2019) have demonstrated that after calibration this extended SWAN model can adequately predict wave attenuation across a salt marsh.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s13157-021-01467-w>.

Acknowledgements We are indebted to the Dutch State Forest Service, and more specifically Erik van der Spek, for allowing access to the salt marsh of nature reserve the Slufter (Dutch Wadden island of Texel). Facilities for the grain size distribution analysis were provided by the Vrije Universiteit Amsterdam. We thank dr. Maarten Prins and Martina Hagen of the Sediment Laboratory for help during the grain size distribution analysis and thank Petra Goessen of Water Board Hoogheemraadschap Hollands Noorderkwartier for facilitating this research. Furthermore, we would like to thank the anonymous reviewers for their helpful comments to improve our paper.

Availability of Data and Material The dataset on plant characteristics and sedimentation generated during and/or analysed during the current study is available at <https://doi.org/10.5281/zenodo.4312767>.

Authors' Contributions JvLS and JL conceived ideas. BB designed the field experiment and performed the field work, lab work and data analysis. JK performed the modelling work. JvLS, JL and RM supervised the project. BB wrote the article with support from JvLS, JL, RM and JK.

Funding Water Board Hoogheemraadschap Hollands Noorderkwartier provided funding for the grain size distribution analysis at the Vrije Universiteit Amsterdam.

Declarations

Conflicts of Interest/Competing Interests Authors declare that they have no conflict of interest.

Ethics Approval The responsible nature management organization gave consent for our field experiments.

Consent to Participate Not applicable.

Consent for Publication Not applicable.

Code Availability XBeach is a public domain model (<https://oss.deltares.nl/web/xbeach/>) and SynBioSys is a public domain information system (<https://www.synbiosys.alterra.nl/>).

References

- Adam P (2002) Saltmarshes in a time of change. *Environmental Conservation* 29:39–61
- Anderson ME, Smith JM (2014) Wave attenuation by flexible, idealized salt marsh vegetation. *Coastal Engineering* 83:82–92
- Augustin LN, Irish JL, Lynett P (2009) Laboratory and numerical studies of wave damping by emergent and near-emergent wetland vegetation. *Coastal Engineering* 56:332–340
- Barbier EB, Hacker SD, Kennedy C, Koch EW, Stier AC, Silliman BR (2011) The value of estuarine and coastal ecosystem services. *Ecological Monographs* 81:169–193
- Baron-Hyppolite C, Lashley CH, Garzon J, Miesse T, Ferreira C, Bricker JD (2019) Comparison of implicit and explicit vegetation representations in SWAN hindcasting wave dissipation by coastal wetlands in Chesapeake Bay. *Geosciences* 9:8
- Barton K (2018) MuMIn: Multi-model inference. R package version 1.42.1. <https://CRAN.R-project.org/package=MuMIn>
- Baustian JJ, Mendelsohn IA, Hester MW (2012) Vegetation's importance in regulating surface elevation in a coastal salt marsh facing elevated rates of sea level rise. *Global Change Biology* 18:3377–3382
- Bouma TJ, de Vries MB, Herman PMJ (2010) Comparing ecosystem engineering efficiency of two plant species with contrasting growth strategies. *Ecological Studies* 91:2696–2704
- Bouma TJ, Temmerman S, van Duren LA, Martini E, Vandenbruwaene W, Callaghan DP, Balke T, Biermans G, Klaassen PC, van Steeg P, Dekker F, van de Koppel J, de Vries MB, Herman PMJ (2013) Organism traits determine the strength of scale-dependent bio-geomorphic feedbacks: a flume study on three intertidal plant species. *Geomorphology* 180–181:57–65
- Bouma TJ, van Belzen J, Balke T, Zhu Z, Airoidi L, Blight AJ, Davies AJ, Galvan C, Hawkins SJ, Hoggart SPG, Lara JL, Losada IJ, Maza M, Ondiviela B, Skov MW, Strain EM, Thompson RC, Yang S, Zanuttigh B, Zhang L, Herman PMJ (2014) Identifying knowledge gaps hampering application of intertidal habitats in coastal protection: opportunities & steps to take. *Coastal Engineering* 87:147–157
- Christiansen T, Wiberg PL, Milligan TG (2000) Flow and sediment transport on a tidal salt marsh. *Estuarine, Coastal and Shelf Science* 50:315–331
- Coleman DJ, Kirwan ML (2019) The effect of a small vegetation dieback event on salt marsh sediment transport. *Earth Surface Processes and Landforms* 44:944–952
- Crosby SC, Sax DF, Palmer ME, Booth HS, Deegan LA, Bertness MD, Leslie HM (2016) Salt marsh persistence is threatened by predicted sea-level rise. *Estuarine, Coastal and Shelf Science* 181:93–99
- Culbertson SD, Foin TC, Collins JN (2004) The role of sedimentation in estuarine marsh development within the San Francisco estuary, California, USA. *Journal of Coastal Research* 20:970–979
- Davidson-Amott RGD, van Proosdij D, Ollerhead J, Schostak L (2002) Hydrodynamics and sedimentation in salt marshes: examples from a macrotidal marsh, bay of Fundy. *Geomorphology* 48:209–231
- Defina A, Bixio AC (2005) Mean flow and turbulence in vegetated open channel flow. *Water Resources Research* 41:W07006
- French JR, Spencer T (1993) Dynamics of sedimentation in a tide-dominated backbarrier salt marsh, Norfolk, UK. *Marine Geology* 110:315–331
- Garzon JL, Maza M, Ferreira CM, Lara JL, Losada IJ (2019a) Wave attenuation by *Spartina* saltmarshes in the Chesapeake Bay under storm surge conditions. *Journal of Geophysical Research, Oceans* 124:5220–5243
- Garzon JL, Miesse T, Ferreira CM (2019b) Field-based numerical model investigation of wave propagation across marshes in the Chesapeake Bay under storm conditions. *Coastal Engineering* 146:32–46
- Gedan KB, Silliman BR, Bertness MD (2009) Centuries of human-driven change in salt marsh ecosystems. *Annual Review of Marine Science* 1:117–141
- Gibeault C, Neumeier U, Bernatchez P (2016) Spatial and temporal sediment dynamics in a subarctic salt marsh (gulf of St. Lawrence, Canada). *Journal of Coastal Research* 32:1344–1361
- Giraudoux P (2018) Pgirmiss: spatial analysis and data mining for field ecologists. R package version 1.6.9. <https://CRAN.R-project.org/package=pgirmiss>

- Gleason ML, Elmer DA, Pien NC, Fisher JS (1979) Effects of stem density upon sediment retention by salt marsh cord grass, *Spartina alterniflora* Loisel. *Estuaries* 2:271–273
- Hennekens SM, Smits JHJ, Schaminée JHJ (2018) SynBioSys Nederland. Alterra, Wageningen UR
- Hothorn T, Bretz F, Westfall P (2008) Simultaneous inference in general parametric models. *Biometrical Journal* 50:346–363
- Kalra TS, Aretxabaleta A, Seshadri P, Ganju NK, Beudin A (2017) Sensitivity analysis of a coupled hydrodynamic-vegetation model using the effectively subsampled quadratures method (ESQM v5.2). *Geoscientific Model Development* 10:4511–4523
- Kirwan ML, Megonigal JP (2013) Tidal wetland stability in the face of human impacts and sea-level rise. *Nature* 504:53–60
- Kirwan ML, Guntenspergen GR, Morris JT (2009) Latitudinal trends in *Spartina alterniflora* productivity and the response of coastal marshes to global change. *Global Change Biology* 15:1982–1989
- Kirwan ML, Temmerman S, Skeehean EE, Guntenspergen GR, Fagherazzi S (2016) Overestimation of marsh vulnerability to sea level rise. *Nature Climate Change* 6:253–260
- Konert M, Vandenberghe J (1997) Comparison of laser grain size analysis with pipette and sieve analysis: a solution for the underestimation of the clay fraction. *Sedimentology* 44:523–535
- Lavorel S, Grigulis K, McIntyre S, Williams NSG, Garden D, Dorrough J, Berman S, Quéfier F, Thébaud A, Bonis A (2008) Assessing functional diversity in the field – methodology matters! *Functional Ecology* 22:134–147
- Leonard LA, Hine AC, Luther ME (1995) Surficial sediment transport and deposition processes in a *Juncus roemerianus* marsh, west-Central Florida. *Journal of Coastal Research* 11:322–336
- Leonardi N, Carnacina I, Donatelli C, Ganju NK, Plater AJ, Schuerch M, Temmerman S (2018) Dynamic interactions between coastal storms and salt marshes: a review. *Geomorphology* 301:92–107
- Li H, Yang SL (2009) Trapping effect of tidal marsh vegetation on suspended sediment, Yangtze Delta. *Journal of Coastal Research* 25:915–924
- Maza M, Lara JL, Losada IJ, Ondiviela B, Trinogga J, Bouma TJ (2015) Large-scale 3-D experiments of wave and current interaction with real vegetation. Part 2: experimental analysis. *Coastal Engineering* 106:73–86
- Mendez FJ, Losada IJ (2004) An empirical model to estimate the propagation of random breaking and nonbreaking waves over vegetation fields. *Coastal Engineering* 51:103–118
- Möller I (2006) Quantifying saltmarsh vegetation and its effect on wave height dissipation: results from a UK east coast saltmarsh. *Estuarine, Coastal and Shelf Science* 69:337–351
- Möller I, Spencer T, French JR, Leggett DJ, Dixon M (1999) Wave transformation over salt marshes: a field and numerical modelling study from North Norfolk, England. *Estuarine, Coastal and Shelf Science* 49:411–426
- Möller I, Kudella M, Rupprecht F, Spencer T, Paul M, van Wesenbeeck BK, Wolters G, Jensen K, Bouma TJ, Miranda-Lange M, Schimmels S (2014) Wave attenuation over coastal salt marshes under storm surge conditions. *Nature Geoscience* 7:727–731
- Morris JT, Sundareshwar PV, Netch CT, Kjerfve B, Cahoon DR (2002) Responses of coastal wetlands to rising sea level. *Ecology* 83:2869–2877
- Nepf HM (2011) Flow over and through biota. In: Wolanski E, McLusky D (eds) *Treatise on estuarine and coastal science*. Academic, Waltham, Massachusetts, pp 267–288
- Nolte S, Koppenaal EC, Esselink P, Dijkema KS, Schuerch M, de Groot AV, Bakker JP, Temmerman S (2013) Measuring sedimentation in tidal marshes: a review on methods and their applicability in biogeomorphological studies. *Journal of Coastal Conservation* 17: 301–325
- Peruzzo P, De Serio F, Defina A, Mossa M (2018) Wave height attenuation and flow resistance due to emergent or near-emergent vegetation. *Water* 10:402
- Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team (2018) nlme: linear and nonlinear mixed effects models. R package version 3.1–137. <https://CRAN.R-project.org/package=nlme>
- Ratliff KM, Braswell AE, Marani M (2015) Spatial response of coastal marshes to increased atmospheric CO₂. *Proceedings of the National Academy of Sciences of the United States of America* 112:15580–15584
- Redfield AC (1965) Ontogeny of a salt marsh estuary. *Science* 147:50–55
- Reed DJ (1989) Patterns of sediment deposition in subsiding coastal salt marshes, Terrebonne Bay, Louisiana: the role of winter storms. *Estuaries* 12:222–227
- Reed DJ, de Luca N, Foote AL (1997) Effect of hydrologic management on marsh surface sediment deposition in coastal Louisiana. *Estuaries* 20:301–311
- Reed DJ, Spencer T, Murray AL, French JR, Leonard L (1999) Marsh surface sediment deposition and the role of tidal creeks: implications for created and managed coastal marshes. *Journal of Coastal Conservation* 5:81–90
- Reef R, Schuerch M, Christie EK, Möller I, Spencer T (2018) The effect of vegetation height and biomass on the sediment budget of a European saltmarsh. *Estuarine, Coastal and Shelf Science* 202: 125–133
- Reitsma JM, de Jong J (2019) Toelichting bij de vegetatiekartering Kwelders Texel 2017: Op basis van false colour-luchtfoto's 1: 5.000. Rijkswaterstaat, Delft
- Roelvink JA (1993) Dissipation in random wave groups incident on a beach. *Coastal Engineering* 19:127–150
- Roelvink D, Reniers A, van Dongeren A, van Thiel de Vries J, McCall R, Lescinski J (2009) Modelling storm impacts on beaches, dunes and barrier islands. *Coastal Engineering* 56:1133–1152
- Schepers L, van Braeckel A, Bouma TJ, Temmerman S (2020) How progressive vegetation die-off in a tidal marsh would affect flow and sedimentation patterns: a field demonstration. *Limnology and Oceanography* 65:401–412
- Shepard CC, Crain CM, Beck MW (2011) The protective role of coastal marshes: a systematic review and meta-analysis. *PLoS One* 6: e27374
- Shi BW, Yang SL, Wang YP, Bouma TJ, Zhu Q (2012) Relating accretion and erosion at an exposed tidal wetland to the bottom shear stress of combined current – wave action. *Geomorphology* 138: 380–389
- Silinski A, Fransen E, Bouma TJ, Meire P, Temmerman S (2016) Unravelling the controls of lateral expansion and elevation change of pioneer tidal marshes. *Geomorphology* 274:106–115
- Silva H, Dias JM, Caçador I (2009) Is the salt marsh vegetation a determining factor in the sedimentation processes? *Hydrobiologia* 621: 33–47
- Stocker TF, Qin D, Plattner G-K, Tignor MMB, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM (2013) *Climate change 2013: the physical science basis, intergovernmental panel on climate change, working group I contribution to the IPCC fifth assessment report (AR5)*. Cambridge University Press, New York
- Suzuki T, Zijlema M, Burger B, Meijer MC, Narayan S (2012) Wave dissipation by vegetation with layer schematization in SWAN. *Coastal Engineering* 59:64–71
- Temmerman S, Govers G, Wartel S, Meire P (2003) Spatial and temporal factors controlling short-term sedimentation in a salt and freshwater tidal marsh, Scheldt estuary, Belgium, SW Netherlands. *Earth Surface Processes and Landforms* 28:739–755
- Temmerman S, Bouma TJ, Govers G, Wang ZB, de Vries MB, Herman PMJ (2005) Impact of vegetation on flow routing and sedimentation patterns: three-dimensional modeling for a tidal marsh. *Journal of Geophysical Research - Earth Surface* 110:F04019

- Temmerman S, Moonen P, Schoelynck J, Govers G, Bouma TJ (2012) Impact of vegetation die-off on spatial flow patterns over a tidal marsh. *Geophysical Research Letters* 39:L03406
- Temmerman S, Meire P, Bouma TJ, Herman PMJ, Ysebaert T, de Vriend HJ (2013) Ecosystem-based coastal defence in the face of global change. *Nature* 504:79–83
- Torio DD, Chmura GL (2013) Assessing coastal squeeze of tidal wetlands. *Journal of Coastal Conservation* 29:1049–1061
- Townend I, Fletcher C, Knappen M, Rossington K (2011) A review of salt marsh dynamics. *Water Environment Journal* 25:477–488
- van der Meijden R (2005) *Heukels' Flora van Nederland*, 23rd edn. Noordhoff Uitgevers, Groningen/Houten, the Netherlands
- van Loon-Steensma JM, Slim PA, Decuyper M, Hu Z (2014) Salt-marsh erosion and restoration in relation to flood protection on the Wadden Sea barrier island Terschelling. *Journal of Coastal Conservation* 18: 415–430
- van Loon-Steensma JM, Hu Z, Slim PA (2016) Modelled impact of vegetation heterogeneity and salt-marsh zonation on wave damping. *Journal of Coastal Research* 32:241–252
- van Rooijen AA, McCall RT, van Thiel de Vries JSM, van Dongeren AR, Reniers AJHM, Roelvink JA (2016) Modeling the effect of wave-vegetation interaction on wave setup. *Journal of Geophysical Research, Oceans* 121:4341–4359
- van Veelen TJ, Fairchild TP, Reeve DE, Karunaratna H (2020) Experimental study on vegetation flexibility as control parameter for wave damping and velocity structure. *Coastal Engineering* 157:103648
- Vuik V, Jonkman SN, Borsje BW, Suzuki T (2016) Nature-based flood protection: the efficiency of vegetated foreshores for reducing wave loads on coastal dikes. *Coastal Engineering* 116:42–56
- Wang A, Gao S, Jia J (2006) Impact of the cord-grass *Spartina alterniflora* on sedimentary and morphological evolution of tidal salt marshes on the Jiangsu coast, China. *Acta Oceanologica Sinica* 25:32–42
- Watson EB, Oczkowski AJ, Wigand C, Hanson AR, Davey EW, Crosby SC, Johnson RL, Andrews HM (2014) Nutrient enrichment and precipitation changes do not enhance resiliency of salt marshes to sea level rise in the northeastern U.S. *Climatic Change* 125:501–509
- Wood N, Hine AC (2007) Spatial trends in marsh sediment deposition within a microtidal creek system, Waccasassa Bay, Florida. *Journal of Coastal Research* 23:823–833
- Yang SL (1999a) Sedimentation on a growing intertidal island in the Yangtze River mouth. *Estuarine, Coastal and Shelf Science* 49: 401–410
- Yang SL (1999b) Tidal wetland sedimentation in the Yangtze Delta. *Journal of Coastal Research* 15:1091–1099
- Yang SL, Eisma D, Ding PX (2000) Sedimentary processes on an estuarine marsh island within the turbidity maximum zone of the Yangtze River mouth. *Geo-Marine Letters* 20:87–92
- Yang SL, Li H, Ysebaert T, Bouma TJ, Zhang WX, Wang YY, Li P, Li M, Ding PX (2008) Spatial and temporal variations in sediment grain size in tidal wetlands, Yangtze Delta: on the role of physical and biotic controls. *Estuarine, Coastal and Shelf Science* 77:657–671
- Yang SL, Shi BW, Bouma TJ, Ysebaert T, Luo XX (2012) Wave attenuation at a salt marsh margin: a case study of an exposed coast on the Yangtze estuary. *Estuaries and Coasts* 35:169–182
- Ysebaert T, Yang SL, Zhang L, He Q, Bouma TJ, Herman PMJ (2011) Wave attenuation by two contrasting ecosystem engineering salt marsh macrophytes in the intertidal pioneer zone. *Wetlands* 31: 1043–1054

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.