

Erratum to: ACMT Position Statement: Guidance for the Use of Intravenous Lipid Emulsion

American College of Medical Toxicology

Received: 3 May 2016 / Accepted: 3 May 2016 / Published online: 25 May 2016
© American College of Medical Toxicology 2016

Erratum to: J. Med. Toxicol.
DOI 10.1007/s13181-016-0550-z

On page 1, the fourth paragraph should read:

“There are no validated, evidence-based dosing regimens. The American Heart Association described a 1.5 mL/kg bolus followed by a 0.25 mL/kg/min infusion, continued for 30–60 minutes to a maximum infusion of 10 mL/kg [11]. *Lipidrescue.org* recommends repeating this initial bolus one to two times for persistent “cardiovascular collapse,” raising the infusion rate to 0.5 mL/kg/min for persistent hypotension, and limiting the total dose to 10–12 mL/kg over the first 30 minutes [2]. Additional dosing recommendations include a loading dose of 1.5 mL/kg, followed by 3–5 minutes of infusion at 0.25 mL/kg, and then a maintenance infusion of 0.025 mL/kg/min [9]. This lower infusion rate may be sufficient to maintain the positive effects of lipids while avoiding lipid overload.”

The dose is correctly stated in the **Recommended Guideline** section on page 2.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s13181-016-0550-z>.

✉ American College of Medical Toxicology
positionstatements@acmt.net

¹ ACMT, Phoenix, AZ, USA