



Correction to: Clinical Impact of Inferior Mesenteric Lymph Node Metastasis in Patients with Cancer of the Sigmoid Colon or Rectum

Rikiya Sato¹ · Shuichiro Matoba² · Hiroya Kuroyanagi² · Masashi Ueno² · Shigeo Toda² · Yutaka Hanaoka² · Kosuke Hiramatsu² · Yusuke Maeda² · Yuki Nishihara²

Published online: 8 September 2021
© Indian Association of Surgical Oncology 2021

Correction to: Indian Journal of Surgical Oncology
<https://doi.org/10.1007/s13193-021-01389-3>

The original version of this article contained mistakes.

- The corresponding author is affiliated to affiliation 2.
- In the first paragraph of the “Discussion” section, the text “, which has made it difficult to demonstrate the prognostic impact of IMLN dissection for patients with IMLN metastasis. In terms of long-term outcomes, 5-year RFS has previously been reported to range from 18.4 to 38.5%, and 5-year OS has been reported to range from 14 to 40% [1,5–7]. Ubukata et al. [11]” was inadvertently deleted.

This is being corrected in this publication.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s13193-021-01389-3>.

✉ Shuichiro Matoba
s-matoba@ac.cyberhome.ne.jp

¹ Department of General and Gastroenterological Surgery, Nanpuh Hospital, 14-3, Nagata-cho, Kagoshima-shi, Kagoshima 892-8512, Japan

² Department of Gastroenterological Surgery, Toranomon Hospital, 2-2-2 Toranomon, Minato-ku, Tokyo 105-8470, Japan