

The recombinant pea defensin Drr230a is active against impacting soybean and cotton pathogenic fungi from the genera *Fusarium*, *Colletotrichum* and *Phakopsora*

Ariane Ferreira Lacerda^{1,2} · Rafael Perseghini Del Sarto^{1,3} ·
Marilia Santos Silva¹ · Erico Augusto Rosas de Vasconcelos¹ ·
Roberta Ramos Coelho¹ · Vanessa Olinto dos Santos^{1,2} ·
Claudia Vieira Godoy⁴ · Claudine Dinali Santos Seixas⁴ ·
Maria Cristina Mattar da Silva¹ · Maria Fatima Grossi-de-Sa^{1,5}

Received: 13 April 2015 / Accepted: 13 June 2015 / Published online: 13 February 2016
© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract Plant defensins are antifungal peptides produced by the innate immune system plants developed to circumvent fungal infection. The defensin Drr230a, originally isolated from pea, has been previously shown to be active against various entomopathogenic and phytopathogenic fungi. In the present study, the activity of a yeast-expressed recombinant Drr230a protein (rDrr230a) was tested against impacting soybean and cotton fungi. First, the gene was subcloned into the yeast expression vector pPICZαA and expressed in *Pichia pastoris*. Resulting rDrr230a exhibited in vitro activity against fungal growth and spore germination of *Fusarium tucumaniae*, which causes soybean sudden death syndrome, and against *Colletotrichum gossypii* var. *cephalosporioides*, which causes cotton ramulosis. The rDrr230a IC₅₀ corresponding to inhibition of fungal growth of *F. tucumaniae* and *C. gossypii* var. *cephalosporioides* was 7.67 and 0.84 μM, respectively, demonstrating moderate activity against *F. tucumaniae* and high potency against *C. gossypii* var.

cephalosporioides. Additionally, rDrr230a at 25 ng/μl (3.83 μM) resulted in 100 % inhibition of spore germination of both fungi, demonstrating that rDrr230a affects fungal development since spore germination. Moreover, rDrr230a at 3 μg/μl (460.12 μM) inhibited 100 % of in vitro spore germination of the obligatory biotrophic fungus *Phakopsora pachyrhizi*, which causes Asian soybean rust. Interestingly, rDrr230a substantially decreased the severity of Asian rust, as demonstrated by *in planta* assay. To our knowledge, this is the first report of a plant defensin active against an obligatory biotrophic phytopathogenic fungus. Results revealed the potential of rDrr230a as a candidate to be used in plant genetic engineering to control relevant cotton and soybean fungal diseases.

Keywords Defensin · *Pisum sativum* · *Pichia pastoris* · *Fusarium tucumaniae* · *Colletotrichum gossypii* var. *cephalosporioides* · *Phakopsora pachyrhizi*

Introduction

Plant diseases caused by fungi affect a broad range of crops worldwide, being responsible for significant losses and for the decrease in quality and safety of agricultural products. In crop plants, fungi cause more economic damage than any other group of microorganisms, with annual losses estimated at more than US\$ 200 billion (Horbach et al. 2011). There are several fungal diseases impacting in tropical areas, which include cotton ramulosis, Asian soybean rust and soybean sudden death syndrome caused by *Colletotrichum gossypii* (South) var. *cephalosporioides* A. S. Costa, *Phakopsora pachyrhizi* H. & P. Syd., and *Fusarium solani* (Mart.) Sacc. f. sp. *glycines* (= *Fusarium tucumaniae* T.) (Aoki et al. 2003), respectively. Currently,

✉ Maria Fatima Grossi-de-Sa
fatima.grossi@embrapa.br

¹ Embrapa Recursos Genéticos e Biotecnologia, PqEB, Avenida W5 Norte (Final), PO 02372, Brasília, DF 70770-917, Brazil
² Universidade Federal do Rio Grande do Norte, Avenida Senador Salgado Filho 3000, Lagoa Nova, Natal, RN 59078-970, Brazil
³ Centro Universitário do Distrito Federal, SEP/SUL EQ 704/ A12 904 Conjunto A, Brasília, DF 70390-045, Brazil
⁴ Embrapa Soja, Rodovia Carlos João Strass, Distrito de Warta, A10 PO Box 231, Distrito de Warta, PR 86001-970, Brazil
⁵ Universidade Católica de Brasília, SGAN 916 Módulo B Avenida W5, Brasília, DF 70790-160, Brazil

there are no cotton or soybean varieties with good agronomic traits, which at the same time present high resistance against these important diseases. In addition, none of these diseases are satisfactorily controlled by fungicides.

To circumvent fungal infection, plants have developed innate immune systems that recognize the presence of pathogens and initiate effective defense responses (Muthamilarasan and Prasad 2013). Therefore, plants produce diverse molecules such as carbohydrates, lipids (Rojas et al. 2014), antimicrobial enzymes (Habib and Fazili 2007), secondary metabolites, (Ahuja et al. 2012; Bednarek 2012) and several defense-related proteins such as chitinases, glucanases, lectins, thionins, and defensins (Carvalho and Gomes 2009).

Defensins, which are pathogenesis-related proteins produced by plants, invertebrate and vertebrate animals, are basic, small (around 50 amino acids, about 5 kDa), cysteine-rich-peptides whose three-dimensional structure presents one α -helix and three antiparallel β -sheets stabilized by an $\alpha\beta$ motif (Lay and Anderson 2005; Carvalho and Gomes 2009). Defensins have a broad range of biological activity, including inhibition of α -amylases (Bloch and Richardson 1991; Pelegrini et al. 2008) and proteases (Wijaya et al. 2000; Melo et al. 2002), antimicrobial activity (Ye and Ng 2001; Wong and Ng 2005; de Zélicourt et al. 2007), inhibition of protein synthesis (Mendez et al. 1990, 1996), blockage of ionic channels (Spelbrink et al. 2004) and interference in cell cycle by DNA binding (Lobo et al. 2007).

Transgenic plants expressing defensins have been shown to be an alternative strategy to protect plants against pathogens (De Bondt et al. 1999; Gao et al. 2000; Parashina et al. 2000; Chen et al. 2002, 2006; Park et al. 2002; Chan et al. 2005; Zhu et al. 2007; Anuradha et al. 2008; Choy et al. 2009; Kovalskaya and Hammond 2009; Abdallah et al. 2010; Jha and Chattoo 2010) specially in case there is no source of natural resistance for breeding programs.

The pea (*Pisum sativum*) defensin Drr230a, transgenically expressed in tobacco, has been previously shown to present antimicrobial activity against the entomopathogenic fungus *Trichoderma reesei* and against the phytopathogenic fungi *F. solani*, *F. oxysporum*, *Ascochyta pisi*, *A. pinodes*, *A. lentis*, *Alternaria alternata*, and *Lep-tosphaeria maculans* (Lai et al. 2002).

Considering the potential of Drr230a to control other relevant fungal diseases in agriculture, in the present study, we describe the expression of the *drr230a* gene in *Pichia pastoris*, the purification of the resulting recombinant Drr230a (rDrr230a) and evaluations of its activity against the impacting cotton and soybean fungi *C. gossypii* var. *cephalosporioides*, *P. pachyrhizi* and *F. tucumaniae*.

Materials and methods

Phytopathogenic fungi

The fungal pathogen *F. solani* (Mart.) Sacc. f. sp. *glycines* (= *F. tucumaniae*), which causes soybean sudden death syndrome, was provided by Dr. Leila Costamilan (Embrapa Trigo, Passo Fundo-RS, Brazil). *C. gossypii* (South) var. *cephalosporioides* A. S. Costa, a fungal pathogen that causes cotton ramulosis, was provided by Dr. Nelson Dias Suassuna (Embrapa Algodão, Campina Grande-PB, Brazil). Both fungi were maintained by in vitro cultivation and spore lyophilization.

The *P. pachyrhizi* H. & P. Syd. uredospores, which cause Asian soybean rust, were collected at soybean fields in Londrina-PR, Brazil and maintained in soybean plants under greenhouse controlled conditions at Embrapa Soja, Londrina-PR, Brazil.

Subcloning of *drr230a* gene into yeast expression vector

The *drr230a* defensin gene (Genebank accession AF525685), provided by Dr. Richard Fobert (National Research Council Canada, Plant Biotechnology Institute), was originally isolated from pea (*P. sativum*) and previously demonstrated to present antifungal activity (Lai et al. 2002). The *drr230a* gene, initially subcloned into a plant transformation vector, provided by Dr Richard Fobert, was amplified by using the primers DRR230a01For (5' GCC GAA TTC AAC ACA TGT GAG AAC 3') and DRR230a02Rev (5' TGG GCG GCC GCT CAA TGA TGA TGA TGA TGA TGG CAG TTT TTA GTA CAC CAA CAG CGA AAG TCA TC 3') designed to insert *EcoRI* and *NotI* restriction sites at the 5' and 3' ends, respectively, to subclone into the *P. pastoris* expression vector pPICZ α [®] (Invitrogen Co.). Moreover, by using the reverse primer DRR230a02Rev a coding sequence of six histidines (6xHis) was added to the 3' end of the *drr230a* gene, to enable purification of the corresponding recombinant rDrr230a protein by immobilized metal affinity chromatography (IMAC) and immunodetection with commercial anti-His antibodies. The amplicon comprising the *drr230a* gene with 6xHis-tag was subcloned into the *EcoRI* and *NotI* sites of pPICZ α [®] in frame with the α -factor secretion peptide signal, downstream of the alcohol oxidase I promoter.

The resulting construct pPICZ α -Drr230a integrity was confirmed by automated ABI sequencing (Perkim-Elmer). The pPICZ α -Drr230a construct was used to transform *P. pastoris* competent cells strain X-33 by using 20 μ g of *SacI* linearized DNA via electroporation (in accordance with the manufacturer). Clones were selected on YPD broth (1 %

yeast extract; 2 % dextrose; 2 % peptone, 2 % agar) plates containing zeocin (100 µg/ml) and by PCR analyses. Clones were then cultivated in plates with higher zeocin concentration (500 µg/ml) to increase chances of selecting those with two gene copies in the yeast genome (Vassileva and Chugh 2001).

Expression of rDrr230a in *P. pastoris* and its purification by immobilized metal affinity chromatography (IMAC)

One positive clone containing the pPICZαA-Drr230a construct was inoculated in 2 ml of YPD broth (1 % yeast extract; 2 % dextrose; 2 % peptone) and incubated over night at 28 °C and 200 rpm (i.e. pre-inoculum). Afterwards, the 2 ml pre-inoculum was diluted in 200 ml of BMG broth (1.34 % de YNB, 1 % de glycerol, 4×10^{-5} % Biotin, 100 mM potassium phosphate buffer, pH 6.0) and incubated for 24 h to reach OD₆₀₀ = 20 (i.e. inoculum). Then, those 200 ml inoculum were centrifuged (1500×g; 25 °C; 5 min) and the resulting pellet was resuspended in the same volume of BMM broth (1.34 % de YNB, 1 % methanol, 4×10^{-5} % Biotin, 100 mM potassium phosphate buffer, pH 6.0) to induce expression of the recombinant protein rDrr230a. During the induced expression, the culture was incubated at 28 °C and 220 rpm agitation for 4 days. Aliquots of the culture supernatant were analyzed by SDS-TRICINE-PAGE 14 % and Western Blot probed with commercial anti-His antibodies (Invitrogen Co.) to detect the His-tag rDrr230a protein (Towbin et al. 1979; Schagger and Von Jagow 1987).

Supernatant from the expression culture was filtered through 0.2 µm membranes (Millex-GV, Millipore), diluted in binding buffer (100 mM sodium phosphate buffer, pH 7.4; 500 mM NaCl) and applied into a His-Trap FF crude column (1.6 × 2.5 cm; 5 ml) (GE Helthcare) previously equilibrated with binding buffer pH 7.4 at 5 ml/min. Binding recombinant rDrr230a protein was eluted with elution buffer (binding buffer containing imidazole to final concentration of 500 mM) and checked for purity by SDS-TRICINE-PAGE 14 % and Western Blot probed with commercial anti-His antibodies (Invitrogen Co.). The fraction containing the recombinant rDrr230a protein was dialyzed, lyophilized and dissolved in deionized water. The N-terminal amino acid sequence of rDrr230a was confirmed by sequencing using Edman's degradation technique.

Antifungal bioassay against non-biotrophic phytopathogenic fungi

Bioassays against the soybean pathogenic fungus *F. tucumaniae* and against the cotton pathogenic fungus *C. gossypii* var. *cephalosporioides* were performed in two

quantitative assays. The first assay, performed to analyze inhibition of fungal growth, consisted of incubating 10 µl of a solution containing 10⁴ spores/ml with 80 µl of liquid PDB (Potato Dextrose Broth) and rDrr230a to different final concentrations, varying from 0.2 up to 60 ng/µl in the case of inhibition of *F. tucumaniae* growth, and varying from 5 up to 50 ng/µl in the case of inhibition of *C. gossypii* var. *cephalosporioides* growth (assay total volume corresponded to 100 µl per sample). The assay was done in a 96-well microplate and the samples were incubated at 28 °C for 48 h in the dark. The inhibition of fungal growth was measured by optical density in a microplate reader (BIORAD) at 600 nm. All treatments were done in triplicates. Distilled water and hydrogen peroxide (70 µM) were used as negative and positive controls, respectively.

The second assay, performed to analyze inhibition of spore germination, was carried out using 5 µl of a solution containing 10⁴ spores/ml and 15 µl of a solution containing rDrr230a. The samples were incubated at 28 °C for 12 h. The germinated spores were counted in a Neubauer chamber upon observation under an optical microscope (Olympus Optical Co.) within several fields of 100 spores/field. Moreover, representative sample fields were photographed with the same equipment. All treatments were done in triplicates. Distilled water and hydrogen peroxide (70 µM) were used as negative and positive controls, respectively.

Antifungal bioassay against obligatory biotrophic phytopathogenic fungus

Bioassays against the obligatory biotrophic soybean pathogenic fungus *P. pachyrhizi* were performed in two assays. The first assay was performed to analyze the disease severity caused *in planta* by *P. pachyrhizi* in presence or absence of rDrr230a. This assay was carried out by inoculating half soybean detached leaflets with 150 µl of a solution containing 8×10^4 uredospores/ml (in sterile distilled water containing 0.01 % v/v Tween 20), by brushing. Fourteen minutes afterwards, the same half leaflets were treated, by brushing, with rDrr230a to final concentration 3 µg/µl. The leaflets were then incubated at 24 °C for 14 days in a humid chamber. The severity was evaluated daily. All treatments were done in triplicates. Distilled water was used as the negative control. In addition, 24 h post inoculation, one half leaflet of each treatment was punched with a microtube cap to collect samples for microscopy analyses of uredospore germination under a Scanning Electron Microscope. For this reason, each collected leaf disk was immediately fixed in glutaraldehyde/paraformaldehyde solution. Afterwards, the leaf disks were further fixed in osmium tetroxide and then in uranyl acetate. Dehydration of the leaf disks was performed in

acetone solutions of serially increasing concentrations. Acetone was removed from the leaf disk samples by critical point drying. Finally, the leaf disks were coated with gold by low-vacuum sputter coating, then observed and photographed under Scanning Electron Microscope (Jeol).

The second assay, performed to analyze inhibition of uredospore germination, was carried out by using 50 μl of a solution containing 10^4 uredospores/ml (in sterile distilled water containing 0.01 % v/v Tween 20) and rDrr230a at final concentration of 3 $\mu\text{g}/\mu\text{l}$, in a final volume of 100 μl . The assay was done in a 96-well microplate and the samples were incubated at 24 °C in a humid chamber overnight. To end up the assay, lactophenol was added. The germinated uredospores were counted in a Neubauer chamber upon observation with an optical microscope (Olympus Optical Co) within several fields of 100 uredospores/field. Moreover, representative sample fields were photographed with the same equipment. All assays were done in triplicates. Distilled water was used as negative control of inhibition of uredospore germination.

Results and discussion

Expression of rDrr230a in *P. pastoris* and its purification by immobilized metal affinity chromatography (IMAC)

To evaluate the biological activity of rDrr230a against phytopathogenic fungi, the corresponding gene was introduced into the *P. pastoris* expression vector pPICZ α A and the rDrr230a protein was expressed under control of the pAOX1 methanol-inducible promoter (Fig. 1). The resulting recombinant protein rDrr230a was secreted due to the α -mating factor signal peptide present in the vector (Fig. 1). A poly-histidine tag (6xHis) was added to the 3' end of the gene sequence to enable purification of the resulting recombinant His-tagged protein (Fig. 1) through IMAC. *P. pastoris* strain X-33 was transformed with the expression construct and resulting colonies were assayed in plates with high concentration of the selection agent zeocin. Zeocin resistant clones were used in a small-scale expression assay. The clone that presented the highest expression level was chosen to proceed to large-scale production of rDrr230a.

The recombinant protein rDrr230a was expressed upon promoter induction with methanol, and untransformed cells were used as negative control sample. Aliquots from the expression set up were collected every 24 h in a time course from 24 h up to 96 h, and analyzed in TRICINE-SDS-PAGE. The production of rDrr230a occurred at 24 h, remaining until 96 h, and rDrr230a presented the expected molecular size of 6.52 kDa (Fig. 2a, b).

The supernatant, containing the secreted rDrr230a, was directly submitted to purification by IMAC. The eluted fractions were analyzed by TRICINE-SDS-PAGE 14 % and Western blot probed with anti-His antibody, confirming the expression of a recombinant protein comprising 6.52 kDa (Fig. 3a, b). The N-terminal amino acid sequence of rDrr230a was confirmed by Edman's degradation sequencing, which revealed additional amino acids corresponding to leftovers of the Kex2 cleavage site (EAEA) and *Eco*RI site (EF) at the N-terminus (Fig. 1). Considering as well the prediction of amino acid sequence based on nucleotide sequence, the resulting protein sequence of the His-tagged recombinant rDrr230a (57 residues) is EAEA EF NTC ENL AGS YKG VCF GGC DRH CRT QEG AIS GRC RDD FRC WCT KNC HHHHHH, as shown in Fig. 1. Quantification of purified rDrr230a, done by 2D-Quant method (Amersham), indicated a recombinant protein recovery estimated in 3.5 mg/l.

rDrr230a activity against non-biotrophic and obligatory biotrophic phytopathogenic fungi

To analyze inhibition of fungal growth upon incubation with the defensin, spores of the non-biotrophic fungi *F. tucumaniae* and *C. gossypii* var. *cephalosporioides* were incubated with rDrr230a in microplate wells in liquid medium for 48 h. This bioassay was performed by using a concentration range of rDrr230a varying from 0.2 ng/ μl up to 60 ng/ μl (0.03 μM up to 9.20 μM) to analyze inhibition of *F. tucumaniae* growth and a range varying from 5 up to 50 ng/ μl (0.77 μM up to 7.67 μM) in the case of *C. gossypii* var. *cephalosporioides* (Fig. 4). Based on the resulting curves of inhibition of fungal growth of both fungi upon incubation with rDrr230a, the calculated IC₅₀ against *F. tucumaniae* was 7.67 μM (50 ng/ μl ; Fig. 4a) and against *C. gossypii* var. *cephalosporioides* corresponded to 0.84 μM (5.5 ng/ μl ; Fig. 4b), indicating that rDrr230a is moderately efficient against *F. tucumaniae* and highly potent against *C. gossypii* var. *cephalosporioides*.

To further investigate the deleterious effect of rDrr230a upon both fungi, the activity of rDrr230a against spore germination of *F. tucumaniae* and *C. gossypii* var. *cephalosporioides* was tested at 25 ng/ μl (3.83 μM) 12 h post incubation, in a similar microplate setup as previously described. At concentration 25 ng/ μl (3.83 μM) the spore germination of both *F. tucumaniae* and *C. gossypii* var. *cephalosporioides* was inhibited in 100 % (Fig. 5a, c, e) despite the fact that the spores of both fungi germinated normally under incubation in water (Fig. 5a, b, d). These observations demonstrate that rDrr230a affects the fungal development of *F. tucumaniae* and *C. gossypii* var. *cephalosporioides* from the very beginning, i.e. since the spore germination.




Fig. 1 Schematic description of the expression cassette construct designed to express Histidine-tagged recombinant Drr230a in yeast (*P. pastoris*). The *drr230a* gene was subcloned into the yeast expression vector pPICZ α A (Invitrogen Co.) under the control of the alcohol oxidase 1 promoter (pAOX1) inducible by methanol. The orientations of the primers DRR230a01For and DRR230a02Rev, including the respective restriction *Eco*RI (resulting in addition of EF residues at the recombinant protein N-terminus) and *Not*I cloning sites (not expressed), are indicated by arrows. The recombinant

protein (sequence depicted above) is expressed in frame with the *Saccharomyces cerevisiae* secretion mating α -factor signal peptide (α -Factor), which is cleaved before protein export at the Kex2 signal cleavage site (Kex2), resulting in the addition of four residues (EAEA) at the N-terminal end of the recombinant protein. A Histidine tag (6xHis; HHHHHH), followed by a stop codon (asterisk) was included at the C-terminal end of the sequence to facilitate recombinant protein purification by Immobilized metal affinity chromatography




Fig. 2 Expression of rDrr230a in *P. pastoris* X-33 cells. **a** SDS-TRICINE-PAGE analysis of the expression of the His-tagged rDrr230a protein from 24 to 96 h after methanol-induction. **b** Western blot analysis of expressed His-tagged rDrr230a protein probed with anti-His antibody (Invitrogen). Molecular mass marker See Blue[®]Plus 2 Prestained Standard, Invitrogen (lane M). Arrows indicate bands corresponding to the expressed rDrr230a (6.52 kDa)

Supernatant from *P. pastoris* X-33 culture expressing rDrr230a 24 h (lane 1), 48 h (lane 2), 72 h (lane 3) and 96 h (lane 4) post-induction. Supernatant from *P. pastoris* X-33 untransformed culture 24 h (lane 5), 48 h (lane 6), 72 h (lane 7) and 96 h (lane 8) post-induction. Arrows indicate bands corresponding to the expressed rDrr230a (6.52 kDa)

The *P. pastoris* system was suitable to produce active rDrr230a (Figs. 2, 3) as shown by biological activity of the resulting recombinant protein against *F. tucumaniae* and *C. gossypii* var. *cephalosporioides* (Fig. 4). There are several other reports corroborating that yeast expression systems are adequate to produce active antimicrobial peptides rich in sulfide-bridges, such as defensins. For instance, *P. pastoris* yeast expression systems were successfully used to produce active plant defensins such as the corn defensin PDC1 (Kant et al. 2009), the dimeric defensin SPE10 (Song et al. 2005), the pea defensin Psd1 (Almeida et al. 2001), the sugar beet defensin AX2 (Kristensen et al. 1999), the mungbean defensin VrD1 (Jrchen et al. 2004) and the wild tobacco defensin NmDef02 (Portieles et al. 2010).

Besides expression of plant defensins in yeast systems, there are reports as well of expression of plant defensins in bacteria, in all cases resulting in production of bioactive peptides against various fungi, such as the wild indigo

defensin TvD1 (Vijayan et al. 2008), the pepper defensin J1-1 (Seo et al. 2014) and the corn defensin PDC1 (Kant et al. 2009). Nevertheless, when expressed in *E. coli* as compared to the *P. pastoris* system, PDC1 expressed in yeast was more efficient to inhibit growth of *F. graminearum* than the same protein expressed in bacteria (Kant et al. 2009), as it is expected, since yeast is an eukaryotic system, what favors correct cysteine-bond formation, protein correct folding and consequently protein full activity (Demain and Vaishnav 2009; Fickers 2014).

Concerning recombinant plant defensins tested against phytopathogenic fungi from the *Colletotrichum* genus, Seo et al. (2014) demonstrated activity of the defensin J1-1, expressed in bacteria, against *C. gloeosporioides* (IC₅₀ 8.3 μ M). The defensin SPE10, expressed in yeast, also presented antifungal activity when tested towards the same fungal species (Song et al. 2005), i.e. *C. gloeosporioides*, despite being less potent than J1-1 (IC₅₀ >20 μ M).




Fig. 3 Purification of rDrr230a expressed in *P. pastoris* X-33 cells. rDrr230a was purified by immobilized-metal affinity chromatography (IMAC). **a** SDS-TRICINE-PAGE analysis of the purification of rDrr230a. **b** Western blot analysis of purification of rDrr230a, probed with anti-His antibody (Invitrogen). Molecular mass marker See Blue[®]Plus 2 Prestained Standard, Invitrogen (lane M). Supernatant

from *P. pastoris* X-33 culture expressing rDrr230a 96 h after methanol-induction (lane 1). Eluate fraction from IMAC, corresponding to *P. pastoris*-expressed rDrr230a 96 h after methanol-induction (lane 2). Arrows indicate bands corresponding to the purified rDrr230a (6.52 kDa)





Fig. 4 Inhibition of fungal growth of the non-biotrophic phytopathogenic fungi *F. tucumaniae* and *C. gossypii* var. *cephalosporioides* by rDrr230a. Bioassays were performed in microplate wells in liquid medium for 48 h. The data are mean \pm standard deviation ($n = 3$). **a** Curve of inhibition of *F. tucumaniae* growth upon incubation with rDrr230a in a concentration range varying from



0.2 ng/ μ l up to 60 ng/ μ l (0.03 μ M up to 9.20 μ M). Calculated IC_{50} against *F. tucumaniae* corresponds to 7.67 μ M (50 ng/ μ l). **b** Curve of inhibition of *C. gossypii* var. *cephalosporioides* growth upon incubation with rDrr230a in a concentration range varying from 5 up to 50 ng/ μ l (0.77 μ M up to 7.67 μ M). Calculated IC_{50} against *C. gossypii* var. *cephalosporioides* corresponds to 0.84 μ M (5.5 ng/ μ l)

Interestingly, rDrr230a is considerably more potent than J1-1 and SPE10 against a target fungus from the *Colletotrichum* genus (*C. gossypii* var. *cephalosporioides*) (IC_{50} 0.84 μ M), what represents around tenfold increase in antifungal potency. This implies that the recombinant rDrr230a expressed in yeast was correctly folded and therefore was biologically active.

In what concerns activity against the *Fusarium* genus, rDrr230a (IC_{50} 7.67 μ M) was shown to be more effective than the defensin PDC1 produced in bacteria (IC_{50} 30 μ M),

though being as effective as the same recombinant defensin expressed in yeast (IC_{50} 7.5 μ M; Kant et al. 2009). Moreover, rDrr230a (IC_{50} 7.67 μ M) was more effective than the defensin SPE10 (IC_{50} > 20 μ M), but less effective than the recombinant defensins VrD1 (IC_{50} 1.0 up to 3.4 μ M against different *F. oxysporum* isolates), NmDef02 (IC_{50} 1.0 μ M against *F. oxysporum*) and TvD1 (IC_{50} < 5.0 μ M against both *F. oxysporum* and *F. moniliforme*) against *Fusarium* species (Jrchen et al. 2004; Song et al. 2005; Vijayan et al. 2008; Portieles et al. 2010).




Fig. 5 Inhibition of spore germination of the non-biotrophic phytopathogenic fungi *F. tucumaniae* and *C. gossypii* var. *cephalosporioides* by rDrr230a. Bioassays were performed in microplate wells in liquid medium for 12 h. **a** Percentage of the inhibition of spore germination of *F. tucumaniae* and *C. gossypii* var. *cephalosporioides* upon incubation with rDrr230a. Fungal spores were incubated either with rDrr230a at 25 ng/ μ l, water or hydrogen peroxide at 70 μ M. The data in **a** are mean \pm standard deviation ($n = 3$). **b** Representative image of *F. tucumaniae* germinating spores (closed arrows) under water incubation. **c** Representative image of *F. tucumaniae* non-germinating spores (open arrows) under rDrr230a incubation. **d** Representative image of *C. gossypii* var. *cephalosporioides* germinating spores (closed arrows) under water incubation. **e** Representative image of *C. gossypii* var. *cephalosporioides* non-germinating spores (open arrows) under rDrr230a incubation

As *P. pachyrhizi* is an obligatory biotrophic phytopathogenic fungus, it is not feasible to assess rDrr230a *in vitro* activity against its growth. Therefore, the impact of rDrr230a upon *P. pachyrhizi* infection was assessed by *in planta* bioassays. For that reason, soybean detached leaflets were inoculated with *P. pachyrhizi* uredospores either in presence or absence of rDrr230a and the disease severity was

evaluated daily for 2 weeks, by using a scale varying from 0 % up to 100 % severity (Godoy et al. 2006). rDrr230a at 3 μ g/ μ l (460.12 μ M) resulted in no disease, i.e. 0 % severity, as compared to average 80 % rust severity in the absence of the recombinant protein, 14 days post inoculation (Fig. 6a, c). A leaf disk sample of each treatment was fixed at 1 day post inoculation for scanning electron microscopy analyses. The corresponding micrographs depict representative images of no uredospore germination onto well preserved plant epidermal cells when rDrr230a is present, whereas there is normal uredospore germination onto damaged epidermal plant cells in absence of rDrr230a (Fig. 5b, d). Moreover, inhibition of *P. pachyrhizi* uredospore germination was evaluated *in vitro*, under similar setup as previously evaluated for *F. tucumaniae* and *C. gossypii* var. *cephalosporioides*. In this case, rDrr230a inhibited 100 % spore germination of *P. pachyrhizi* at 3 μ g/ μ l (460.12 μ M) (Fig. 7). Despite the considerably high concentration used that inhibited *in vitro* *P. pachyrhizi* spore germination and decreased *in planta* disease severity, to our knowledge, this is the first report of a plant defensin being active against an obligatory biotrophic phytopathogenic fungus.

The antifungal effect of wild type Drr230a protein (also named as DRR206) was firstly reported by Wang and Nowak (1999) who showed that transgenic canola (*Brassica napus*)




Fig. 6 Plant disease severity (Asian soybean rust) caused by the obligatory biotrophic phytopathogenic fungus *P. pachyrhizi* upon inoculation with rDrr230a. **a**, **c** Representative half detached soybean leaflets inoculated with *P. pachyrhizi* uredospores either in presence rDrr230a at 3 μ g/ μ l (**a**) or absence of rDrr230a (**c**) 14 days post inoculation. **b**, **d** Sampled areas from inoculated detached leaflets fixed at 1 day post inoculation and analyzed by scanning electron microscopy. Representative micrographs depict no uredospore (S) germination onto well preserved plant epidermal cells when rDrr230a is present (**b**), whereas there is normal uredospore (S) germination onto damaged epidermal plant cells (yellow arrows) in absence of rDrr230a (**d**)




Fig. 7 Inhibition of spore germination of the obligatory biotrophic phytopathogenic fungus *P. pachyrhizi* by rDrr230a. Bioassays were performed in microplate wells in liquid medium for 12 h. **a** Representative image of *P. pachyrhizi* germinating spores (arrows) under water incubation. **b** Representative image of *P. pachyrhizi* non-germinating spores under incubation with rDrr230a at 3 $\mu\text{g}/\mu\text{l}$

expressing Drr230a was active against the fungus *L. maculans*, which causes blackleg disease on Brassica crops. Wang and Fristensky 2001 additionally demonstrated that Drr230a expressed in transgenic canola was effective against the phytopathogenic fungi *Rhizoctonia solani* (soil-borne pathogen that causes various plant diseases in a wide host range such as collar rot, root rot, damping off and wire stem) and *Sclerotinia sclerotiorum* (that causes white mold in a wide host range of plants).

In 2002 Lai and cols showed that leaf extracts from transgenic tobacco plants expressing Drr230a were active against the entomopathogenic fungus *T. reesei* and the phytopathogenic fungi *F. solani*, *F. oxysporum* (both *Fusarium* species that cause various diseases in plants and have extremely broad host range), *A. pisi*, *A. pinodes*, *A. lentis* (all *Ascochyta* species that cause blight in legume crops), *A. alternata* (that causes leaf spot in numerous plant hosts) and *L. maculans* (that causes blackleg on *Brassica* spp.). It is not possible to compare the potency of Drr230a against the target fungi herein tested and other affected fungi previously described because of lack of previous literature IC_{50} data available. Nevertheless, the previously reported antifungal activities of Drr230a expressed in transgenic plants is compatible with the herein presented data. Furthermore, the present data demonstrates that the recombinant rDrr230a expressed in yeast retained its antimicrobial property, interestingly against different fungal species as previously tested and with remarkable high potency against *C. gossypii* var. *cephalosporioides*.

The effect of extra residues upon antimicrobial activity of recombinant defensins is a case-wise issue. For instance, Kant et al. (2009) showed a minor deleterious effect of 6xHis-tag upon the antifungal activity of the defensin PDC1

against *F. graminearum*. Nevertheless, the activity of the defensin Psd1 against *Aspergillus niger*, a saprophytic fungus that causes post-harvest black mold in fruits and vegetables, was abolished in the presence of the four leftover residues from the Kex2 cleavage (EAEA) at the N-terminal of the recombinant protein (Cabral et al. 2003). In spite of the fact that rDrr230a expressed in yeast has extra residues, due to the Kex2 cleavage site left over (EAEA) and *EcoRI* site (EF) at the N-terminus and the presence of 6xHis-tag at the C-terminus (Fig. 1), the antifungal activity of rDrr230a was considerably retained, what is indicated by the remarkable high antimicrobial efficiency (IC_{50} 0.84 μM) against *C. gossypii* var. *cephalosporioides* (Fig. 4).

There are several reports of plant defensins transgenically expressed in plants for protection against pathogenic fungi (De Bondt et al. 1999; Gao et al. 2000; Parashina et al. 2000; Chen et al. 2002, 2006; Park et al. 2002; Chan et al. 2005; Zhu et al. 2007; Anuradha et al. 2008; Choy et al. 2009; Kovalskaya and Hammond 2009; Abdallah et al. 2010; Jha and Chattoo 2010), including the Drr230a defensin, which is object of the present study (Wang and Nowak 1999; Wang and Fristensky 2001; Lai et al. 2002). Nonetheless, to our knowledge, this is the first report of a plant defensin active against *P. pachyrhizi*, *F. tucumaniae* and *C. gossypii* var. *cephalosporioides*, which are relevant tropical phytopathogenic fungi of soybean and cotton crops. Since there is neither satisfactory chemical control nor natural sources of resistance against *P. pachyrhizi*, *F. tucumaniae* and *C. gossypii* var. *cephalosporioides*, these results point to rDrr230a as a promising tool for the development of transgenic soybean and cotton plants against this set of impacting tropical diseases.

Conclusions

The successful expression and purification of the defensin rDrr230a by using *P. pastoris* yeast system, as well as the antimicrobial activity of rDrr230a against impacting cotton and soybean pathogenic fungi was demonstrated in this article. Notably it was demonstrated the inhibition of fungal growth and spore germination of the economically important phytopathogenic fungi *F. tucumaniae*, *P. pachyrhizi* and *C. gossypii* var. *cephalosporioides*, which cause soybean sudden death syndrome, Asian soybean rust and cotton ramulosis, respectively. The data herein presented indicate moderate activity of rDrr230a against *F. tucumaniae* and high potency against *C. gossypii* var. *cephalosporioides*. rDrr230a also drastically decreased the severity of Asian soybean rust. Moreover, rDrr230a affects the development of the three target fungi tested since spore germination. These results reveal the potential of rDrr230a for plant genetic engineering to control relevant tropical fungal diseases of cotton and soybean, for which there are

neither satisfactory chemical control nor resistant varieties commercially available.

Acknowledgments The authors thank Dr. Pierre R. Fobert (National Research Council Canada, Plant Biotechnology Institute) for kindly providing the *drr230a* gene, Dr. Leila Costamilan (Embrapa Trigo) and Dr. Nelson Dias Suassuna (Embrapa Algodão) for providing fungal isolates. The authors are grateful for the scientific support, technical assistance and critical review of the manuscript performed by Dr. Thales Lima Rocha (Embrapa Recursos Genéticos e Biotecnologia). Authors thank as well Thiago de Jesus Costa (Universidade de Brasília) and João Victor Mendanha Costa (Universidade Católica de Brasília) for the technical support and Sônia Bão (Universidade de Brasília) who provided facilities at the Laboratory of Electron Microscopy. They also thank Dr. Andrew Murray (Embrapa Recursos Genéticos e Biotecnologia) for comments and suggestions on the English language review done of the manuscript. Thanks to FINEP, CNPq, CAPES and Embrapa for financial support.

Compliance with ethical standards

Conflict of interest The authors declare no financial or commercial conflict of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

Abdallah NA, Shah D, Abbas D, Madkour M (2010) Stable integration and expression of a plant defensin in tomato confers resistance to *Fusarium* wilt. *GM Crops* 1–5:344–350

Ahuja I, Kissen R, Bones AM (2012) Phytoalexins in defense against pathogens. *Trends Plant Sci* 2:73–90

Almeida MS, Cabral KS et al (2001) cDNA cloning and heterologous expression of functional cysteine-rich antifungal protein Psd1 in the yeast *Pichia pastoris*. *Arch Biochem Biophys* 395:199–207

Anuradha S, Divya K, Jami SK, Kirti PB (2008) Transgenic tobacco and peanut plants expressing a mustard defensin show resistance to fungal pathogens. *Plant Cell Rep* 27:1777–1786

Aoki T, O'donell K, Homma Y, Lattanzi AR (2003) Sudden-death syndrome of soybean is caused by two morphologically and phylogenetically distinct species within the *Fusarium solani* species complex—*F. virguliforme* in North America and *F. tucumaniae* in South America. *Mycologia* 95:660–684

Bednarek P (2012) Chemical warfare or modulators of defense responses—the function of secondary metabolites in plant immunity. *Curr Opin Plant Biol* 15:407–414

Bloch C Jr, Richardson M (1991) A new family of small (5 kDa) protein inhibitors of insect α -amylases from seeds or sorghum (*Sorghum bicolor* (L) Munch) have sequence homologies with wheat γ -purothionins. *FEBS Lett* 279:101–104

Cabral KMS, Almeida MS, Valente AP, Almeida FCL, Kurtenbach E (2003) Production of the active antifungal *Pisum sativum* defensin 1 (Psd1) in *Pichia pastoris*: overcoming the inefficiency of the STE13 protease. *Protein Expr Purif* 31:115–122

Carvalho AO, Gomes VM (2009) Plant defensins—prospects for the biological functions and biotechnological properties. *Peptides* 30:1007–1020

Chan YL, Prasad V et al (2005) Transgenic tomato plants expressing an Arabidopsis thionin (Thi2.1) driven by fruit-inactive promoter battle against phytopathogenic attack. *Planta* 221:386–393

Chen KC, Lin CY et al (2002) Cloning and characterization of a cDNA encoding an antimicrobial protein from mung bean seeds. *Bot Bull Acad Sin* 43:251–259

Chen SC, Liu AR, Zou ZR (2006) Overexpression of glucanase gene and defensin gene in transgenic tomato enhances resistance to *Ralstonia solanacearum*. *Russia J Plant Physiol* 53:756–763

Choy MS, Kim YH et al (2009) Expression of Br D1, a plant defensin from Brassica rape, confers resistance against brown plant hopper (*Nilaparvata lugens*) in transgenic rices. *Mol Cells* 28:131–137

De Bondt A, Zaman S et al (1999) Genetic transformation of apple (*Malus pumila* Mill.) for increased fungal resistance: in vitro antifungal activity in protein extracts of transgenic apple expressing Rs-AFP2 or AceAMP1. *Acta Hort* 484:565–570

de Zélicourt A, Letousey P et al (2007) Ha-DEF1, a sunflower defensin, induces cell death in Orobanchae parasitic plants. *Planta* 226:591–600

Demain AL, Vaishnav P (2009) Production of recombinant proteins by microbes and higher organisms. *Biotechnol Adv* 27:297–306

Fickers P (2014) *Pichia pastoris*: a workhorse for recombinant protein production. *Curr Res Microbiol Biotechnol* 2:354–363

Gao AG, Hakimi SM et al (2000) Fungal pathogen protection in potato by expression of a plant defensin peptide. *Nat Biotechnol* 18:1307–1310

Godoy CV, Koga LJ, Canteri MG (2006) Diagrammatic scale for assessment of soybean rust severity. *Fitopatol Bras* 31:63–68

Habib H, Fazili KM (2007) Plant protease inhibitors: a defense strategy in plants. *Biotechnol Mol Biol Rev* 2:68–85

Horbach R, Navarro-Quesada AR, Deising HB (2011) When and how to kill a plant cell: infection strategies of plant pathogenic fungi. *J Plant Physiol* 168:51–62

Jha S, Chattoo BB (2010) Expression of a plant defensin in rice confers resistance to fungal phytopathogens. *Transg Res* 19:373–384

Jrchen J, Hongchen G, Chinghsu H, Shingli S, Sanchen C (2004) Cloning and functional expression of a mungbean defensin VrD1 in *Pichia pastoris*. *Agric Food Chem* 52:2256–2261

Kant P, Liu W, Peter PK (2009) PDC1, a corn defensin peptide expressed in *Escherichia coli* and *Pichia pastoris* inhibits growth of *Fusarium graminearum*. *Peptides* 30:1593–1599

Kovalskaya N, Hammond RW (2009) Expression and functional characterization of the plant antimicrobial snakin-1 and defensin recombinant proteins. *Protein Expr Purif* 63:12–17

Kristensen AK, Brunstedt J et al (1999) Processing, disulfide pattern, and biological activity of a sugar beet defensin, AX2, expressed in *Pichia pastoris*. *Protein Expr Purif* 16:377–387

Lai FM, DeLong C, Mei K, Wignes T, Fobert PR (2002) Analysis of the DRR230 family of pea defensins: gene expression pattern and evidence of broad host-range antifungal activity. *Plant Sci* 163:855–864

Lay FT, Anderson MA (2005) Defensins—components of the innate immune system in plants. *Curr Protein Pept* 6:85–101

Lobo DS, Pereira IB et al (2007) Antifungal *Pisum sativum* defensin 1 interacts with *Neurospora crassa* cyclin F related to the cell cycle. *Biochemistry* 46:987–996

Melo FR, Rigden DJ, Franco OL, Mello LV, Ary MB, Grossi de Sa MF, Bloch C Jr (2002) Inhibition of trypsin by cowpea thionin: characterization, molecular modeling, and docking. *Proteins* 48:311–319

Mendez E, Moreno A et al (1990) Primary structure and inhibition of protein synthesis in eukaryotic cell-free system of a novel

- thionin, gamma-hordothionin, from barley endosperm. *Eur J Biochem* 194:533–539
- Mendez E, Rocher A et al (1996) Primary structure of omega-hordothionin, a member of a novel family of thionins from barley endosperm, and its inhibition of protein synthesis in eukaryotic and prokaryotic cell-free systems. *Eur J Biochem* 239:67–73
- Muthamilarasan M, Prasad M (2013) Plant innate immunity: an updated insight into defense mechanism. *J Biosci* 38:433–449
- Parashina EV, Serdobinskii LA et al (2000) Genetic engineering of oilseed rape and tomato plants expressing a radish defensin gene. *Russ J Plant Physiol* 47:417–423
- Park HC, Kang YH et al (2002) Characterization of a stamen-specific cDNA encoding a novel plant defensin in Chinese cabbage. *Plant Mol Biol* 50:57–68
- Pelegrini PB, Lay FT, Murad AM, Anderson MA, Franco OL (2008) Novel insights on the mechanism of action of α -amylase inhibitors from the plant defensin family. *Proteins* 73:719–729
- Portieles R, Ayra C et al (2010) NmDef02, a novel antimicrobial gene isolated from *Nicotiana megalosiphon* confers high-level pathogen resistance under greenhouse and field conditions. *Plant Biotechnol* 8:678–690
- Rojas CM, Senthil-Kumar M, Tzin V, Mysore KS (2014) Regulation of primary plant metabolism during plant–pathogen interaction and its contribution to plant defense. *Front Plant Sci* 10:5–17
- Schagger H, Von Jagow G (1987) Tricine-sodium dodecyl sulfate-polyacrylamide gel electrophoresis for the separation of proteins in the range from 1 to 100 kDa. *Anal Biochem* 166:368–379
- Seo HH, Park S et al (2014) Overexpression of a defensin enhances resistance to a fruit-specific anthracnose fungus in pepper. *PLoS ONE* 9:e97936
- Song X, Wang J, Wu F, Li X, Teng M, Gong W (2005) cDNA cloning, functional expression and antifungal activities of a dimeric plant defensin SPE10 from *Pachyrhizus erosus* seeds. *Plant Mol Biol* 57:13–20
- Spelbrink RG, Dilmac N et al (2004) Differential antifungal and calcium channel-blocking activity among structurally related plant defensins. *Plant Physiol* 135:2055–2067
- Towbin H, Staehelin T, Gordon J (1979) Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets—procedure and some applications. *PNAS* 76:4350–4354
- Vassileva A, Chugh DA (2001) Effect of copy number on the expression levels of hepatitis B surface antigen in the methylotrophic yeast *Pichia pastoris*. *Protein Expr Purif* 21:71–80
- Vijayan S, Guruprasad L, Kirti PB (2008) Prokaryotic expression of a constitutively expressed *Tephrosia villosa* defensin and its potent antifungal activity. *Appl Microbiol Biotechnol* 80:1023–1032
- Wang YP, Fristensky B (2001) Transgenic canola lines expressing pea defense gene DRR206 have resistance to aggressive blackleg isolates and to *Rhizoctonia solani*. *Mol Breed* 8:263–271
- Wang Y, Nowak G (1999) Constitutive expression of pea defense gene DRR206 confers resistance to blackleg (*Leptosphaeria maculans*) disease in transgenic canola (*Brassica napus*). *Mol Plant Microbe Interact* 12:410–418
- Wijaya R, Neumann GM, Condron R, Hughes AB, Polya GM (2000) Defense proteins from seed of *Cassia fistula* include a lipid transfer protein homologue and a protease inhibitor plant defensin. *Plant Sci* 159:243–255
- Wong JH, Ng TB (2005) Sesquin, a potent defensin-like antimicrobial peptide from ground beans with inhibitory activities toward tumor cells and HIV-1 reverse transcriptase. *Peptides* 26:1120–1126
- Ye XY, Ng TB (2001) Peptides from pinto bean and red bean with sequence homology to cowpea 10-kDa protein precursor exhibit antifungal, mitogenic, and HIV-1 reverse transcriptase-inhibitory activities. *Biochem Biophys Res Commun* 285:424–429
- Zhu YJ, Agbayani R, Moore PH (2007) Ectopic expression of *Dahlia merckii* defensin DmAMP1 improves papaya resistance to *Phytophthora palmivora* by reducing pathogen vigor. *Planta* 226:87–97