

RESEARCH ARTICLE

Quantitative proteomic analysis of S-nitrosated proteins in diabetic mouse liver with ICAT switch method

Xu Zhang^{1,2*}, Bo Huang^{1,2*}, Xixi Zhou^{1,2}, Chang Chen¹✉

¹ National Laboratory of Biomacromolecules, Institute of Biophysics, Chinese Academy of Sciences, Beijing 100101, China

² Graduate School of the Chinese Academy of Sciences, Beijing 100049, China

✉ Correspondence: changchen@moon.ibp.ac.cn

Received June 9, 2010 Accepted June 22, 2010

ABSTRACT

In this study we developed a quantitative proteomic method named ICAT switch by introducing isotope-coded affinity tag (ICAT) reagents into the biotin-switch method, and used it to investigate S-nitrosation in the liver of normal control C57BL/6J mice and type 2 diabetic KK-Ay mice. We got fifty-eight S-nitrosated peptides with quantitative information in our research, among which thirty-seven had changed S-nitrosation levels in diabetic mouse liver. The S-nitrosated peptides belonged to forty-eight proteins (twenty-eight were new S-nitrosated proteins), some of which were new targets of S-nitrosation and known to be related with diabetes. S-nitrosation patterns were different between diabetic and normal mice. Gene ontology enrichment results suggested that S-nitrosated proteins are more abundant in amino acid metabolic processes. The network constructed for S-nitrosated proteins by text-mining technology provided clues about the relationship between S-nitrosation and type 2 diabetes. Our work provides a new approach for quantifying S-nitrosated proteins and suggests that the integrative functions of S-nitrosation may take part in pathophysiological processes of type 2 diabetes.

KEYWORDS ICAT switch, mass spectrometry, quantitative, S-nitrosation, type 2 diabetes

INTRODUCTION

It is well established that nitric oxide (NO) can regulate a diverse array of signal transduction pathways which play

important roles in both physiologic and pathological processes, partly acting through modification of cysteine residues of proteins (S-nitrosation) (Hess et al., 2005). Methods such as biotin-switch and SNOSID have been developed for studying S-nitrosation (Jaffrey and Snyder, 2001; Hao et al., 2006). However, these approaches will not be powerful enough for those researches concerning quantitative analysis of S-nitrosation. As known, the biotin switch method includes three key steps: blocking free cysteines, specific reduction of nitrosated cysteines (S-nitrosothiol, SNO) by ascorbate, and biotinylation of released free cysteines by biotinylation reagents such as biotin-HPDP. Then the biotinylated proteins or peptides can be purified by streptavidin-agarose for further analysis. Isotope-coded affinity tag (ICAT) reagents, reactive to free thiols, have been used extensively in quantitative proteomics (Gygi et al., 1999). Here, we replaced the biotin-HPDP in the original biotin-switch method with ICAT reagents following with mass spectrometry detection, allowing quantification of S-nitrosated proteins in samples at the proteomic level. This method was named ICAT switch.

Type 2 diabetes mellitus (T2DM) is now a major threat to human health (Zimmet, 2000). Although the pathology of the development of T2DM is complicated, chronic inflammation induced by obesity is thought to play a key role (Wellen and Hotamisligil, 2005). The expression of inducible NO synthase (iNOS) increases under inflammation conditions (Xie et al., 1994), and targeted disruption of iNOS protects against obesity-linked insulin resistance in muscle tissue (Perreault and Marette, 2001). Endogenous NO produced abnormally by iNOS has a variety of effects on some processes in the development of diabetes (Perreault and Marette, 2001; Sugita

*These authors contributed equally to the work.

et al., 2002; Fujimoto et al., 2005) and its complications (Ceriello et al., 2002), and alterations in the amount of S-nitrosated proteins such as insulin receptors, insulin receptor substrate 1, and Akt (Carvalho-Filho et al., 2005; Yasukawa et al., 2005; Wadham et al., 2007) are reported to be involved in these diseases. Liver is an important organ for many metabolic processes, such as gluconeogenesis, glycogenesis, fatty acid and triglyceride metabolism, many of which are abnormal in diabetes. Research on S-nitrosated proteins in liver at the proteomic level will help to uncover the role of S-nitrosation in diabetes.

In the present study we developed the ICAT switch method to identify and quantify S-nitrosated proteins in the liver of control C57BL/6J mice and diabetic model KK-Ay mice. The relationship between S-nitrosated proteins and diabetes was subsequently analyzed using bioinformatics.

RESULTS

Construction and validation of ICAT switch method

We labeled the S-nitrosated thiols in C57BL/6J samples with

light ICAT reagents and the S-nitrosated thiols in KK-Ay samples with heavy ICAT reagents. The peptides containing both quantitative and SNO site information were then identified by mass spectrometry. The flow chart of ICAT switch is shown schematically in Fig. 1A (ICAT structure was shown in Supplemental Fig. 1), in which ICAT reagents were used to replace biotin-HPDP to label the S-nitrosated Cys residues. Taking one result as an example, the peaks of peptide AGCQVVPSDMMDGR labeled with ICAT reagents can be seen in precursor MS spectrum as shown in Fig. 1B. The difference in m/z units between light and heavy ICAT labeled peptides is 9 Da for univalent ions, 4.5 Da for divalent ions, and 3 Da for trivalent ions. This spectrum indicated that ICAT reagent could be used to identify S-nitrosation with mass spectrometry.

To test the accuracy of ICAT switch, we applied this method in quantifying S-nitrosation induced by S-nitrosoglutathione (GSNO) *in vitro*, whose ratio could be controlled. The liver lysates of normal mice were treated with 500 μ M GSNO for protein S-nitrosation followed by ICAT switch. Half was biotinylated by L (light) and another half by H (heavy) ICAT reagent. Since the levels of S-nitrosation in both halves of the




Figure 1. Construction and validation of the ICAT switch method. (A) Schematic overview of ICAT switch. Proteins were extracted from mouse liver and added to a blocking buffer, in which the free Cys-thiols were methylthiolated by MMTS. S-NO bonds in proteins from C57/BL6 and KK-Ay mouse liver were then separately reduced by ascorbate and labeled with ICAT light/heavy reagents. Samples from C57/BL6 and KK-Ay mouse liver were then combined and digested by trypsin at 37°C overnight. Peptides labeled with ICAT reagents were purified by monomeric avidin agarose, and then acid-cleaved by cleaving reagents. Finally, peptides were analyzed by LC-MS/MS. (B) Representative mass spectrum of the ICAT labeled peptides in full scan with three kinds of mass intervals. The difference of mass values between the isotopic peptides was 9 Da for univalent ions, 4.5 Da for divalent ions, and 3 Da for trivalent ions. (C) Box plot of light/heavy ratios of S-nitrosated peptides is shown. The middle line in the box indicates the median of light/heavy ratios in this group of data. The box indicates 50% members of the group.

lysates were the same, the expected value of the ratios was 1.0 (L/H = 1:1). 127 S-nitrosated peptides were identified with an average ratio 0.94 and the distribution was shown in Fig. 1C. These results guaranteed the accuracy of our method used afterward.

S-nitrosated peptides and proteins identified in C57BL/6J and KK-Ay diabetic mouse liver

We analyzed the different S-nitrosation patterns between normal and diabetic mouse liver by ICAT switch. The probability above 0.8 for every peptide acquired by Peptideprophet was set as the criteria for peptide acceptance. Filtration of S-nitrosated peptide was shown in Fig. 2A. Employing a threshold of Probability above 0.8, the percent of false S-nitrosated peptides identified in this experiment is below 3.6%. In this state, the fraction of all correct assignments passing the minimal probability thresholds (MPT) filter is nearly 75% (sensitivity). The accepted peptides contained Cys were quantified afterward by ASAPRatio, which was part of the TPP software package. The peptide AGCQVPSDMMMDGR in aminolevulinic acid delta dehydratase (gi 34328485) was taken as an example, in which heavy ICAT reagents were labeled on the cysteine (Fig. 2B). There was a shift of 236 (the mass of the heavy ICAT tag) in the m/z from the b_3 to the b_{13} ion and an ICAT diagnostic peak at m/z 294.9 (c1). Fig. 2C shows quantification of the peptide AGCQVPSDMMMDGR by ASAPRatio. The ratio of signals from light (C57BL/6J) versus heavy (KK-Ay) ICAT labeled peptides was determined to be 0.46 ± 0.03 , meaning that the amount of S-nitrosated peptides in KK-Ay increased to about 2.2 fold ($1/0.46$) that of C57BL/6J. S-nitrosated peptides whose amount is changed in diabetic mouse liver ($L/H < 0.7$ or $L/H > 1.3$) were listed in Table 1, and the others ($0.7 \leq L/H \leq 1.3$) were shown in Supplemental Table 1. After checking the MS/MS spectra and the SIC (single ion chromatogram) manually by ASAPRatio, 58 peptides were considered as S-nitrosated peptides containing quantitative information. The distribution of the peptides ratios is shown in Fig. 3A. Of the 58 peptides identified, 24 peptides whose L/H ratio was less than 0.7 were considered as S-nitrosation increased in KK-Ay mouse liver, 13 peptides whose L/H ratio was more than 1.3 were considered as S-nitrosation decreased, and 21 peptides whose L/H ratio was between 0.7 and 1.3 were considered to show no change in S-nitrosation status, because changes below $\pm 30\%$ were considered as not significantly different in accordance with the ICAT kit instructions. An L/H ratio of 0 meant that the S-nitrosated peptide was only present in KK-Ay mouse liver, and an L/H ratio of 999 meant that the S-nitrosated peptide was only present in C57BL/6J mouse liver.

Bioinformatic analysis of S-nitrosated proteins

To identify which biological processes were more likely to be

regulated by quantitative changes in S-nitrosated proteins, we performed GO enrichment of these proteins using the BiNGO plugin in the Cytoscape platform. Thirty proteins with altered levels of S-nitrosation in KK-Ay mouse liver were analyzed by GO enrichment tool BiNGO. The enriched GO terms were visualized using a DAG (directed acyclic graph) graphical representation (Fig. 3B). The "amino acid metabolic processes" had a low *p*-value, which meant this process had higher probability to be regulated by S-nitrosation proteins. Other biologic processes enriched are listed in Supplemental Table 2.

Since S-nitrosated proteins may affect signaling pathways indirectly through their impact on other proteins (Benhar and Stamler, 2005; Qu et al., 2007), we constructed a protein network containing 14 S-nitrosated proteins identified in KK-Ay mouse liver and other related proteins by virtue of text-mining technology to find out the latent functions of the S-nitrosated proteins (Fig. 4A). Green, white and red nodes stand for S-nitrosated proteins, and the gradient from green to red reflects the S-nitrosation change from increasing to decreasing. Grey nodes stand for related proteins to S-nitrosated targets. The edge between two nodes stands for the relationship between them. For example, Fig. 4B shows a subnet composed of an S-nitrosated protein and its related proteins. In this subnet, the green color of the node for serine/threonine kinase receptor-associated protein (STRAP), which was identified as an S-nitrosated protein in our research, means that the amount of S-nitrosated STRAP increased in KK-Ay mouse liver. Other gray nodes stand for proteins related with STRAP. Of the related proteins, the edge between Cdkn1a (an alias of p21) and STRAP shows that stable expression of STRAP induces downregulation of the cyclin-dependent kinase inhibitor p21 (Halder et al., 2006); the edge between trp53 (an alias of p53) and STRAP shows that STRAP activates p53 (Jung et al., 2007).

DISCUSSION

Here we developed the ICAT switch method for quantitative proteomic research on S-nitrosation in diabetic mouse liver. By virtue of the ICAT switch in quantification, we solved the problem of false negative identification of S-nitrosation targets. In our study there are 20 peptides with $0 < L/H < 0.7$ and seven peptides with $1.3 < L/H < 999$, which were identified as new S-nitrosation targets and their quantitative variation was determined. However, using former methods lacking quantitative information, these proteins would have been eliminated from the S-nitrosated protein target list since they were presented in both control C57/BL6 and the KK-Ay mouse liver, therefore, leading to false negative identification of S-nitrosation targets. Some S-nitrosated targets reported in other studies, such as transaldolase (Lefièvre et al., 2007), glutathione S-transferase (Sies et al., 1998; Lindermayr et al., 2005; Hao et al., 2006; Lefièvre et al., 2007), D-dopachrome tautomerase (Han and Chen, 2008),




Figure 2. Filtering, identification and quantification of S-nitrosated peptides. (A) Filtration of S-nitrosated peptide. The MPT for false positive (error) rate below 0.05 is 0.68. Employing a threshold of Probability (PeptideProphet) above 0.8, we make sure the percent of false S-nitrosated peptides identified in this experiment is below 3.6%. In this state, the fraction of all correct assignments passing MPT filter is nearly 75% (sensitivity). (B) Identification of the peptide AGCQVVPSDMMMDGR in aminolevulinatase delta dehydratase. Red dots indicate matching of b-ions, and blue dots indicate y-ions. Orange dots labeled with b* indicate dehydrated b-ions. The purple dot at m/z 294.9 shows the ICAT diagnostic peak. (C) Reconstructed ion chromatogram of the precursor ion from peptide AGCQVVPSDMMMDGR in its light and heavy versions using ASAPRatio. The ratio of light (C57/BL6) versus heavy (KK-Ay) ICAT labeled peptides was determined to be 0.46 ± 0.03 .

Table 1 List of S-nitrosated proteins, S-nitrosated sites and quantitative information of S-nitrosated peptides, whose amount is changed in diabetic mouse liver (L/H < 0.7 or L/H > 1.3), as revealed by ICAT switch.

sequence	probability (peptide-prophet)	ASAP ratio ± SD (L/H)	gi accession ^a	Entrez Gene ID ^b	gene symbol ^c	Z	MH +	ΔM	protein description	probability (protein-prophet)
R.ADPHLQDFLES HYLDK.E ^d	1.0000	0.00 ± 0.00	82982537	none	LOC634386	3	2141.228	1.00871	ferritin light chain 1	0.9975
K.LACADITINSIY GLFVLVSTFGMD LLFIFLSYVILR.S	0.9926	0.00 ± 0.00	22128815	none	OLFR641	3	4401.91	1.75489	olfactory receptor 641	0.979
-MNTDGGGAR K.R	0.9098	0.00 ± 0.00	82917032	none	SORBS2	2	1393.298	0.89466	Arg/Abi-interacting protein ArgBP2,isoforms	0.8691
K.MVATGVCR.S	1.0000	0.00 ± 0.00	6724311	11522	ADH1	2	1072.496	0.678	alcohol dehydrogenase 1 (class I)	0.9982
K.ALAGCDFLTIS PK.L	0.9883	0.13 ± 0.02	33859640	21351	TALDO1	2	1572.706	0.0032	transaldolase 1	0.9771
K.AAGCDFN ^W K.T	0.9036	0.18 ± 0.00	40807498	15473	HRSP12	2	1365.115	0.29113	heat-responsive protein 12	1.0000
K.VHITLS ^W THECA GLSER.D	1.0000	0.26 ± 0.03	13384608	13180	PCBD1	3	1990.277	0.18839	pterin 4 alpha carbinola- mine dehydratase/dimeri- zation cofactor of hepatocyte nuclear factor 1 alpha (TCF1) 1	0.9975
K.TEIQVNC ^W PK.V	0.8409	0.35 ± 0.03	41282044	19027	SYPL	2	1260.224	0.90051	pantophysin 2	1.0000
K.NPADGAC ^W LE K.Y	1.0000	0.35 ± 0.03	19526926	70984	4931406C07 RIK	2	1359.174	0.76213	pantophysin 1 hypothetical protein LOC70984	1.0000 0.9985
R.GLGCSISSGPIQ K.P	1.0000	0.37 ± 0.06	12963769	72088	USH1C	3	1481.43	2.14405	harmonin isoform a1	0.9990
K.CLDAFP ^W NLR.D ^d	0.8582	0.38 ± 0.01	33468899	14864	GSTM	2	1275.969	0.38671	glutathione S-transferase, isoforms	0.9939
R.AYPIDQYPCR.T	1.0000	0.44 ± 0.03	21450101	243537	UROC1	2	1454.282	0.76857	urocanase domain con- taining 1	1.0000
K.AGCQV ^W VAPSD MMDGR.V	1.0000	0.46 ± 0.03	34328485	17025	ALAD	2	1773.221	0.67436	aminolevulinatase, delta-, dehydratase	0.9989
K.FPDVPGFS ^W WT PCISAK.D	0.9517	0.47 ± 0.01	7106255	11846	ARG1	2	2078.209	1.06322	arginase 1, liver	1.0000

(Continued)

sequence	probability (peptide-prophet)	ASAP ratio ± SD (L/H)	gi accession ^a	Entrez Gene ID ^b	gene symbol ^c	Z	MH +	ΔM	protein description	probability (protein-prophet)
K.EGLALINGTQM ITSLGQEALER.A	0.9996	0.54 ± 0.03	6754152	638196	HAL	3	2272.681	0.81572	histidine ammonia lyase	1.0000
K.LVQSGPLTTCR.I	0.8747	0.61 ± 0.08	6754098	110006	GUSB	2	1402.201	0.31011	glucuronidase, beta	1.0000
K.GHFGPIHCVR.F	1.0000	0.61 ± 0.05	6755682	20901	STRAP	3	1351.122	0.67925	serine/threonine kinase receptor associated pro- tein	0.9989
R.NAGGTYQNLID MVSSAIGCQLG K.D	1.0000	2.53 ± 0.27	31981882	223978	C530044N13 RIK	3	2565.488	1.73676	hypothetical protein LOC223978	0.9975
R.LEECKDAGLV K.S	0.9128	3.06 ± 0.21	7304879	27384	AKR1C13	2	1441.221	0.31294	aldo-keto reductase family 1, member C13	0.9166
R.NLRPGDLCDR.D	1.0000	3.15 ± 0.33	94395919	none	no match	3	1386.38	0.04859	aldo-keto reductase family 1, member C12	0.9166
K.CPVPSNMTRIR.F	0.8791	31.76 ± 3.80	12738842	64817	SVEP1	2	1518.314	0.64523	hypothetical protein LOC75300	0.9989
R.HYAHTDCPGH ADYVKN	1.0000	999 ± 0.00	94408128	none	no match	3	1726.699	0.29563	Sushi, von Willebrand factor type A, EGF and pentraxin domain con- taining 1	0.9841
K.PTGQRFQINSV ALKFKPSK.P	0.8143	999 ± 0.00	21313208	76467	MSRB2	3	2348.931	0.71841	tripartite motif protein TRIM5	0.9990
K.YECLPGYGRGI SR.M	1.0000	999 ± 0.00	31542275	12269	C4BP	3	1706.295	1.4983	pilin-like transcription fac- tor	0.8152
K.HSCPDVSTIVQ EKRRPK.R	1.0000	999 ± 0.00	9506867	54562	LRRC6	3	2352.299	2.23723	complement component 4 binding protein	0.9985
R.CVLGN SAVS VEQWKA	1.0000	999 ± 0.00	19526800	13872	ERCC3	2	1863.128	2.10713	leucine rich repeat con- taining 6	0.9985
R.CASGAE LRQPP R.D	1.0000	999 ± 0.00	51829389	none	6430704N06	3	1522.385	0.79939	excision repair cross- complementing rodent repair deficiency, comple- mentation group 3	0.9989
R.CVILPDSVR.N	0.9654	0.13 ± 0.00	21450071	12411	CBS	2	1328.297	0.17505	5T4 oncofetal trophoblast glycoprotein cystathionine beta- synthase, isoform 1	0.9997

(Continued)

sequence	probability (peptide-prophet)	ASAP ratio ± SD (L/H)	gi accession ^a Gene ID ^b	Entrez Gene ID ^b	gene symbol ^c Z	MH + ΔM	protein description	probability (protein-prophet)
R_CIVMPEK.M	0.9357	0.24 ± 0.01	30089694 12411	12411	CBS 2	1328.297 0.17505	cystathionine beta-synthase, isoform 2	0.9997
R_NASNPLAHYD DTAEELQQCDG K.L	1.0000	0.64 ± 0.06	21450071 30089694 12411	12411	CBS 2	1160.42 0.07845	cystathionine beta-synthase, isoform 1	0.9997
R_NASNPLAHYD DTAEELQQCDG K.L	1.0000	0.64 ± 0.06	21450071 30089694 12411	12411	CBS 3	2761.603 0.80564	cystathionine beta-synthase, isoform 2	0.9997
R_LCAATATILDK PEDR.V	0.9998	0.42 ± 0.03	30089694 6753618 13202	12411	CBS 2	2761.603 0.80564	cystathionine beta-synthase, isoform 2	0.9997
K_STEPCAHLLVS SIGVVGTAEQNR.T	1.0000	0.60 ± 0.05	6753618 6753618 13202	13202	DDT 3	1844.962 2.16118	D-dopachrome tautomerase	1.0000
R_CMLDRDEDML ITGGR.H	0.9681	0.69 ± 0.10	77682555 22436	22436	DDT 2	1961.772 0.34849	xanthine dehydrogenase	1.0000
R_AVQNACQILM K.R	0.9866	1.93 ± 0.50	13994153 71724	71724	AOX2 2	1446.235 0.39549	aldehyde oxidase structural homolog 2	1.0000
R_MACEDQFTNL VPQTDSK.C	1.0000	2.61 ± 0.34	13994153 71724	71724	AOX2 2	2155.56 0.29901	aldehyde oxidase structural homolog 2	1.0000
R_NHPEPSTEQIM ETLGGNLCR.C	0.9999	3.06 ± 0.20	13994153 71724	71724	AOX2 2	2463.614 0.02224	aldehyde oxidase structural homolog 2	1.0000

S-nitrosated peptides are shown in the sequence column, and the cysteine residues of the peptides are the S-nitrosated sites. The ASAP Ratio ± SD (L/H) stands for the ratio of S-nitrosated peptides (C57/BL6: KK-Ay). S-nitrosated peptides whose amount is not changed significantly ($0.7 \leq L/H \leq 1.3$) are shown in Supplemental Table 1.

^a The id from the NCBI database is used for identification.

^b Entrez Gene IDs were transformed from gi accessions and used for GO enrichment.

^c Gene Symbols were transformed from gi accessions, and used for text mining.

^d A group of proteins containing more than 3 isoforms. Only a member was listed.




Figure 3. Analysis of S-nitrosated peptides and proteins identified in the normal and diabetic mouse liver using ICAT switch. (A) The number of S-nitrosated peptides distributed in different L/H (C57/BL6 versus KK-Ay) ratio groups. L/H = 0: S-nitrosated peptide only presents in KK-Ay mouse liver; $0 < L/H < 0.7$: S-nitrosation increased in KK-Ay mouse liver; $0.7 \leq L/H \leq 1.3$: no change in S-nitrosated peptides from KK-Ay mouse liver; $1.3 < L/H < 999$: S-nitrosation decreased in KK-Ay mouse liver; L/H = 999: S-nitrosated peptides only present in C57BL/6J mouse liver. (B) GO enrichment result of proteins with changed S-nitrosation level. Numbers in the nodes represented IDs of GO terms. The color from yellow to red (dependent on p -value) reflected their degree of enrichment.

aspartate aminotransferase (Lefièvre et al., 2007), and glutathione peroxidase 1 (Koh et al., 2001; Lindermayr et al., 2005), were also detected in our study, indicating the efficiency of the ICAT switch.

With this quantitative proteomic method, we were able to find a number of new S-nitrosated targets that are known to be involved in diabetes and thus merit special investigation. (A) Arg/Abl binding protein (ArgBP2), this multi-adaptor protein interacts with many important proteins in cell signaling (Wang et al., 1997), including Akt (Yuan et al., 2005) and cbl (Soubeyran et al., 2003), which are involved in the insulin-induced glucose transport pathway. S-nitrosated ArgBP2 was only detected in KK-Ay mouse liver, suggesting that S-nitrosation of this protein may contribute to dysfunction of the glucose transport pathway. (B) Hepatocyte pterin 4 alpha carbinolamine dehydratase/dimerization cofactor (PCBD1), also known as dimerizing cofactor for HNF1 (DCoH), can bind to the HNF1 family of transcriptional activators, nuclear factor 1 alpha (TCF1) 1 (Mendel et al., 1991). The HNF1-DCoH complex regulates the expression of a large group of genes related to metabolism and the insulin pathway in liver. Defects

in the expression of these genes underlie maturity-onset diabetes of the young (MODY) (Yamagata et al., 1996a, b), and mice lacking DCoH display hyperphenylalaninemia and an impaired glucose tolerance (Bayle et al., 2002). The His80 and Glu81 residues (Rose et al., 2004) are thought to be the active site of DCoH1, and the 3.9-fold increase in S-nitrosation of Cys82 in diabetic mouse liver detected in our experiment may affect the structure near the active site and cause dysfunction. (C) Pantophysin (SYPL) (Leube, 1994), it is reported to associate with GLUT4-containing vesicles and may regulate vesicle transport in adipocytes (Brooks et al., 2000). Here we found that S-nitrosation of this protein increased 2.9 folds in diabetic mouse liver. Whether this S-nitrosation affects Pantophysin's role in GLUT translocation is expected for further research. (D) Aminolevulinic acid delta dehydratase (ALAD), its activity decreased in the liver of high glucose-induced diabetic rats (Folmer et al., 2002) and the oxidation of cysteines is thought to be one possible reason (Afonso et al., 1996). Here we found that S-nitrosation of ALAD increased 2.17 folds in diabetic mouse liver, suggesting one possible contribution to the decrease of




Figure 4. Integrative analysis of the network of S-nitrosated proteins. (A) Text-mining network constructed by S-nitrosated proteins. The nodes colored green, white or red stand for S-nitrosated proteins, and the gradient from green to red reflects the increasing ratio of light (C57/BL6) versus heavy (KK-Ay) ICAT labeled peptides. Grey nodes are proteins that affect or are affected by S-nitrosated proteins. (B) Topological connections of the S-nitrosated protein STRAP. Grey nodes stand for proteins related to STRAP by text-mining. The edge between two nodes stands for the relationship. Cdkn1a and trp53 are known to be regulated by STRAP from the information provided by this subnet, and these two proteins are associated with diabetes.

ALAD activity. (E) Aldo-keto reductase family 1, member C12/C13 (AKR1C12/C13) (S-nitrosation decreased by 67%), this superfamily is related to the pathogenesis of secondary complications associated with diabetes (Bhatnagar and

Srivastava, 1992). Aldose reductase (AR) is another member of the aldo-keto reductase superfamily, and the inhibition of AR is thought to prevent diabetic complications. AR is very sensitive to nitric oxide donors and increasing NO

availability can usefully inhibit the polyol pathway via down-regulating AR activity and alleviating the development of diabetes complications (Dixit et al., 2001; Chandra et al., 2002). The functional role of the decrease in the extent of S-nitrosation of AKR1C12/C13 might be one possible factor for the development of diabetes. (F) Alcohol dehydrogenase 1 (class I) (gi6724311), which was S-nitrosated only in KK-Ay mouse liver, could depress the phosphorylation of insulin receptor (Tyr1146) and reduced phosphorylation of Akt and GSK-3 β in cardiomyocytes from ethanol-fed mice (Li and Ren, 2008). These are similar to the abnormal insulin signaling pathway in liver or peripheral diabetic tissues (Cohen, 2006).

In addition, with the help of GO enrichment, the integrative analysis results showed that the S-nitrosated proteins in KK-Ay mouse liver were enriched in processes related with amino acid metabolism (Fig. 3B). For example, one S-nitrosated target is arginase 1 (Entrez Gene ID: 11846), which is responsible for hydrolyzing L-arginine to form L-ornithine and urea, is involved in the nitrogen metabolism pathway and urea cycle. There are also reports that amino acid metabolism was abnormal in diabetes (Wijekoon et al., 2004; Noguchi et al., 2006). Excess amino acids in the liver can act as substrates for gluconeogenesis (Felig, 1975), interfere with the inhibitory effect of insulin on glucose production (Boden and Tappy, 1990), and change the insulin/glucagon ratio (Roden et al., 1996), all of which may increase the output of hepatic glucose. Moreover, abnormal amino acid metabolism in the liver may change levels of amino acids throughout the whole body. Some reports suggest that high levels of amino acids can have both positive (Tremblay and Marette, 2001; Nobukuni et al., 2005) and negative (Armstrong et al., 2001; Bogan et al., 2001) effects on glucose metabolism and insulin action in peripheral tissues. Amino acid levels can also affect insulin secretion in pancreatic islet β -cells (Newsholme et al., 2007). Therefore, it is possible that S-nitrosation interferes with amino acid metabolic processes and thus contributes to diabetes.

Meanwhile, we noticed that not every S-nitrosated protein identified in our research can be linked with diabetes or its complications directly. Thus, we used text-mining network to help to reveal the relationships. As shown in Fig. 4B, this subnet composed by STRAP and nodes connected to it, such as p21 and p53. p21 is known to promote adipose tissue expansion and leads to insulin resistance (Inoue et al., 2008), and many proteins affecting the insulin pathway are regulated by p53 (Feng et al., 2007). This subnet indicates that S-nitrosation of STRAP may play an indirect role in diabetes via p21 and p53. Therefore, the network constructed for S-nitrosated proteins provided clues about the relationship between S-nitrosation and type 2 diabetes.

With the new developed ICAT switch method, we got the quantitative proteomic profile of S-nitrosation in C57/BL6 and KK-Ay mouse liver. The quantitative information, together with

the network analysis of S-nitrosated targets contributed to explaining the relationship between S-nitrosation and diabetes. Our work provides a solution for quantification of multiple S-nitrosation, makes it possible to elucidate the integrative correlation of S-nitrosation with pathophysiologic processes. The mode of this work can be used to study S-nitrosation profiles in other physiologic and pathological processes.

MATERIALS AND METHODS

Reagents

Cleavable ICATTM reagent kits were purchased from Applied Biosystems (Foster City, CA, USA), Methyl methanethiosulfonate (MMTS), biotin-(N-(6-(Biotinamido)hexyl)-3'-(2'-pyridyl)thio)-propionamide (biotin-HPDP), and BCATM protein assay kits were from Pierce (Rockford, IL, USA). PlusOneTM urea was from GE Healthcare (Piscataway, NJ, USA). Protease inhibitor cocktail tablets (Complete, Mini, EDTA-free) were from Roche Applied Sciences (Indianapolis, IN, USA). Sequencing-grade modified trypsin (V5113) was from Promega (Madison, WI, USA). Solvents used in LC/MS analysis, including formic acid (FA), acetonitrile (ACN) and methanol, were from J.T. Baker Chemicals (Philipsburg, NJ, USA). All other chemicals were from Sigma-Aldrich (St. Louis, MO, USA).

Animals

Eight-week-old female lean wild-type C57BL/6J and KK-Ay mice were obtained from the Institute of Experimental Animals, Beijing (Beijing, China). The mice were housed in mesh cages in a room maintained at 25°C and illuminated in 12:12-h light/dark cycles. They were provided with a standard rodent diet and water. All animal experimental procedures were conducted according to the Animal Welfare Act 2006.

Protein Preparation

Livers from C57BL/6J and KK-Ay mice were homogenized separately in detergent-free lysis buffer (50 mM Tris, pH 7.7, 2.5 mM EDTA, 0.1 mM neocuproine, 8 M Urea, 20 mM MMTS and protease inhibitor cocktail) on ice, as described in the literature (Han and Chen, 2008). Lysates were centrifuged at 12,000 g for 20 min at 4°C to remove the debris. All steps were performed in the dark. Protein concentration was determined using a BCATM protein assay kit and adjusted to 1 mg/mL, after removal of MMTS from the protein solution by TCA/acetone precipitation.

In each individual experiment, one control mouse and one diabetic KK-ay mouse were used. Independent experiments were conducted for 9 times and representative data were presented.

ICAT switch method for isolation of S-nitrosated peptides

Protein samples were dissolved in blocking buffer (20 mM Tris, pH 7.7, 1 mM EDTA, 0.1 mM neocuproine, 8 M urea, 20 mM MMTS) and incubated for 1 h at room temperature in dark (Han and Chen, 2008). After TCA/acetone precipitation, the pellets were dissolved in

labeling buffer (20 mM Tris, pH 7.7, 1 mM EDTA, 0.1 mM neocuproine, 4 M urea and a vial of ICAT reagents) in the presence of sodium ascorbate. Protein obtained from C57BL/6J and KK-Ay mice was labeled with isotopically light and heavy ICAT reagents, respectively, and incubated for 2 h at room temperature. Labeled preparations were combined and precipitated by TCA/acetone. Pellets were dissolved in enzymolysis buffer (50 mM NH_4HCO_3 , pH 8.3, 1.5 M urea) and digested with trypsin overnight at 37°C using an enzyme-to-protein ratio of 1:50. The resulting peptide mixture was cleaned using a cation-exchange cartridge. Cleaned peptides were collected and subjected to affinity purification using monomeric avidin agarose. Peptides labeled with ICAT reagents were enriched and then cleaved with cleaving reagent at 37°C for 2 h. The cation-exchange cartridge, cleaving reagent and relevant buffers were from a cleavable ICATTM reagents kit, and procedures for purifying peptides were according to the manufacturer's guidelines. ICAT labeled peptides were dried by vacuum centrifuge and resolubilized in 0.1% formic acid for LC-MS/MS analysis.

LC/MS/MS analysis

Peptides from which the biotin tag had been removed were separated by reverse-phase capillary liquid chromatography (RP-C18, Column Technology Inc, 0.15 mm × 150 mm) at a flow rate of 200 nL/min. The eluent was directly analyzed by a Thermo LTQ linear trap instrument equipped with a Thermo micro-electrospray source, a Thermo Surveyor pump and an autosampler (Thermo Finnigan, San Jose, CA, USA). A survey scan followed by 5 CID (collision induced dissociation) events was used. Peptide identification by CID was carried out in automated mode using the 3-min dynamic exclusion option.

Database search, data validation and quantification

MS/MS spectra were searched against a database downloaded from the National Center for Biotechnology Information website (NCBI; mouse; May 4, 2007; 46,903 entries) using SEQUEST (Yates et al., 1995) in Bioworks 3.2 (Thermo Finnigan). The SEQUEST database search criteria included a static modification of cysteine residues of 227 Da (light cleavable ICAT reagent) and a variable modification of 9 Da for cysteines (for the heavy cleavable ICAT reagent). The tolerance of the precursor ion and fragment ion were 2 and 1 amu, respectively. Peptides were permitted to have up to two missed cleavages. The .dta and .out files obtained from SEQUEST were validated using PeptideProphet (Keller et al., 2002) and ProteinProphet (Nesvizhskii et al., 2003). The minimal probability thresholds (MPT) were set to make false-positive error rates for both peptides and proteins were below 0.05. The peptides passing the MPT statistic filter in corresponding experiment were accepted and quantified. The minimal intensity of precursor ions peak was set to 1000 when acquiring MS/MS data. Relative quantification with SIC (single ion chromatogram) was performed using ASAPRatio (Li et al., 2003). ASAPRatio grouped abundance ratios of same sequence peptide (including different isotopic forms, charge states, repeats) in one RP elution peak together to calculate the mean and standard deviation first, then abundance ratios of same sequence peptide in different RP elution peaks were grouped together to get the final mean and standard deviation for a unique peptide. Software packages were utilized via the Trans-Proteomics Pipeline (ISB, Seattle, WA, USA).

The resulting peptide spectra of proteins were manually checked for qualitative and quantitative results.

Bioinformatic analysis of the results of S-nitrosation

The database ID (gi Accession) of the S-nitrosated proteins was converted into an Entrez Gene ID using the DAVID Gene ID Conversion Tool (Dennis et al., 2003). Entrez Gene IDs were used to do GO enrichment. S-nitrosated proteins were visualized with Cytoscape (Shannon et al., 2003) and used to do GO enrichment using the BiNGO (Maere et al., 2005) plugin with the Gene Ontology (GO) annotation (Ashburner et al., 2000). The Benjamini and Hochberg False Discovery Rate (FDR) correction was chosen as the testing correction, and a *p*-value threshold of 0.05 was selected for Biological Process categories in the BiNGO settings. A text-mining network was constructed for S-nitrosated proteins and their related proteins via the Agilent Literature Search (Vailaya et al., 2005) plugin of Cytoscape in order to analyze the contribution of the S-nitrosated proteins to diabetes through their impact on other proteins. In the Agilent Literature Search tool, Gene Symbols of S-nitrosated proteins were used to search the Pubmed database; "Mus musculus" was set as the concept lexicon; Max Engine Matches was set to 10; and aliases were used. The network was displayed by Cytoscape, and relationships between nodes were checked manually. Wrong nodes and edges were removed, and nodes standing for S-nitrosated proteins were colored using a green-to-red gradient according to the amount of S-nitrosated peptides.

ACKNOWLEDGEMENTS

We are grateful to Prof. Rong Zeng, Prof. Taijiao Jiang and Prof. John Yates III for valuable suggestions and discussions. This research was supported by the National Basic Research Program (973 program) of China (Grant Nos. 2006CB911001, 2005CB522804), the National Natural Science Foundation of China (Grant Nos. 90606020 and 30770512) and the Knowledge Innovation Program of the Chinese Academy of Sciences.

ABBREVIATIONS

biotin-HPDP, biotin-(N-(6-(biotinamido)hexyl)-3'-(2'-pyridyldithio)-propionamide; CID, collision induced dissociation; DAG, directed acyclic graph; GLUT, glucose transporter; GO, gene ontology; GSNO, S-nitrosoglutathione; ICAT, isotope-coded affinity tag; iNOS, inducible NO synthase; MMTS, methyl methane thiosulfonate; MODY, maturity onset diabetes of the young; MPT, minimal probability threshold; NO, nitric oxide; RNS, Reactive nitrogen species; SNOSID, SNO-site identification; SNO, S-nitrosothiol; STRAP, serine/threonine kinase receptor-associated protein; T2DM, type 2 diabetes mellitus

REFERENCES

- Afonso, S.G., Polo, C.F., Enriquez de Salamanca, R., and Batlle, A. (1996). Mechanistic studies on uroporphyrin I-induced photoinactivation of some heme-enzymes. *Int J Biochem Cell Biol* 28, 415–420.
- Armstrong, J.L., Bonavaud, S.M., Toole, B.J., and Yeaman, S.J. (2001). Regulation of glycogen synthesis by amino acids in cultured human muscle cells. *J Biol Chem* 276, 952–956.

- Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight, S.S., Eppig, J.T., *et al.* (2000). Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet* 25, 25–29.
- Bayle, J.H., Randazzo, F., Johnen, G., Kaufman, S., Nagy, A., Rossant, J., and Crabtree, G.R. (2002). Hyperphenylalaninemia and impaired glucose tolerance in mice lacking the bifunctional DCoH gene. *J Biol Chem* 277, 28884–28891.
- Benhar, M., and Stamler, J.S. (2005). A central role for S-nitrosylation in apoptosis. *Nat Cell Biol* 7, 645–646.
- Bhatnagar, A., and Srivastava, S.K. (1992). Aldose reductase: congenial and injurious profiles of an enigmatic enzyme. *Biochem Med Metab Biol* 48, 91–121.
- Boden, G., and Tappy, L. (1990). Effects of amino acids on glucose disposal. *Diabetes* 39, 1079–1084.
- Bogan, J.S., McKee, A.E., and Lodish, H.F. (2001). Insulin-responsive compartments containing GLUT4 in 3T3-L1 and CHO cells: regulation by amino acid concentrations. *Mol Cell Biol* 21, 4785–4806.
- Brooks, C.C., Scherer, P.E., Cleveland, K., Whittemore, J.L., Lodish, H.F., and Cheatham, B. (2000). Pantophysin is a phosphoprotein component of adipocyte transport vesicles and associates with GLUT4-containing vesicles. *J Biol Chem* 275, 2029–2036.
- Carvalho-Filho, M.A., Ueno, M., Hirabara, S.M., Seabra, A.B., Carneiro, J.B., de Oliveira, M.G., Velloso, L.A., Curi, R., and Saad, M.J. (2005). S-nitrosation of the insulin receptor, insulin receptor substrate 1, and protein kinase B/Akt: a novel mechanism of insulin resistance. *Diabetes* 54, 959–967.
- Ceriello, A., Quagliaro, L., D'Amico, M., Di Filippo, C., Marfella, R., Nappo, F., Berrino, L., Rossi, F., and Giugliano, D. (2002). Acute hyperglycemia induces nitrotyrosine formation and apoptosis in perfused heart from rat. *Diabetes* 51, 1076–1082.
- Chandra, D., Jackson, E.B., Ramana, K.V., Kelley, R., Srivastava, S. K., and Bhatnagar, A. (2002). Nitric oxide prevents aldose reductase activation and sorbitol accumulation during diabetes. *Diabetes* 51, 3095–3101.
- Cohen, P. (2006). The twentieth century struggle to decipher insulin signalling. *Nat Rev Mol Cell Biol* 7, 867–873.
- Dennis, G. Jr, Sherman, B.T., Hosack, D.A., Yang, J., Gao, W., Lane, H.C., and Lempicki, R.A. (2003). DAVID: Database for Annotation, Visualization, and Integrated Discovery. *Genome Biol* 4, 3–13.
- Dixit, B. L., Ramana, K. V., Chandra, D., Jackson, E. B., Srivastava, S., Bhatnagar, A., and Srivastava, S. K. (2001). Metabolic regulation of aldose reductase activity by nitric oxide donors. *Chem Biol Interact* 130–132, 573–581.
- Felig, P. (1975). Amino acid metabolism in man. *Annu Rev Biochem* 44, 933–955.
- Feng, Z., Hu, W., de Stanchina, E., Teresky, A.K., Jin, S., Lowe, S., and Levine, A.J. (2007). The regulation of AMPK beta1, TSC2, and PTEN expression by p53: stress, cell and tissue specificity, and the role of these gene products in modulating the IGF-1-AKT-mTOR pathways. *Cancer Res* 67, 3043–3053.
- Folmer, V., Soares, J.C., and Rocha, J.B. (2002). Oxidative stress in mice is dependent on the free glucose content of the diet. *Int J Biochem Cell Biol* 34, 1279–1285.
- Fujimoto, M., Shimizu, N., Kunii, K., Martyn, J.A., Ueki, K., and Kaneki, M. (2005). A role for iNOS in fasting hyperglycemia and impaired insulin signaling in the liver of obese diabetic mice. *Diabetes* 54, 1340–1348.
- Gygi, S.P., Rist, B., Gerber, S.A., Turecek, F., Gelb, M.H., and Aebersold, R. (1999). Quantitative analysis of complex protein mixtures using isotope-coded affinity tags. *Nat Biotechnol* 17, 994–999.
- Halder, S.K., Anumanthan, G., Maddala, R., Mann, J., Chytil, A., Gonzalez, A.L., Washington, M.K., Moses, H.L., Beauchamp, R. D., and Datta, P.K. (2006). Oncogenic function of a novel WD-domain protein, STRAP, in human carcinogenesis. *Cancer Res* 66, 6156–6166.
- Han, P., and Chen, C. (2008). Detergent-free biotin switch combined with liquid chromatography/tandem mass spectrometry in the analysis of S-nitrosylated proteins. *Rapid Commun Mass Spectrom* 22, 1137–1145.
- Hao, G., Derakhshan, B., Shi, L., Campagne, F., and Gross, S.S. (2006). SNOSID, a proteomic method for identification of cysteine S-nitrosylation sites in complex protein mixtures. *Proc Natl Acad Sci U S A* 103, 1012–1017.
- Hess, D.T., Matsumoto, A., Kim, S.O., Marshall, H.E., and Stamler, J. S. (2005). Protein S-nitrosylation: purview and parameters. *Nat Rev Mol Cell Biol* 6, 150–166.
- Inoue, N., Yahagi, N., Yamamoto, T., Ishikawa, M., Watanabe, K., Matsuzaka, T., Nakagawa, Y., Takeuchi, Y., Kobayashi, K., Takahashi, A., *et al.* (2008). Cyclin-dependent kinase inhibitor, p21WAF1/CIP1, is involved in adipocyte differentiation and hypertrophy, linking to obesity and insulin resistance. *J Biol Chem* 283, 21220–21229.
- Jaffrey, S.R., and Snyder, S.H. (2001). The biotin switch method for the detection of S-nitrosylated proteins. *Sci. STKE* 2001, L1.
- Jung, H., Seong, H. A., and Ha, H. (2007). NM23-H1 tumor suppressor and its interacting partner STRAP activate p53 function. *J Biol Chem* 282, 35293–35307.
- Keller, A., Nesvizhskii, A.I., Kolker, E., and Aebersold, R. (2002). Empirical statistical model to estimate the accuracy of peptide identifications made by MS/MS and database search. *Anal Chem* 74, 5383–5392.
- Koh, Y.H., Suzuki, K., Che, W., Park, Y.S., Miyamoto, Y., Higashiyama, S., and Taniguchi, N. (2001). Inactivation of glutathione peroxidase by NO leads to the accumulation of H₂O₂ and the induction of HB-EGF via c-Jun NH₂-terminal kinase in rat aortic smooth muscle cells. *FASEB J* 15, 1472–1474.
- Lefièvre, L., Chen, Y., Conner, S.J., Scott, J.L., Publicover, S.J., Ford, W.C., and Barratt, C.L. (2007). Human spermatozoa contain multiple targets for protein S-nitrosylation: an alternative mechanism of the modulation of sperm function by nitric oxide? *Proteomics* 7, 3066–3084.
- Leube, R.E. (1994). Expression of the synaptophysin gene family is not restricted to neuronal and neuroendocrine differentiation in rat and human. *Differentiation* 56, 163–171.
- Li, S.Y., and Ren, J. (2008). Cardiac overexpression of alcohol dehydrogenase exacerbates chronic ethanol ingestion-induced myocardial dysfunction and hypertrophy: role of insulin signaling and ER stress. *J Mol Cell Cardiol* 44, 992–1001.
- Li, X.J., Zhang, H., Ranish, J.A., and Aebersold, R. (2003). Automated statistical analysis of protein abundance ratios from data generated by stable-isotope dilution and tandem mass spectrometry. *Anal Chem* 75, 6648–6657.
- Lindermayr, C., Saalbach, G., and Durner, J. (2005). Proteomic

- identification of S-nitrosylated proteins in Arabidopsis. *Plant Physiol* 137, 921–930.
- Maere, S., Heymans, K., and Kuiper, M. (2005). BiNGO: a Cytoscape plugin to assess overrepresentation of gene ontology categories in biological networks. *Bioinformatics* 21, 3448–3449.
- Mendel, D.B., Khavari, P.A., Conley, P.B., Graves, M.K., Hansen, L.P., Admon, A., and Crabtree, G.R. (1991). Characterization of a cofactor that regulates dimerization of a mammalian homeodomain protein. *Science* 254, 1762–1767.
- Nesvizhskii, A.I., Keller, A., Kolker, E., and Aebersold, R. (2003). A statistical model for identifying proteins by tandem mass spectrometry. *Anal Chem* 75, 4646–4658.
- Newsholme, P., Bender, K., Kiely, A., and Brennan, L. (2007). Amino acid metabolism, insulin secretion and diabetes. *Biochem Soc Trans* 35, 1180–1186.
- Nobukuni, T., Joaquin, M., Roccio, M., Dann, S.G., Kim, S.Y., Gulati, P., Byfield, M.P., Backer, J.M., Natt, F., Bos, J.L., *et al.* (2005). Amino acids mediate mTOR/raptor signaling through activation of class 3 phosphatidylinositol 3OH-kinase. *Proc Natl Acad Sci U S A* 102, 14238–14243.
- Noguchi, Y., Zhang, Q.W., Sugimoto, T., Furuhashi, Y., Sakai, R., Mori, M., Takahashi, M., and Kimura, T. (2006). Network analysis of plasma and tissue amino acids and the generation of an amino index for potential diagnostic use. *Am J Clin Nutr* 83, 513S–519S.
- Perreault, M., and Marette, A. (2001). Targeted disruption of inducible nitric oxide synthase protects against obesity-linked insulin resistance in muscle. *Nat Med* 7, 1138–1143.
- Qu, J., Liu, G.H., Wu, K., Han, P., Wang, P., Li, J., Zhang, X., and Chen, C. (2007). Nitric oxide destabilizes Pias3 and regulates sumoylation. *PLoS ONE* 2, e1085.
- Roden, M., Perseghin, G., Petersen, K. F., Hwang, J. H., Cline, G. W., Gerow, K., Rothman, D. L., and Shulman, G. I. (1996). The roles of insulin and glucagon in the regulation of hepatic glycogen synthesis and turnover in humans. *J Clin Invest* 97, 642–648.
- Rose, R.B., Pullen, K.E., Bayle, J.H., Crabtree, G.R., and Alber, T. (2004). Biochemical and structural basis for partially redundant enzymatic and transcriptional functions of DCoH and DCoH2. *Biochemistry* 43, 7345–7355.
- Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B., and Ideker, T. (2003). Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res* 13, 2498–2504.
- Sies, H., Dafré, A.L., Ji, Y., and Akerboom, T.P. (1998). Protein S-thiolation and redox regulation of membrane-bound glutathione transferase. *Chem Biol Interact* 111–112, 177–185.
- Soubeyran, P., Barac, A., Szymkiewicz, I., and Dikic, I. (2003). Cbl-ArgBP2 complex mediates ubiquitination and degradation of c-Abl. *Biochem J* 370, 29–34.
- Sugita, H., Kaneki, M., Tokunaga, E., Sugita, M., Koike, C., Yasuhara, S., Tompkins, R.G., and Martyn, J.A. (2002). Inducible nitric oxide synthase plays a role in LPS-induced hyperglycemia and insulin resistance. *Am J Physiol Endocrinol Metab* 282, E386–E394.
- Tremblay, F., and Marette, A. (2001). Amino acid and insulin signaling via the mTOR/p70 S6 kinase pathway. A negative feedback mechanism leading to insulin resistance in skeletal muscle cells. *J Biol Chem* 276, 38052–38060.
- Vailaya, A., Bluvast, P., Kincaid, R., Kuchinsky, A., Creech, M., and Adler, A. (2005). An architecture for biological information extraction and representation. *Bioinformatics* 21, 430–438.
- Wadham, C., Parker, A., Wang, L., and Xia, P. (2007). High glucose attenuates protein S-nitrosylation in endothelial cells: role of oxidative stress. *Diabetes* 56, 2715–2721.
- Wang, B., Golemis, E.A., and Kruh, G.D. (1997). ArgBP2, a multiple Src homology 3 domain-containing, Arg/Abl-interacting protein, is phosphorylated in v-Abl-transformed cells and localized in stress fibers and cardiocyte Z-disks. *J Biol Chem* 272, 17542–17550.
- Wellen, K.E., and Hotamisligil, G.S. (2005). Inflammation, stress, and diabetes. *J Clin Invest* 115, 1111–1119.
- Wijekoon, E.P., Skinner, C., Brosnan, M.E., and Brosnan, J.T. (2004). Amino acid metabolism in the Zucker diabetic fatty rat: effects of insulin resistance and of type 2 diabetes. *Can J Physiol Pharmacol* 82, 506–514.
- Xie, Q.W., Kashiwabara, Y., and Nathan, C. (1994). Role of transcription factor NF-kappa B/Rel in induction of nitric oxide synthase. *J Biol Chem* 269, 4705–4708.
- Yamagata, K., Furuta, H., Oda, N., Kaisaki, P.J., Menzel, S., Cox, N. J., Fajans, S.S., Signorini, S., Stoffel, M., and Bell, G.I. (1996a). Mutations in the hepatocyte nuclear factor-4alpha gene in maturity-onset diabetes of the young (MODY1). *Nature* 384, 458–460.
- Yamagata, K., Oda, N., Kaisaki, P.J., Menzel, S., Furuta, H., Vaxillaire, M., Southam, L., Cox, R.D., Lathrop, G.M., Boriraj, V.V., *et al.* (1996b). Mutations in the hepatocyte nuclear factor-1alpha gene in maturity-onset diabetes of the young (MODY3). *Nature* 384, 455–458.
- Yasukawa, T., Tokunaga, E., Ota, H., Sugita, H., Martyn, J.A., and Kaneki, M. (2005). S-nitrosylation-dependent inactivation of Akt/protein kinase B in insulin resistance. *J Biol Chem* 280, 7511–7518.
- Yates, J.R. 3rd, Eng, J.K., McCormack, A.L., and Schieltz, D. (1995). Method to correlate tandem mass spectra of modified peptides to amino acid sequences in the protein database. *Anal Chem* 67, 1426–1436.
- Yuan, Z.Q., Kim, D., Kaneko, S., Sussman, M., Bokoch, G.M., Kruh, G.D., Nicosia, S.V., Testa, J.R., and Cheng, J.Q. (2005). ArgBP2gamma interacts with Akt and p21-activated kinase-1 and promotes cell survival. *J Biol Chem* 280, 21483–21490.
- Zimmet, P. (2000). Globalization, coca-colonization and the chronic disease epidemic: can the Doomsday scenario be averted? *J Intern Med* 247, 301–310.