

RESEARCH ARTICLE

Origin and molecular characterization of the human-infecting H6N1 influenza virus in Taiwan

Weifeng Shi¹, Yi Shi^{2,3}, Ying Wu², Di Liu^{2,4}✉, George F. Gao^{2,3,5}

¹ School of Medical Sciences, Taishan Medical College, Taian 271016, China

² CAS Key Laboratory of Pathogenic Microbiology and Immunology, Institute of Microbiology, Chinese Academy of Sciences, Beijing 100101, China

³ Beijing Institutes of Life Sciences, Chinese Academy of Sciences, Beijing 100101, China

⁴ Network Information Center, Institute of Microbiology, Chinese Academy of Sciences, Beijing 100101, China

⁵ Office of Director-General, Chinese Center for Disease Control and Prevention (China CDC), Beijing 102206, China

✉ Correspondence: liud@im.ac.cn

Received September 11, 2013 Accepted September 22, 2013

ABSTRACT

In June 2013, the first human H6N1 influenza virus infection was confirmed in Taiwan. However, the origin and molecular characterization of this virus, A/Taiwan/2/2013 (H6N1), have not been well studied thus far. In the present report, we performed phylogenetic and coalescent analyses of this virus and compared its molecular profile/characteristics with other closely related strains. Molecular characterization of H6N1 revealed that it is a typical avian influenza virus of low pathogenicity, which might not replicate and propagate well in the upper airway in mammals. Phylogenetic analysis revealed that the virus clusters with A/chicken/Taiwan/A2837/2013 (H6N1) in seven genes, except PB1. For the PB1 gene, A/Taiwan/2/2013 was clustered with a different H6N1 lineage from A/chicken/Taiwan/A2837/2013. Although a previous study demonstrated that the PB2, PA, and M genes of A/Taiwan/2/2013 might be derived from the H5N2 viruses, coalescent analyses revealed that these H5N2 viruses were derived from more recent strains than that of the ancestor of A/Taiwan/2/2013. Therefore, we propose that A/Taiwan/2/2013 is a reassortant from different H6N1 lineages circulating in chickens in Taiwan. Furthermore, compared to avian isolates, a single P186L (H3 numbering) substitution in the hemagglutinin H6 of the human isolate might increase the mammalian receptor binding and, hence, this strain's pathogenicity in humans. Overall, human infection with this virus seems an accidental event and is unlikely to cause an influenza pandemic. However, its co-circulation and potential reassortment with other influenza subtypes are still worthy of attention.

sortment with other influenza subtypes are still worthy of attention.

KEYWORDS molecular characterization, phylogenetic analysis, coalescent analysis, H6N1, influenza virus, Taiwan

INTRODUCTION

On June 21, 2013, the Taiwan Centers for Disease Control formally confirmed the first human infection with an avian influenza A (H6N1) virus, and named it A/Taiwan/2/2013 (<http://www.cdc.gov.tw/english/info.aspx?treeid=bc2d4e89b154059b&nowtreeid=ee0a2987cfba3222&tid=E36A5E9AB3D3A216>). The patient was a 20-year-old female, diagnosed with pneumonia due to an unconfirmed type of influenza virus, which was subsequently proven to be of the H6N1 subtype. The patient has fully recovered, and no human-to-human transmission has been identified.

Avian influenza A (H6N1) virus has been circulating throughout North America (Senne, 2003) and Eurasia (Lee et al., 2006) for many years. In Taiwan, the H6N1 virus has been circulating for over 30 years and is extremely prevalent among poultry, with approximately half of the layers and 30% of the broilers having antibodies against H6N1 influenza viruses (Lee et al., 2006). In southern China, H6N1 has established itself in minor poultry species (Cheung et al., 2007) and has been proposed to be the potential progenitor of the human-infecting H5N1 influenza virus A/Hong Kong/156/97 (Hoffmann et al., 2000). Although human infection with this virus subtype was never reported prior to the Taiwanese case, H6-specific




Figure 1. Maximum likelihood phylogenetic trees of HA and NA. The human-infecting H6N1 and chicken isolates from 2013 are in red, one of the 2004/05 poultry H6N1 clade viruses is in green, and the 2009/10 poultry H6N1 viruses are in cyan. (A) HA phylogenetic tree. (B) NA phylogenetic tree.

antibodies have been detected in live animal market workers in China and in veterinarians exposed to birds in the United States (Shortridge 1992; Myers et al., 2007). Furthermore, two of 11 healthy human volunteers inoculated with the H6N1 virus displayed mild upper respiratory symptoms, suggesting that this virus has the potential to infect mammals (Beare and Webster, 1991). In addition, some Taiwanese H6N1 viruses can replicate in mice without pre-adaptation (Lee et al., 2006). Therefore, clinically asymptomatic or unreported human infections of H6 viruses may have occurred.

In the present report, we performed a molecular and phylogenetic analysis of the first laboratory-confirmed human-infecting H6N1 influenza virus and closely related virus strains. Our results revealed that *A/Taiwan/2/2013* is a low-pathogenicity H6N1 influenza virus whose genes may have been derived from different H6N1 lineages circulating in Taiwan.

RESULTS

The human-infecting H6N1 virus is a reassortant of different lineages of poultry H6N1 viruses circulating in Taiwan

To examine the origin of the human-infecting *A/Taiwan/2/2013* H6N1 virus, we performed a phylogenetic analysis of related influenza virus genomes. Except for the PB1 gene, the phylogenetic trees of the remaining genes indicated that *A/chicken/Taiwan/A2837/2013* (H6N1) is closely related to

A/Taiwan/2/2013 (Figs. 1 and 2). This suggests that the H6N1 virus in poultry is the source of the human virus, though changes have occurred. From the HA phylogeny, the 2013 viruses fell into the clade of the poultry H6N1 viruses isolated in 2004 and 2005 in Taiwan (Fig. 1A). Nevertheless, the branches between the 2013 viruses and 2004/2005 viruses were quite long, indicating the lack of surveillance data. In the NA phylogenetic tree, the 2013 isolates fell into another poultry H6N1 lineage, which consisted of isolates from 2009 to 2010 in Taiwan (Fig. 1B). This lineage is distant to that of the 2004/2005 isolates in the NA tree, indicating that the 2013 H6N1 influenza viruses were possibly reassortants of varied H6N1 lineages.

A comparison of the phylogenetic trees of the internal genes showed that all but the PA gene could be grouped with the 2009/2010 isolates and close to *A/Taiwan/2/2013*. The closest PA gene to the 2013 isolates was that from *A/chicken/Taiwan/0101/2012* (H5N2) (Fig. 2). Then, the clade clustered with earlier H6N1 strains in Taiwan.

Co-circulating H5N2 influenza viruses reassorted with H6N1 viruses in poultry

Consistent with recently published data (Yuan et al., 2013), in the PB2, PA, and M trees, the closest strains of *A/Taiwan/2/2013* and *A/chicken/Taiwan/A2837/2013* belonged to the H5N2 subtype rather than H6N1 (Figs. 2 and 3), with the closest strains being *A/chicken/Taiwan/0101/2012* (H5N2) for PB2 and PA, and *A/chicken/Taiwan/A1997/2012* (H5N2) for M.




Figure 2. Phylogenetic trees of the six internal genes. The trees were generated by the maximum likelihood methods with 1000x bootstrap analyses. The Taiwan 2013 H6N1 isolates are in red, while the 2009/2010 H6N1 isolates are in cyan. The subtypes other than H6N1 are noted.

Table 1. tMRCAs of the two H6N1 strains isolated in 2013 in Taiwan and their closely related H5N2 viruses

Gene	Sequences ^a	Models			Burn-in ^b
		Constant size	Logistic growth	Exponential growth	
PB2	a + b	Aug, 2010 (Dec, 2008, Dec, 2011)	Oct, 2009 (Feb, 2008, Jul, 2011)	May, 2010 (Oct, 2008, Dec, 2011)	4 million
	a + b + c	May, 2008 (Apr, 2006, Apr, 2010)	Jul, 2007 (Jul, 2005, Jul, 2009)	Dec, 2007 (Feb, 2006, Apr, 2010)	4 million
PA	a + b	Jan, 2011 (Dec, 2009, Dec, 2011)	Dec, 2010 (Oct, 2009, Nov, 2011)	Dec, 2010 (Jan, 2010, Dec, 2011)	2 million
	a + b + c	Jun, 2007 (Sep, 2005, Mar, 2009)	Apr, 2007 (Jul, 2005, Feb, 2009)	May, 2007 (Aug, 2005, Feb, 2009)	2 million
M	a + b	Oct, 2010 (Sep, 2008, Sep, 2012)	Aug, 2010 (Apr, 2008, Sep, 2012)	Sep, 2010 (Jun, 2008, Aug, 2012)	4 million
	a + b + d	Jun, 2008 (Jan, 2006, Feb, 2011)	Mar, 2008 (Jun, 2005, Mar, 2011)	June, 2008 (Sep, 2005, Jan, 2011)	4 million

^a In this column, “a”, “b”, “c”, and “d” represent A/Taiwan/2/2013 (H6N1), A/chicken/Taiwan/A2837/2013 (H6N1), A/chicken/Taiwan/0101/2012 (H5N2), and A/chicken/Taiwan/A1997/2012 (H5N2), respectively.

^b Bayesian Markovchain Monte Carlo analysis was run for 40 million steps, but different steps were removed as burn-in.




Figure 3. BEAST trees of the PB2, PA, and M genes of the human-infecting H6N1 and related viruses. The isolates from 2013 are in red. The H5 subtype viruses are in cyan. The bars represent the 95% highest posterior density of the estimation of the dates.




Figure 4. Structural modeling of the receptor binding site and sequence alignment of the different sites of the HA gene. (A) Receptor binding sites of two isolates from 2013 and the ancestral virus *A/shearwater/Australia/1/1972*. (B) Amino acid sites that differ from the 2004/2005 H6N1 viruses or 2009/2010 H6N1 viruses. The arrowheads point to the position in *A/Taiwan/2/2013* that differs from all other strains.

However, the H5N2 viruses did not form independent lineages in the trees (Fig. 3). Instead, they scattered within H6N1 lineages that have been circulating in Taiwan for over 10 years. Moreover, the branch length between the H6N1 Taiwanese strains from 2013 and their H5N2 relatives was very long.

To further test whether the PB2, PA, and M genes of *A/Taiwan/2/2013* and *A/chicken/Taiwan/A2837/2013* originated from H5N2 strains, we performed a coalescent analysis and calculated the estimated time to most recent common ancestor (tMRCA) for these viruses. The tMRCA of the two H6N1 strains was estimated as early 2011 for the PA gene and before 2011 for the PB2 and M genes (Table 1), whereas the two H5N2 viruses were isolated in 2012, suggesting that they are more recent strains than the ancestor of the novel human-infecting H6N1 virus. Furthermore, the tMRCA of the two H6N1 strains and the corresponding H5N2 strains was roughly 2008 for the PB2 and M genes, and 2007 for the PA gene (Table 1), also suggesting that recent reassortment from H5N2 to H6N1 is unlikely.

The proline to leucine substitution at the receptor-binding site tends to increase the affinity to bind mammalian receptors

To investigate the molecular basis for human infectivity and

pathogenicity with Taiwanese H6N1, we compared the receptor-binding sites of the 2013 isolates with an earlier avian strain *A/shearwater/Australia/1/1972* (Fig. 4A). At the receptor-binding site, the human isolate had a proline to leucine substitution at position 186 (P186L, H3 numbering), which was not observed in the earlier strains or the recent circulating H6N1 lineages. Because of the stronger hydrophobicity of leucine, this substitution is hypothesized to increase the hydrophobicity at the receptor-binding site and thus make HA more likely to bind the mammalian receptor (Shi et al. 2013, Xiong et al. 2013). This may explain how this H6N1 virus was able to infect humans even though position 226 is a glutamine. The mechanism of the receptor binding switch for the H6N1 virus may be different from that for the H5N1 virus (Zhang et al. 2013). Moreover, when compared to the receptor-binding site of *A/shearwater/Australia/1/1972*, the 2013 isolates also contained an E190V substitution, which could also increase the hydrophobicity at the receptor-binding site. These data indicate that the 2013 isolates have evolved from an earlier avian isolate to adapt to human infection.

Aside from the P186L substitution in HA, there is only one difference at position 287 that *A/Taiwan/2/2013* encodes relative to the poultry strains (Fig. 4B). However, whether the A287T substitution increases the binding to mammalian recep-

eral key features that could influence the pathogenesis of the virus, including the presence of the avian-signature Gln226 in the HA receptor-binding site, the lack of multiple basic amino acids at the cleavage site, two deletions in the NA stalk, and Glu627 and Asp701 in PB2 (avian-signatures). Therefore, A/Taiwan/2/2013 tends to be of low pathogenicity, prefers to infect avian hosts, and has adapted well within terrestrial poultry. The Arg292 in NA and Ser31 in M2 indicate that the virus is sensitive to oseltamivir but insensitive to adamantane. The molecular features of the novel H6N1 virus suggest that it is not able to replicate well in human cells and cannot be efficiently transmitted within mammalian hosts (Glu627 and Asp701 in PB2). Nevertheless, the P186L substitution has provided an increased hydrophobic environment in the sialic acid receptor-binding pocket of HA, and this change possibly enhances its ability to bind to the human receptor. Hence, the receptor binding affinity of the novel human-infecting H6N1 virus requires further examination in the future. In light of this molecular characterization it is reasonable to assume that the single human infection with this virus in Taiwan was most likely an unfortunate rare event.

A recent publication proposes that A/Taiwan/2/2013 (H6N1) originated from a reassortment between H5N2 (PB2, PA, and M) and H6N1 subtypes based on phylogenetic analyses (Yuan et al., 2013). However, the formation of the novel H6N1 influenza viruses from H5N2 viruses was not fully supported by our phylogenetic and coalescent analyses. First, based on surveillance data, the H5N2 viruses did not form independent lineages in the PB2, PA, and M trees. On the contrary, they fell within an H6N1 lineage circulating in Taiwan for approximately 10 years, indicating that the H5N2 viruses gained gene segments from H6N1 viruses at an earlier time. Second, the long branch between the H6N1 Taiwanese strains from 2013 and the H5N2 relative suggests that unknown evolutionary events may have occurred between the H5N2 to H6N1 viruses, and therefore, direct reassortment is unlikely to have occurred. Finally, coalescent analysis showed that the H5N2 viruses (isolated from 2012) were more recent strains than the ancestor of the novel human-infecting H6N1 virus. This indicates that the ancestral strain transferred the gene to both H5N2 and H6N1 viruses, and from the view of the phylogenetic trees, the ancestor strain is more likely to be an H6 than an H5 virus. Therefore, we hypothesize that A/Taiwan/2/2013 is likely a reassortant from different H6N1 lineages.

Although the emergence of human-infecting H6N1 viruses in Taiwan is unlikely to be due to direct reassortment from H5N2 viruses, the co-circulation of H6N1 and H5N2 viruses in Taiwan still requires close surveillance. Analysis of phylogenetic trees indicates that distinct H6-H5 reassortment events have occurred during the past 10 years. The reassortment of varied subtypes of influenza viruses has the potential to cause rapid adaptation and lead to an unexpected evolutionary road (Liu et al., 2013; Morens et al., 2013). Therefore, extensive surveillance is needed in the future.

MATERIALS AND METHODS

The genome sequences of A/Taiwan/2/2013 (H6N1) and A/chicken/Taiwan/A2837/2013 (H6N1) were obtained from the Global Initiative on Sharing Avian Influenza Data (GISAID) database, and closely related sequences were downloaded from BLAST searches against GenBank. Multiple sequence alignment for each gene was performed using MUSCLE (Edgar, 2004). Phylogenetic trees were reconstructed using RAxML (Stamatakis, 2006) using the GTRGAMMA model.

The tMRCAs were estimated using the Bayesian Markov chain Monte Carlo method, implemented in BEAST (Drummond and Rambaut, 2007). To investigate the extent to which dating estimates are affected by the demographic model chosen, we repeated our analyses using constant size, logistic growth, and exponential growth models, respectively. Bayesian Markov chain Monte Carlo analyses were run for 40 million steps, but different steps were removed as burn-in. Trees and other parameters were sampled every 10,000 steps.

ACKNOWLEDGEMENTS

This study was supported by the National Basic Research Program (973 Program) (Nos. 2010CB530303, 2011CB504703, and 2012CB955501) and an intramural special grant for influenza virus research from the Chinese Academy of Sciences (KSZD-EW-Z-002). WS was supported by the Doctoral Starting up Foundation of Taishan Medical College. GFG is a leading principal investigator of the Innovative Research Group of the National Natural Science Foundation of China (Grant No. 81021003). We thank the scientists from Taiwan who submitted the nucleotide sequences of the novel avian H6N1 influenza viruses to GISAID database, Joel Haywood for critical reading and English editing, and Dr. Haixia Xiao for helpful discussion.

COMPLIANCE WITH ETHICS GUIDELINES

Weifeng Shi, Yi Shi, Ying Wu, Di Liu, George F. Gao declare that there is no conflict of interest.

This article does not contain any studies with human or animal subjects performed by any of the authors.

REFERENCES

- Beare, A.S., and Webster, R.G. (1991). Replication of avian influenza viruses in humans. *Arch Virol* 119, 37–42.
- Chen, W., Calvo, P.A., Malide, D., Gibbs, J., Schubert, U., Bacik, I., Basta, S., O'Neill, R., Schickli, J., Palese, P., et al. (2001). A novel influenza A virus mitochondrial protein that induces cell death. *Nat Med* 7, 1306–1312.
- Cheung, C.L., Vijaykrishna, D., Smith, G.J., Fan, X.H., Zhang, J.X., Bahl, J., Duan, L., Huang, K., Tai, H., Wang, J., et al. (2007). Establishment of influenza A virus (H6N1) in minor poultry species in southern China. *J Virol* 81, 10402–10412.
- Drummond, A.J., and Rambaut, A. (2007). BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evol Biol* 7, 214.
- Edgar, R.C. (2004). Muscle: Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32, 1792–1797.
- Ha, Y., Stevens, D.J., Skehel, J.J., and Wiley, D.C. (2001). X-ray structures of H5 avian and H9 swine influenza virus hemagglutinins bound to avian and human receptor analogs. *Proc Natl Acad Sci U S A* 98, 11181–11186.

- Hatta, M., Gao, P., Halfmann, P., and Kawaoka, Y. (2001). Molecular basis for high virulence of Hong Kong H5N1 influenza A viruses. *Science* 293, 1840–1842.
- Hoffmann, E., Stech, J., Leneva, I., Krauss, S., Scholtissek, C., Chin, P.S., Peiris, M., Shortridge, K.F., and Webster, R.G. (2000). Characterization of the influenza A virus gene pool in avian species in southern China: was H6N1 a derivative or a precursor of H5N1? *J Virol* 74, 6309–6315.
- Holsinger, L., Nichani, D., Pinto, L., and Lamb, R. (1994). Influenza A virus M2 ion channel protein: a structure-function analysis. *J Virol* 68, 1551–1563.
- Kiso, M., Ozawa, M., Le, M.T., Imai, H., Takahashi, K., Kakugawa, S., Noda, T., Horimoto, T., and Kawaoka, Y. (2011). Effect of an asparagine-to-serine mutation at position 294 in neuraminidase on the pathogenicity of highly pathogenic H5N1 influenza A virus. *J Virol* 85, 4667–4672.
- Labadie, K., Dos Santos Afonso, E., Rameix-Welti, M.A., van der Werf, S., Naffakh, N. (2007). Host-range determinants on the PB2 protein of influenza A viruses control the interaction between the viral polymerase and nucleoprotein in human cells. *Virology* 362, 271–282.
- Lee, M.S., Chang, P.C., Shien, J.H., Cheng, M.C., Chen, C.L., and Shieh, H.K. (2006). Genetic and pathogenic characterization of H6N1 avian influenza viruses isolated in Taiwan between 1972 and 2005. *Avian Dis* 50, 561–571.
- Liu, D., Shi, W., Shi, Y., Wang, D., Xiao, H., Li, W., Bi, Y., Wu, Y., Li, X., Yan, J., et al. (2013). Origin and diversity of novel avian influenza A H7N9 viruses causing human infection: phylogenetic, structural, and coalescent analyses. *Lancet* 381, 1926–1932.
- Massin, P., van der Werf, S., and Naffakh, N. (2001). Residue 627 of PB2 is a determinant of cold sensitivity in RNA replication of avian influenza viruses. *J Virol* 75, 5398–5404.
- Matrosovich, M., Zhou, N., Kawaoka, Y., and Webster, R.G. (1999). The surface glycoproteins of H5 influenza viruses isolated from humans, chickens and wild aquatic birds have distinguishable properties. *J Virol* 73, 1146–1155.
- McAuley, J.L., Hornung, F., Boyd, K.L., Smith, A.M., McKeon, R., Benink, J., Yewdell, J.W., and McCullers, J.A. (2007). Expression of the 1918 influenza A virus PB1-F2 enhances the pathogenesis of viral and secondary bacterial pneumonia. *Cell Host Microbe* 2, 240–249.
- Meunier, I., and von Messling, V. (2012). PB1-F2 modulates early host responses but does not affect the pathogenesis of H1N1 seasonal influenza virus. *J Virol* 86, 4271–4278.
- Morens, D.M., Taubenberger, J.K., and Fauci, A.S. (2013). Pandemic influenza viruses—hoping for the road not taken. *N Engl J Med* 368, 2345–2348.
- Myers, K.P., Setterquist, S.F., Capuano, A.W., and Gray, G.C. (2007). Infection due to 3 avian influenza subtypes in United States veterinarians. *Clin Infect Dis* 45, 4–9.
- Pinto, L., Holsinger, L., and Lamb, R. (1992). Influenza virus M2 protein has ion channel activity. *Cell* 69, 517–528.
- Senne, D.A. (2003). Avian influenza in the Western Hemisphere including the Pacific Islands and Australia. *Avian Dis* 47, 798–805.
- Shi, Y., Zhang, W., Wang, F., Qi, J., Wu, Y., Song, H., Gao, F., Bi, Y., Zhang, Y., Fan, Z., et al. (2013). Structures and Receptor Binding of Hemagglutinins from Human-Infecting H7N9 Influenza Viruses. *Science*. (In Press).
- Shortridge, K.F. (1992). Pandemic influenza: a zoonosis? *Semin Respir Infect* 7, 11–25.
- Stamatakis, A. (2006). RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22, 2688–2690.
- Xiong, X., Martin, S.R., Haire, L.F., Wharton, S.A., Daniels, R.S., Bennett, M.S., McCauley, J.W., Collins, P.J., Walker, P.A., Skehel, J.J., et al. (2013). Receptor binding by an H7N9 influenza virus from human. *Nature* 499, 496–499.
- Yuan, J., Zhang, L., Kan, X., Jiang, L., Yang, J., Guo, Z., and Ren, Q. (2013). Origin and molecular characteristics of a novel 2013 avian influenza A H6N1 virus causing human infection in Taiwan. *Clin Infect Dis*. (In Press).
- Zamarin, D., Ortigoza, M.B., and Palese, P. (2006). Influenza A virus PB1-F2 protein contributes to viral pathogenesis in mice. *J Virol* 80, 7976–7983.
- Zell, R., Krumbholz, A., Eitner, A., Krieg, R., Halbhuber, K.J., and Wutzler, P. (2007). Prevalence of PB1-F2 of influenza A viruses. *J Gen Virol* 88, 536–546.
- Zhang, W., Shi, Y., Lu, X., Shu, Y., Qi, J., and Gao, G.F. (2013). An airborne transmissible avian influenza H5 hemagglutinin seen at the atomic level. *Science* 340, 1463–1467.