CORRECTION

Correction to: Efficient derivation of extended pluripotent stem cells from NOD-*scid* *Il2rg*^{-/-} mice

Yaqin Du¹, Ting Wang¹, Jun Xu¹, Chaoran Zhao¹, Haibo Li¹, Yao Fu³, Yaxing Xu², Liangfu Xie³, Jingru Zhao², Weifeng Yang⁴, Ming Yin⁴, Jinhua Wen¹✉, Hongkui Deng^{1,2,3}✉

¹ Peking University Stem Cell Research Center, Department of Cell Biology, School of Basic Medical Sciences, Peking University Health Science Center, Beijing 100191, China

² Peking University-Tsinghua University-National Institute of Biological Sciences Joint Graduate Program, College of Life Sciences, Peking University, Beijing 100871, China

³ The MOE Key Laboratory of Cell Proliferation and Differentiation, College of Life Sciences, Peking-Tsinghua Center for Life Sciences, Peking University, Beijing 100871, China

⁴ Beijing Vitalstar Biotechnology, Beijing 100012, China

✉ Correspondence: jhwen@bjmu.edu.cn (J. Wen), hongkui_deng@pku.edu.cn (H. Deng)

CORRECTION TO: PROTEIN CELL

[HTTPS://DOI.ORG/10.1007/S13238-018-0558-Z](https://doi.org/10.1007/s13238-018-0558-z)

In the original publication Fig. 1D and supplementary material is incorrect. The correct Fig. 1D and supplementary material is provided in this correction.

The original article can be found online at <https://doi.org/10.1007/s13238-018-0558-z>.

Electronic supplementary material The online version of this article (doi:<https://doi.org/10.1007/s13238-018-0571-2>) contains supplementary material, which is available to authorized users.

Figure 1. Generation of NOD-scid *Il2rg*^{-/-} extended pluripotent stem cells. (A) Schematic of two approaches used for generating NOD-scid *Il2rg*^{-/-} extended pluripotent stem cells: *de novo* derivation from blastocysts (upper panels) and chemical reprogramming from embryonic fibroblasts (lower panels). (B) Phase-contrast images of *de novo* derived outgrowth and EPS colonies for 17 passages in LCDM medium. Scale bars, 100 μ m. (C) qRT-PCR analysis of XEN marker genes expression during the chemical induction process (day 16). Error bars indicate SEM ($n = 2$). (D) Co-immunostaining of XEN marker genes during the chemical induction process (day 16). Upper panels: GATA6 and SALL4; lower panels: SOX17 and SALL4. Scale bars, 100 μ m. (E) Immunofluorescence of OCT4-positive primary colonies at the end of the chemical induction (day 40). Scale bars, 100 μ m. (F) Typical karyotypes of EPS (passage 22) and cEPS (passage 23) cells. Each cell line counts 30 cells.

OPEN ACCESS

This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and

reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.