



Marked Spatial Point Processes: Current State and Extensions to Point Processes on Linear Networks

Matthias ECKARDT^{id} and Mehdi MORADI^{id}

Within the applications of spatial point processes, it is increasingly becoming common that events are labelled by marks, prompting an exploration beyond the spatial distribution of events by incorporating the marks in the undertaken analysis. In this paper, we first consider marked spatial point processes in \mathbb{R}^2 , where marks are either integer-valued, real-valued, or object-valued, and review the state-of-the-art to analyze the spatial structure and type of interaction/correlation between marks. More specifically, we review cross/dot-type summary characteristics, mark-weighted summary characteristics, various mark correlation functions, and frequency domain approaches. We also propose novel cross/dot-type higher-order summary characteristics, mark-weighted summary characteristics, and mark correlation functions for marked point processes on linear networks. Through a simulation study, we show that ignoring the underlying network gives rise to erroneous conclusions about the interaction/correlation between marks. Finally, we consider two applications: the locations of two different proteins on the membranes of cells infected with the influenza virus and the locations of public trees along the street network of Vancouver, Canada, where trees are labelled by their diameters at breast height.

Key Words: Cross-type summary characteristics; Influenza virus; Mark correlation functions; Mark-weighted summary characteristics; Point spectra; Public street trees

1. INTRODUCTION

Significant advancements in data collection and storage capacities have led to vast point pattern data availability from diverse sources, which often give access to (precise) spatial

A commentary to this article is available at <https://doi.org/10.1007/s13253-024-00609-x>, <https://doi.org/10.1007/s13253-024-00608-y>, <https://doi.org/10.1007/s13253-024-00610-4>, <https://doi.org/10.1007/s13253-024-00606-0>, <https://doi.org/10.1007/s13253-024-00607-z>.

M. Eckardt, Chair of Statistics, Humboldt-Universität zu Berlin, Berlin, Germany
(E-mail: m.eckardt@hu-berlin.de).

M. Moradi (✉), Department of Mathematics and Mathematical Statistics, Umeå University, Umeå, Sweden
(E-mail: mehdi.moradi@umu.se).

© 2024 The Author(s)
Journal of Agricultural, Biological, and Environmental Statistics
<https://doi.org/10.1007/s13253-024-00605-1>

locations and time occurrences of events together with further valuable point-specific information, i.e., marks. In some cases, spatial locations might be restricted by the entities of some spatially embedded relational systems, which consequently limit where events could occur. Thus, the state space that accommodates the events plays a vital role in advanced statistical analysis. Not surprisingly, the growing availability and accessibility of point pattern data, which find applications in distinct scientific fields, generally stimulated an increasing interest in developing suitable statistical/mathematical tools for the analysis of point patterns. Within the literature, applications often include astronomy (Feigelson and Babu 2012), zoology (Russell et al. 2016), ecology (Daniel et al. 2020), forestry (Gavrikov and Stoyan 1995; Yazigi et al. 2021), geology (Flagg et al. 2020), and health (Bayisa et al. 2023). Within these applications, typical examples of marks, which are point-specific, include, e.g., the shape of galaxies, animal sightings of (non)invasive species, the diameter of trees at breast height (dbh), habitat characteristics, the magnitude of earthquakes, and exposure levels to pollutants. In any such application, the interesting aim is not only to make inferences about the spatial distribution of events and their potential interaction but also to understand the association among the corresponding marks. For instance, in the case of forestry and ecology, spatial variation of dbh measurements for pairs of distinct trees and spatial correlation between sighting locations for pairs of different animal species might be interesting. Note that understanding how the dbh values differ across spatial locations might provide valuable insights into tree growth patterns, resource distribution, and ecological interactions within the forest ecosystem. Also, studying sighting locations might uncover relationships and dependencies between animal distributions, shedding light on potential habitat preferences, interspecies dynamics, and ecological coexistence patterns. These objectives are usually addressed by employing different so-called mark summary characteristics, which are expected to account for the specificity of the marks and, potentially, the constrained spatial domain of the points. This paper discusses the existing methodologies and state-of-the-art for marked spatial point processes in \mathbb{R}^2 and proposes some novelties in the context of point processes on linear networks. Throughout the paper, we consider three types of marks which are frequently seen in practice, namely, qualitative marks, quantitative marks, and non-scalar marks. In what follows, these are referred to as *discrete and integer-valued marks*, *real-valued marks*, and *object-valued marks*, respectively.

The literature for spatial point processes usually employs diverse exploratory tools, such as summary characteristics, to e.g., investigate both pairwise and higher-order interactions between points and/or marks, as well as to validate fitted models. As a general construction principle, any such summary characteristic quantitatively assesses the average interrelation between points and/or marks within a specific interpoint distance. Initially, the main attention was paid to unmarked stationary and homogeneous point processes, where the spatial distribution of points is translation invariant, and the intensity function remains constant over the corresponding state space, leading to the development of the K -function (Ripley 1976), used to study the pairwise relationships between points, and the J -function (van Lieshout and Baddeley 1999), which goes beyond the second-order analysis. Both these summary characteristics are used to identify clustering and/or inhibition among points. These were later extended to marked stationary point processes, developing cross/dot-type second/higher-order summary characteristics for point processes with integer-valued marks (Lotwick and

Silverman 1982; van Lieshout 2006), and mark-weighted second-order summary characteristics for real-valued marks (Penttinen et al. 1992; Schlather 2001). A different line of research focused on the pairwise association/variation of real-valued marks, including the mark variogram (Cressie 1993; Walder and Stoyan 1996; Stoyan and Walder 2000), which has similarities with the (semi-)variogram commonly used in geostatistical contexts, and Stoyan’s mark correlation function (Stoyan and Stoyan 1994). Similar to the classic point process characteristics, any such tool is constructed based on the marks for pairs of points at arbitrary interpoint distances. While, in practice, points might exhibit inhomogeneity, most of these summary characteristics for real-valued marks are only defined for stationary point processes, and their inhomogeneous versions remain an open topic for future research. Other approaches to investigating the dependencies between marks and spatial locations are proposed by Schlather et al. (2004); Guan (2006); Guan and Afshartous (2007). Moreover, some recent developments for the analysis of stationary spatial point processes, which simultaneously possess both integer-valued and real-valued marks, include some graphical model approaches, and partial characteristics proposed by Eckardt and Mateu (2019a,b). Instead of considering a spatial domain perspective, they applied a frequency domain representation to the marked points to compute different (partial) spectral density characteristics. This highlights the energy distribution of the point patterns over a range of frequencies, unlike the summary characteristics, which focus on pointwise spatial interactions. Although frequency domain methods offer a highly flexible and computationally efficient way to investigate the structural interrelation of complex marked point processes, methodological contributions and practical applications within the spatial point process literature remain limited.

Turning back to unmarked point processes, Baddeley et al. (2000) and van Lieshout (2011) extended the second- and higher-order summary characteristics, i.e., the K - and J -functions, to inhomogeneous settings for particular classes of point processes, namely second-order intensity-reweighted stationary and intensity-reweighted moment stationary processes; the former is a particular case of the latter. The extended versions of these summary characteristics to take marks into account are proposed by Moller and Waagepetersen (2003); Cronie and van Lieshout (2016); Ghorbani et al. (2021). Note that one could, based on some rules, construct different mark subsets out of real-valued marks and employ cross/dot-type summary characteristics (Illian et al. 2008).

In an effort to study non-integer/real-valued marks, the focus is directed toward proposing novel methodologies capable of handling diverse forms of marks. More specifically, by borrowing ideas from functional data analysis (Ramsay and Silverman 1997), extensions of Stoyan’s mark correlation function (Stoyan and Stoyan 1994) to function-valued marks are proposed by Comas et al. (2011, 2013) for stationary point processes. Ghorbani et al. (2021) proposed a framework for functional marked point processes together with some mark-weighted reduced moment measures. Moreover, Eckardt et al. (2023) and Eckardt et al. (2023) extended some summary characteristics for spatial point processes with integer-valued marks to the case of multivariate point processes with multivariate function-valued marks and constrained vector-valued quantities. These non-scalar valued marks are summarized into the class of object-valued marks, where instead of a scalar mark, each point is augmented by a non-scalar mark, i.e., an object-valued attribute, which lives on a suitable mark space whose precise form depends on the object under study. Suitable choices for the

mark space include the Banach/Hilbert space for function-valued marks and the simplex for constrained vector-valued quantities. As in standard mark point process investigations, these extended tools aim to explore the spatial variation/association of the specific objects, e.g., curves, for pairs of points at arbitrary distances.

In the past two decades, significant attention has been given to point processes on linear networks, where spatial locations of events are limited to a linear network. [Okabe and Yamada \(2001\)](#) and [Xie and Yan \(2008\)](#) proposed a network-based version of Ripley's K -function and a kernel-based intensity estimator, which replaced the Euclidean distance with the shortest-path distance without taking the geometry of the underlying network into account, leading to biased results; note that the underlying network itself often has a non-uniform distribution. As for taking the geometry of the underlying network into account, novel methodologies focusing on intensity function and summary characteristics were developed. Utilizing shortest-path distances, [Ang et al. \(2012\)](#) proposed geometrically corrected second-order summary characteristics for both homogeneous and inhomogeneous point processes on linear networks. The developed framework by [Ang et al. \(2012\)](#) was subsequently expanded to the case of multivariate/multitype point processes on linear networks, taking integer-valued marks into account ([Baddeley et al. 2014](#)), and spatio-temporal point processes on linear networks ([Moradi and Mateu 2020](#)). Moreover, [Rakshit et al. \(2017\)](#) proposed the consideration of regular distances on linear networks and defined novel versions of the summary characteristics proposed by [Ang et al. \(2012\)](#) based on a more general class of metrics. Later, employing regular distances, [Cronie et al. \(2020\)](#) proposed higher-order summary characteristics for point processes on linear networks. In terms of intensity functions, various techniques are proposed, which are either based on kernel functions ([McSwiggan et al. 2017](#); [Moradi et al. 2018](#)) or Voronoi tessellations ([Moradi et al. 2019](#); [Mateu et al. 2020](#)). Computational issues became evident on large networks, leading to a fast kernel-based intensity estimator ([Rakshit et al. 2019](#)) and an efficient way to compute the K -functions for point processes on linear networks ([Rakshit et al. 2019](#)). Excluding the second-order summary characteristics for multivariate/multitype (integer-valued marks) point processes on linear networks, proposed by [Baddeley et al. \(2014\)](#), and a kernel-based smoothing approach for scalar marks on linear networks by [Rakshit et al. \(2019\)](#), the literature certainly lacks mark-based methodologies to analyze events occurring on linear networks. In particular, no specific contributions go beyond integer-valued marks.

Knowing the limitations mentioned above, we propose novel methodologies for spatial point processes on linear networks that possess either integer- or real-valued marks. More specifically, in the case of real-valued marks, we present extensions of various marked summary characteristics, e.g., Stoyan's mark correlation and mark-weighted summary characteristics, to linear network settings. In the cases of multivariate/multitype point processes on linear networks, we take the higher-order summary characteristics, defined by [Cronie et al. \(2020\)](#), and propose their cross/dot-type versions, which are of great use to reveal the type of interactions between points with different marks; these can also be used to study real-valued marks by defining mark subsets. In Sect. 2, we present two motivating datasets, one on a planar state space and one on a linear network. We then, in Sect. 3, provide a detailed overview of the state-of-the-art for the analysis of marked spatial point processes in \mathbb{R}^2 . In particular, we start by presenting various summary characteristics for inhomogeneous

geneous multivariate/multitype point processes and then present several mark summary characteristics for stationary point processes with real-valued marks. This section further covers mark characteristics for object-valued marks as well as distinct frequency domain methods. Section 4 starts by reviewing second-order summary characteristics for inhomogeneous multivariate/multitype point processes on linear networks together with defining their mark-weighted versions and then proposes higher-order summary characteristics for such point processes. It then extends several mark summary characteristics for point processes with real-valued marks to settings where events happen on linear networks, accompanied by a numerical evaluation highlighting the importance of considering the underlying networks. Lastly, in Sect. 5, we present the results obtained from the analysis of two considered datasets and close the paper with a discussion in Sect. 6.

2. DATA

To illustrate the use of both the existing methodologies and our novel contributions, we consider two spatial point pattern datasets sourced from publicly accessible open data repositories. These datasets are locations of two different proteins on the membranes of cells infected with the influenza virus (Fig. 1), and data concerning public street trees in Vancouver, Canada (Fig. 2).

The influenza virus protein data are accessible through the R package `spatstat.data` (Baddeley et al. 2015), provided by Chen et al. (2008). It gives replicated spatial point patterns of the locations of two different proteins on the membranes of cells infected with the influenza virus. The data contain 41 replicates wherein marked point patterns concern the association of matrix proteins and glycoproteins of the viral membrane - either on matrix proteins 1 (M1) and 2 (M2) or M2 and the hemagglutinin (HA) proteins. While the M2 governs the pH during viral maturation and is essential for virus replication, the HA spike protein is regularly spaced in the membrane and is the antigenic determinant of the influenza virus. We here only focus on the 37th pattern, which consists of 845 locations of the two proteins M2 (303 points) and HA (542 points).

Figure 2 presents the spatial distribution of public trees, excluding park trees, located along the street network of Vancouver, Canada, in 2016. Attributes such as tree species and their corresponding diameter at breast height (dbh) in inches are attached to each location. The original data is sourced from the open data portal¹ of Vancouver, Canada, and consists of 136,574 places of public trees categorized into 282 different species. However, we only consider five species *Aquifolium*, *Arnold*, *Bignonioides*, *Involucrata*, and *Populus*, which together include 1045 trees with a dbh varying between 2 and 94 inches. The street network comprises 49,928 vertices and 55,221 segments and spans a total length of 1,779.547 km.

¹<https://opendata.vancouver.ca>




Figure 1. Locations of two different proteins, M2 and HA, on the membranes of cells infected with influenza virus.




Figure 2. Public street trees in Vancouver, Canada, in 2016. Left: different shapes show species of trees; right: colors show diameter at breast height in inches .

3. MARKED SPATIAL POINT PROCESSES IN \mathbb{R}^2

Let $\mathbf{x} = \{x_i, m(x_i)\}_{i=1}^n$ be an observed finite marked point pattern within a window $W \subset \mathbb{R}^2$, where x_i is the spatial location of i -th event and $m(x_i)$ is some associated mark living on a complete separable metric space \mathbb{M} . It is assumed that \mathbf{x} has been generated by an underlying random mechanism X , called a marked spatial point process, on the product space $\mathbb{R}^2 \times \mathbb{M}$, where \mathbb{R}^2 is equipped with the spatial distance $d(u, v) = \|u - v\|$, $u, v \in \mathbb{R}^2$, and the Lebesgue measure $|\cdot|$. There is also an appropriate reference measure on the mark

space, most notably the counting measure when we deal with multitype point processes. For an unmarked version of X , denoted by \check{X} , it is said that \check{X} is a homogeneous point process with constant intensity λ if $\mathbb{E}[N(\check{X} \cap A)] = \lambda|A|$, $A \subset \mathbb{R}^2$, where N is a count function. Otherwise, \check{X} is an inhomogeneous process, and

$$\mathbb{E}[N(\check{X} \cap A)] = \int_A \lambda(u)du, \quad A \subset \mathbb{R}^2, \quad (1)$$

so that $\lambda(\cdot)$ governs the spatial distribution of points. Heuristically, $\lambda(u)|du|$, $u \in \mathbb{R}^2$, is the probability of finding a point of \check{X} within a vicinity of u of size $|du|$. From now on, whenever we refer to the intensity function, we are specifically referring to the intensity function of an unmarked point process. Furthermore, according to Campbell's formulas, for any non-negative measurable function $f : \mathbb{R}^{2m} \rightarrow \mathbb{R}$,

$$\mathbb{E} \left[\sum_{x_1, \dots, x_m \in \check{X}}^{\neq} f(x_1, \dots, x_m) \right] = \int_{\mathbb{R}^{2m}} f(u_1, \dots, u_m) \lambda^{(m)}(u_1, \dots, u_m) du_1 \cdots du_m, \quad (2)$$

where $\lambda^{(m)}(\cdot)$ is called the m -th order product intensity of \check{X} , and \neq means that the sum is taken over distinct m -tuples. Note that (1) is a particular case of (2) when $f(x) = \mathbf{1}\{x \in A\}$ where $\mathbf{1}$ is an indicator function. In a similar manner, $\lambda^{(m)}(u_1, \dots, u_m) \prod_{i=1}^m |du_i|$ is the probability of having points of \check{X} in m infinitesimal disjoint areas around u_1, \dots, u_m with sizes $|du_1|, \dots, |du_m|$.

As the product intensity functions of \check{X} are not designed to study the correlation among points, if all m -th order product intensities exist, one may wish to employ the correlation function

$$g_m(u_1, \dots, u_m) = \frac{\lambda^{(m)}(u_1, \dots, u_m)}{\lambda(u_1) \cdots \lambda(u_m)} = \sum_{j=1}^m \sum_{D_1, \dots, D_j} \xi_{N(D_1)}(\{u_i : i \in D_1\}) \cdots \xi_{N(D_j)}(\{u_i : i \in D_j\}), \quad (3)$$

where \sum_{D_1, \dots, D_j} ranges over all partitions $\{D_1, \dots, D_j\}$ of $\{1, \dots, m\}$ into j non-empty and disjoint sets, $N(D_j)$ is the cardinality of the index set D_j , and the ξ -functions are translation invariant (van Lieshout 2011). If such, the point process \check{X} is called intensity-reweighted moment stationary. Note that, if $m = 2$ it reduces to the so-called second-order intensity-reweighted stationary (Baddeley et al. 2000), where one can obtain $g_2(u, v) - g_1(u) = (\xi_2(u, v) + 1) - 1 = \xi_2(u, v)$, where $g_2(u, v) = \rho(u, v)$ is the so-called pair correlation function and only depends on $d(u, v)$. A more stringent condition, entailing that \check{X} has an identical distribution to $\check{X} + a = \{x + a : x \in \check{X}\}$, $a \in \mathbb{R}^2$, meaning that the spatial distribution of \check{X} is invariant under translations, leads to the concept of stationarity.

Next, by increasing marks' complexity, we present different summary characteristics designed to investigate the interactions between marks subject to an interpoint distance $0 < r < \infty$. Under specific assumptions, such as mark independence, these summary characteristics have predetermined values, making them valuable benchmarks for indicating interactions between marks. Thus, they can be effectively utilized in the hypothesis

testing framework, enabling us to draw inferences about the spatial structure of marks and identify potential deviations from mark independence. Depending on the data in question, null models include, e.g., the superposition of some independent point processes and the random labelling of points generated by subsequent processes.

Throughout the paper, we consider three types of marks which are frequently seen in practice, namely, qualitative marks, quantitative marks, and non-scalar marks. In the case of qualitative marks, points are categorized into distinct types where each type can be labelled by, e.g., an integer value or a letter, see the influenza virus protein data in Fig. 1; point processes with such marks are usually called multitype/multivariate point processes. In the case of quantitative marks, the marks live on a continuous scale, i.e., they are real numbers; see public street trees in Fig. 2. Note that these two cases may be translated into one another based on some rules. For instance, quantitative marks may be translated into qualitative marks based on some data aggregation leading to subsets of marks (Illian et al. 2008; Baddeley et al. 2015). In what follows, we refer to the qualitative marks as *discrete and integer-valued marks* and the quantitative marks as *real-valued marks*. In a more complicated scenario where marks do not belong to either of these categories, marks may have non-scalar forms such as functions, compositions, graphs, and manifolds (Ghorbani et al. 2021; Eckardt et al. 2023). All of these non-scalar marks are summarized into a generic class called *object-valued marks*.

3.1. DISCRETE AND INTEGER-VALUED MARKS

When marks represent a set of countably many entities, which correspond to the simplest case of all potential marked point process settings, each point is assigned a single label $1 \leq k \leq n$, meaning that an integer-valued quantity categorizes points. The underlying process is called a multivariate/multitype point process in this case. More specifically, the observed point pattern \mathbf{x} will be denoted as a collection $\{\mathbf{x}_1, \dots, \mathbf{x}_k\}$, $k \geq 2$, where for each point pattern \mathbf{x}_i , $i = 1, \dots, k$, we have $m(\cdot) = i$, meaning that each point pattern \mathbf{x}_i consists of some points with a specific mark. Within the literature, multivariate/multitype point patterns are commonly analyzed through the so-called cross/dot-type extensions of the classical summary characteristic such as the K -functions and the pair correlation functions ρ (Ripley 1976; Baddeley et al. 2000), the nearest-neighbor distance distribution function H , the empty space function F , and the J -function (van Lieshout and Baddeley 1996; van Lieshout 2011). The cross-type summary characteristics concern pairs of points with distinct marks, whereas the dot-type summary characteristics focus on the relationship between points with a particular mark and all the remaining points. These summary characteristics were initially developed for homogeneous marked spatial point processes (Lotwick and Silverman 1982; Harkness and Isham 1983; van Lieshout and Baddeley 1999). However, since, in practice, homogeneity is rarely satisfied, we here only present the inhomogeneous versions of such summary characteristics, which reduce to their homogeneous counterparts when intensity is constant. We note that the inhomogeneous K -functions and pair correlation functions for multivariate/multitype point processes in \mathbb{R}^2 are defined for second-order intensity-reweighted stationary processes (Møller and Waagepetersen 2003), whereas the inhomogeneous nearest-neighbor distance distribution function H , the inhomogeneous

empty space function F , and the inhomogeneous J -function demand intensity-reweighted moment stationarity (Cronie and van Lieshout 2016).

For a second-order intensity-reweighted stationary multivariate/multitype point process X in \mathbb{R}^2 , the cross-type K -function is given as

$$K_{ij}^{\text{inhom}}(r) = \mathbb{E} \left[\sum_{x \in X_j} \frac{\mathbf{1}\{d(u, x) \leq r\}}{\lambda_j(x)} \middle| u \in X_i \right], \quad i, j = 1, \dots, k, \quad r > 0, \quad (4)$$

where X_i, X_j are point processes with marks i, j , respectively, and $\lambda_j(\cdot)$ is the intensity function of X_j . If X_i does not depend on X_j , then $K_{ij}^{\text{inhom}}(r) = \pi r^2$, which can serve as a criterion to discover independence between sub-processes of different types. If $K_{ij}^{\text{inhom}}(r) > \pi r^2$, one can conclude that the expected number of points of type j around points of type i , within an interpoint distance r , is more than the expected number of such points under mark independence, indicating a tendency to occur around points of types i . On the contrary, $K_{ij}^{\text{inhom}}(r) < \pi r^2$ means that points of type j tend to maintain an interpoint distance from points of type i . Note that $K_{ii}^{\text{inhom}}(\cdot)$ gives the K -function for X_i . Moreover, the dot-type K -function is of the form

$$K_{i\bullet}^{\text{inhom}}(r) = \mathbb{E} \left[\sum_{\substack{x \in X \\ u \neq x}} \frac{\mathbf{1}\{d(u, x) \leq r\}}{\lambda(x)} \middle| u \in X_i \right], \quad i = 1, \dots, k, \quad r > 0, \quad (5)$$

where $\lambda(\cdot)$ is the intensity function of X . In practice, (5) measures the expected number of points of any type around the points with type i . Furthermore, one could obtain the cross/dot-type versions of both the inhomogeneous pair correlation function and the inhomogeneous L -function as $\rho_{ij}^{\text{inhom}}(r) = \partial K_{ij}^{\text{inhom}}(r)/2\pi r$ and $L_{ij}^{\text{inhom}}(r) = \sqrt{K_{ij}^{\text{inhom}}(r)/\pi}$; see Baddeley et al. (2015, Chapter 14) for more details.

Apart from the above cross/dot-type second-order summary characteristics, when X is further intensity-reweighted moment stationary, there exist the cross-type inhomogeneous nearest-neighbor distance distribution function

$$H_{ij}^{\text{inhom}}(r) = 1 - \mathbb{E} \left[\prod_{x \in X_j} \left(1 - \frac{\bar{\lambda}_j}{\lambda_j(x)} \mathbf{1}\{d(u, x) \leq r\} \right) \middle| u \in X_i \right], \quad (6)$$

and the cross-type inhomogeneous J -function

$$J_{ij}^{\text{inhom}}(r) = \frac{1 - H_{ij}^{\text{inhom}}(r)}{1 - F_j^{\text{inhom}}(r)}, \quad F_j^{\text{inhom}}(r) \neq 1, \quad (7)$$

where $\bar{\lambda}_j = \inf_{u \in X_j} \lambda_j(u)$, and

$$F_j^{\text{inhom}}(r) = 1 - \mathbb{E} \left[\prod_{x \in X_j} \left(1 - \frac{\bar{\lambda}_j}{\lambda_j(x)} \mathbf{1}\{d(u, x) \leq r\} \right) \right], \quad u \in \mathbb{R}^2, \quad (8)$$

which is the inhomogeneous empty space function of X_j and does not depend on the choice u (Cronie and van Lieshout 2016). Note that these functions are not symmetric in i and j meaning that, e.g., $H_{ij}^{\text{inhom}}(r) \neq H_{ji}^{\text{inhom}}(r)$, and by setting $j = i$ in (6) and (7) one obtains the nearest-neighbor distance distribution function and the J -function of X_i . Furthermore,

$$H_{i\bullet}^{\text{inhom}}(r) = 1 - \mathbb{E} \left[\prod_{\substack{x \in X \\ u \neq x}} \left(1 - \frac{\bar{\lambda}}{\lambda(x)} \mathbf{1}\{d(u, x) \leq r\} \right) \middle| u \in X_i \right], \quad (9)$$

$$J_{i\bullet}^{\text{inhom}}(r) = \frac{1 - H_{i\bullet}^{\text{inhom}}(r)}{1 - F^{\text{inhom}}(r)}, \quad F^{\text{inhom}}(r) \neq 1, \quad (10)$$

where $F^{\text{inhom}}(r)$ is the inhomogeneous empty space function of X . Due to the properties of J -functions under superposition of stationary and independent processes, van Lieshout and Baddeley (1999) introduced the I -function, which its analogous for inhomogeneous processes becomes $I^{\text{inhom}}(r) = \sum_{i=1}^k p_i J_i^{\text{inhom}}(r) - J^{\text{inhom}}(r)$ where p_i is the probability of type i , and J_i^{inhom} and $J^{\text{inhom}}(r)$ are J -functions of the point process of type i , and the entire process ignoring marks, respectively. Under independence of marks, the I -function becomes zero, whereas deviations from zero suggest either positive or negative associations among the components.

If point processes X_i and X_j are independent, then one expects to have $J_{ij}^{\text{inhom}}(r) = 1$, at least for small distances, while $J_{ij}^{\text{inhom}}(r) > 1$ points to a tendency for point of type j to happen near points of type i , and $J_{ij}^{\text{inhom}}(r) < 1$ means that points of type j prefer not to occur close to points of type i showing an inhibition between the two types (van Lieshout 2011; Cronie and van Lieshout 2016). It is worth mentioning that Cronie and van Lieshout (2016) further extended the Lotwick-Silverman test for random labelling (Lotwick and Silverman 1982) to inhomogeneous settings; see Baddeley et al. (2015, Chapter 14.7) for details on the practical use of cross/dot-type summary characteristics for testing random labelling and mark independence. In general terms, cross-type summary characteristics are employed to examine whether the presence of a specific type can influence the occurrence of points belonging to another type.

3.1.1. Mark Connection and Mingling Functions

Within the context of discrete/integer-valued marks, a variety of more accessible empirical summary characteristics are available for stationary point processes. For instance, the mark connection functions $p_{ij}(r) = p_i p_j \rho_{ij}(r) / \rho(r)$, where p_i is the probability of type i and $\rho(\cdot)$ is the pair correlation function, can be intuitively interpreted as the conditional probability that two points at a distance $r > 0$ from each other possess marks i and j provided that these points belong to the point process X ; under the independence of X_i and X_j , we have $p_{ij}(r) = p_i p_j$ (Stoyan and Stoyan 1994; Illian et al. 2008; Baddeley 2010). For $p_{ij}(r) > p_i p_j$, the mark connection function indicates a tendency of aggregation of components i and j around each other while $p_{ij}(r) < p_i p_j$ suggests repulsion. Moreover,

the normalized mark mingling function is given by

$$v(r) = \frac{1}{c} \mathbb{E} \left[\sum_{x,y \in X}^{\neq} \mathbf{1}\{m(x) \neq m(y)\} \mathbf{1}\{d(x,y) \leq r\} \right], \quad (11)$$

where $c = \sum_{i=1}^k (n_i(n - n_i)) / (n(n - 1))$ is a normalizing constant, n_i is the number of points of type i , and \sum^{\neq} means that points are distinct (Lewandowski and Pommerening 1997; Pommerening et al. 2011; Hui and Pommerening 2014). The mark mingling function $v(r)$ may be used to disclose aggregation of points with (dis)similar marks so that under the mark independence we have $v(r) \approx 1$, while $v(r) > 1$ indicates heterospecific attraction, and $v(r) < 1$ suggests conspecific attraction. Note that the main difference between the mingling function and the cross/dot-type summary characteristics is that the mingling function quantifies the general closeness of points with distinct types.

3.2. REAL-VALUED MARKS

Unlike discrete/integer-valued marks, which are investigated by reformulating a marked point process into $k > 1$ components, real-valued marks are commonly analyzed through a test function $t_f : \mathbb{M} \times \mathbb{M} \rightarrow \mathbb{R}^+$, which is usually a function of the marks of two points located at a distance of $r > 0$ from each other (Penttinen and Stoyan 1989; Schlather 2001). Moreover, in contrast to the presented summary characteristics in Sect. 3.1, which address the aggregation/repulsion between points of different types, test functions t_f analyze numerical differences between the marks of distinct points as a function of distance. For instance, neighboring points may exhibit (dis)similar mark values and specific points might exert dominance over the marks of neighboring points, having large marks while other points in their vicinity possess smaller marks (Illian et al. 2008, Chapter 5).

For stationary point processes, the t_f -correlation function $\kappa_{t_f}(r)$ is given as

$$\kappa_{t_f}(r) = \frac{\mathbb{E} [t_f(m(x), m(y)) \mid x, y \in X]}{c_{t_f}}, \quad d(x, y) = r, \quad (12)$$

where c_{t_f} is a normalizing factor; note that the numerator in (12) is a conditional expectation with respect to the joint distribution of marks. Thus, one can see how marks of the points, located at a distance r from each other, interact in comparison to the average behavior of marks (Penttinen and Stoyan 1989; Baddeley 2010). Indeed, the explicit form of these mark characteristics relies on the choice of test functions, which often account for the spatial variation of marks as well as their pairwise relationships. Within the literature, denoting the average and the variance of all marks as μ_m and σ_m^2 , different test functions and normalizing factors are proposed, which are given in Table 1. We note that Stoyan and Stoyan (1994) proposed the so-called nearest-neighbor indexes, which are similarly constructed through different test functions but only accounting for the marks at a point $x \in X$ and its nearest-neighbor, rather than considering all potential pairs of points within a distance r from x .

The mark variogram $\gamma_{mm}(r)$ (Cressie 1993; Walder and Stoyan 1996; Stoyan and Walder 2000) measures the local half-squared differences among marks of a pair of points while

Table 1. Common test functions for point processes with real-valued marks. For Schlather's I , $\mu_m(r)$ is the mean of the marks for all points located at a distance r from x , and for the mark differentiation function $c_{t_f}^\Delta = 1 - (2/n(n-1)) \sum_{i=1}^n R_i/m(x_{(i)})$ where $m(x_{(i)})$ is the i -th increasingly-ordered mark, $R_1 = 0$ and $R_i = \sum_{j=1}^{i-1} m(x_{(j)})$, for $2 \leq i \leq n$

Name of the function	Symbol	Test function (t_f)	Normalizing factor (c_{t_f})
Mark variogram	$\gamma_{mm}(r)$	$0.5[m(x) - m(y)]^2$	σ_m^2
Stoyan's mark correlation function	$\kappa_{mm}(r)$	$m(x)m(y)$	μ_m^2
r -mark correlation function	$\kappa_{m\bullet}(r)$	$m(x)$	μ_m
r -mark correlation function	$\kappa_{\bullet m}(r)$	$m(y)$	μ_m
Beisbart/Kerscher's $\kappa_{mm}(r)$	$\kappa_{mm}^{\text{Bei}}(r)$	$m(x) + m(y)$	$2\mu_m$
Isham's $\kappa_{mm}(r)$	$\kappa_{mm}^{\text{Ish}}(r)$	$m(x)m(y) - \mu_m^2$	σ_m^2
Stoyan's mark covariance function	$\text{Cov}_{mm}(r)$	$m(x)m(y) - \mu_m^2$	1
Schlather's I	$I_{mm}(r)$	$(m(x) - \mu_m(r))(m(y) - \mu_m(r))$	σ_m^2
Shimanti's I	$I_{mm}^{\text{Shi}}(r)$	$(m(x) - \mu_m)(m(y) - \mu_m)$	σ_m^2
Mark differentiation function	$\Delta_{mm}(r)$	$1 - \frac{\min(m(x), m(y))}{\max(m(x), m(y))}$	$c_{t_f}^\Delta$

Stoyan's mark correlation function $\kappa_{mm}(r)$ (Stoyan and Stoyan 1994) focuses on the mean of the local product of such marks. In other terms, the objective of the mark variogram $\gamma_{mm}(r)$ is to assess the pairwise differences between the marks associated with a pair of points separated by a distance r , which can provide valuable insights into the spatial relationship and variability of marks within a specified distance range. Concerning Stoyan's mark correlation function $\kappa_{mm}(r)$, the pairwise product is expected to coincide with the global squared mean of the marks μ_m^2 under the independence assumption such that $\kappa_{mm}(r) = 1$. If nearby points have smaller (larger) marks, their average product will also be small (large) and deviate from 1. We note that Stoyan's mark covariance function $\text{Cov}_{mm}(r)$ (Stoyan 1984) is indeed a linear transformation of his mark correlation function such that both $\text{Cov}_{mm}(r)$ and $\kappa_{mm}(r)$ essentially convey the same message (Schlather 2001). Regarding the use of **r**-mark correlation functions, $\kappa_{m\bullet}(r)$ and $\kappa_{\bullet m}(r)$, both reflect the average of the mark for either of the two considered points with respect to a distance r . Except for mark independence, the averaged **r**-mark correlation will not be equal to the overall mark mean since either the mark of the first or second point is chosen. Shedding some light on Beisbart's and Kerscher's version of $\kappa_{mm}(r)$, to have a large contribution in (12), only one of the points is required to have a large mark. However, under mark independence, this mark correlation function coincides with twice the mark mean yielding $\kappa_{mm}^{\text{Bei}}(r) = 1$. In contrast to $\kappa_{mm}(r)$ and $\kappa_{mm}^{\text{Bei}}(r)$, Isham's function $\kappa_{mm}^{\text{Ish}}(r)$ reveals a Pearson-type correlation for marks. Schlather's function $I_{mm}(r)$ gives insight into how marks, for a pair of points with an interpoint distance r , are related to each other (Schlather et al. 2004). Basically, this function centers each mark by the conditional mean mark $\mu_m(r)$, i.e., the **r**-mark correlation function, and normalizes the product by the mark variance σ_m^2 . Shimanti's function $I_{mm}^{\text{Shi}}(r)$ is constructed similarly but uses the global mark mean instead of the conditional mark mean (Shimatani 2002). Finally, the mark differentiation function $\Delta_{mm}(r)$ shows how the ratio of marks varies with respect to a distance r . In the case of mark independence, the minimum and the maximum of marks

are assumed to be, on average, similar such that the proportion becomes 1, i.e., $\Delta_{mm}(r) = 0$ (Pommerening et al. 2011; Hui and Pommerening 2014).

Aiming at defining a K -function for stationary point processes with real-valued marks and a constant intensity λ , which accounts for the correlation between points as well as between marks, Penttinen et al. (1992) proposed a mark-weighted K -function as

$$K_{t_f}(r) = \frac{1}{\lambda c_{t_f}} \mathbb{E} \left[\sum_{\substack{x \in X \\ u \neq x}} t_f(m(u), m(x)) \mathbf{1}\{d(u, x) \leq r\} \middle| u \in X \right], \quad (13)$$

where c_{t_f} is the expected value of $t_f(\cdot)$ under mark independence for which $K_{t_f}(r)$ reduces to the so-called Ripley's K -function (Baddeley et al. 2015, Chapter 15); (13) has already been extended to inhomogeneous cases (Ghorbani et al. 2021). Penttinen et al. (1992) initially considered the Stoyan's mark correlation function in (13), i.e., $t_f(\cdot) = \kappa_{mm}(\cdot)$, and commonly denoted $K_{t_f}(\cdot)$ as $K_{mm}(\cdot)$, however, one may use any of the test functions presented in Table 1. Note that, in the case of $K_{mm}(\cdot)$, by taking the pairwise product of marks as weight into the original (not-mark-weighted) K -function, the estimated curves become scaled versions of the original K -function, except under mark independence. More specifically, for pairwise positively correlated marks, i.e., if the pairwise product of marks exceeds the overall expected value, the empirical curves are up-scaled versions of K -function. Otherwise, they are down-scaled. To better understand the impact of marks, it is advisable to compare the original K -functions with their mark-weighted counterparts. Further, normalizing $K_{t_f}(r)$ by its unmarked counterpart, i.e., the Ripley's K -function, one can obtain a cumulative mark correlation function corresponding to the mean value of the employed test function for a pair of points with an interpoint distance r (Wiegand and Moloney 2013). We note that D'Angelo et al. (2023) recently discussed local mark-weighted cumulative summary statistics, which account for the contribution of each marked point to the global second-order summary characteristics.

A further summary statistics for a stationary point process X , with an intensity function λ and pair correlation $\rho(\cdot)$, is

$$U(r) = \lambda^2 \rho(r) \kappa_{t_f}(r) |dx| |dy|, \quad (14)$$

where $|dx|, |dy|$ are sizes of two infinitesimal small areas around x and y separated by a distance r . In a similar manner as $K_{t_f}(r)$ in (13), $U(\cdot)$ considers the correlation between points and between marks. Since under complete spatial randomness $\rho(\cdot) = 1$, and for independent marks $\kappa_{t_f}(\cdot) = 1$, thus, in the combination of those cases, $U(\cdot)$ reduces to the second-order product density of X (Capobianco and Renshaw 1998; Renshaw 2002). If at least one of these two does not vanish, then $U(\cdot)$ is a version of the second-order product density of X weighted by the spatial interaction between the points via $\rho(\cdot)$ and/or association between the marks via $\kappa_{t_f}(\cdot)$. We note that further summary characteristics, considered within the literature, include the (non)cumulative density correlation functions (Fedriani et al. 2015).

Turning to the case where each data point has two distinct marks $m_i(\cdot)$, $i = 1, 2$, by considering a bivariate test function $\mathfrak{t}_f(m_1(x), m_2(y)) = m_1(x)m_2(y)$, $x, y \in X$, [Stoyan \(1987\)](#) proposed a bivariate mark correlation function with a normalizing constant $\mu_1\mu_2$ where μ_i , $i = 1, 2$, is the mean of mark $m_i(\cdot)$, $i = 1, 2$; see also ([Raventós et al. 2011](#); [Wiegand and Moloney 2013](#)). Note that such an idea can be used to construct other bivariate and multivariate (nearest-neighbor-based) test functions ([Pommerening and Grabarnik 2019](#); [Eckardt et al. 2023](#)). In addition, the cross/dot-type pair correlation function and the bivariate/multivariate mark summary characteristics can also be used to derive cross/dot-type and multivariate versions of $U(r)$ ([Eckardt and Mateu 2019a](#); [Eckardt et al. 2021](#)). Such extended versions not only help to investigate the interrelations between different types of points but also allow to understand the association between different real-valued quantities over space.

3.3. FREQUENCY DOMAIN APPROACHES

Next, we discuss characterizations of marked spatial point processes through frequency domain methods, all of which could be translated into their corresponding distance-based characteristics, e.g., $U(\cdot)$, given in (14), by using the inverse Fourier theorem. This close relationship of the frequency and distance-based approaches allows to derive planar partial mark characteristics which reflect the interrelation between two components X_i, X_j , $i, j = 1, \dots, k$, conditional on all the remaining types.

3.3.1. Frequency Domain Approaches for Discrete and Integer-Valued Marks

In contrast to previous marked summary characteristics, frequency domain characteristics for marked spatial point processes have received limited attention. Bartlett's complete covariance density function ([Bartlett 1964](#)) is given as $\mathfrak{K}_{ij}(u, v) = \lambda_i(u)\delta_{ij}(u - v) + \vartheta_{ij}(u, v)$, $i, j = 1, \dots, k$, where $\vartheta_{ij}(\cdot)$ is a covariance density function ([Muggestone and Renshaw 1996](#), Equation 2) and $\delta_{ij}(\cdot)$ is a two-dimensional Dirac delta-function. Similar to the covariance density function, $\mathfrak{K}_{ij}(u, v)$ may be used to describe the cross-second-order behavior of points within the corresponding space. However, unlike the covariance function, it controls for multiple coincident points via δ_{ij} . Under stationarity, for the components X_i, X_j , $i, j = 1, \dots, k$, and at frequencies $\boldsymbol{\omega} = (\omega_1, \omega_2)$, [Muggestone and Renshaw \(1996\)](#), using a discrete Fourier transform representation, introduced the cross-spectral density function $f_{ij}(\boldsymbol{\omega})$ as

$$f_{ij}(\boldsymbol{\omega}) = \int_{-\infty}^{\infty} \mathfrak{K}_{ij}(r) \exp(-i\boldsymbol{\omega}^\top r) dr, \quad \mathfrak{K}_{ij}(r) = \lambda_i \delta_{ij}(r) + \vartheta_{ij}(r),$$

where $r = d(u, v)$, $u, v \in \mathbb{R}^2$, $\boldsymbol{\omega}^\top$ is the transpose of $\boldsymbol{\omega}$, and $i = \sqrt{-1}$. Note that, in this case, $\mathfrak{K}_{ij}(\cdot)$ only depends on distances between points. Two components are said to be independent if the cross-spectral density function equals zero for all frequencies. We note that by applying Theorem 8.3.1 of [Brillinger \(1981\)](#), the above spectrum could be translated into a partial version $f_{ij|X \setminus \{i, j\}}(\boldsymbol{\omega})$, i.e., computing the Schur complement of the cross-spectral densities ([Eckardt and Mateu 2019a](#)).

The cross-spectral function $f_{ij}(\cdot)$ further leads to the definition of some interesting spectral functions, including the spectral coherence function $R_{ij}^2(\boldsymbol{\omega}) = f_{ij}^2(\boldsymbol{\omega}) / (f_{ii}(\boldsymbol{\omega})f_{jj}(\boldsymbol{\omega}))$, the cross-amplitude spectrum $\zeta_{ij}(\boldsymbol{\omega}) = \text{mod}\{f_{ij}(\boldsymbol{\omega})\}$, the cross-phase spectrum $\wp_{ij}(\boldsymbol{\omega}) = \arg(f_{ij}(\boldsymbol{\omega}))$, the gain function $\mathfrak{G}_{i|j}(\boldsymbol{\omega}) = \sqrt{(f_{ij}(\boldsymbol{\omega})R_{ij}(\boldsymbol{\omega}))/f_i(\boldsymbol{\omega})}$, and also their corresponding partial versions (Mugglestone and Renshaw 1996; Eckardt and Mateu 2019b). To offer some insight into these functions, the cross-phase spectrum $\wp(\boldsymbol{\omega})$ measures the similarity between two point patterns to see if the spectrum of one pattern is a linear shift of that of the other one, while the cross-amplitude spectrum $\zeta_{ij}(\boldsymbol{\omega})$ encodes the relative magnitude of frequencies for two point patterns. Thus, both functions are useful tools to investigate the characteristics of the empirical spectrum with respect to the frequencies $\boldsymbol{\omega}$. The gain spectrum $\mathfrak{G}_{i|j}(\boldsymbol{\omega})$, in contrast, can be interpreted as a regression coefficient in a linear regression at frequency $\boldsymbol{\omega}$, for two corresponding components. Additionally, one could make directional inferences by transforming the spectral density functions into the polar form; see e.g., Renshaw and Ford (1983, 1984).

3.3.2. Frequency Domain Approaches for Real-Valued Marks

The accessibility of frequency domain approaches for real-valued marks is more constrained than that of discrete marks. The auto/cross-spectral density functions for real-valued marks, at frequencies $\boldsymbol{\omega} = (\omega_1, \omega_2)$, can be obtained from a discrete Fourier transformation of the corresponding $U(r)$ functions, as

$$f_{ii}^m(\boldsymbol{\omega}) = \int U_{ii}(r) \exp(-i\boldsymbol{\omega}^\top r) dr, \quad f_{ij}^m(\boldsymbol{\omega}) = \int U_{ij}(r) \exp(-i\boldsymbol{\omega}^\top r) dr,$$

where $U_{ii}(r)$ and $U_{ij}(r)$ are the auto/cross-type versions of (14), respectively; see Renshaw (2002); Eckardt and Mateu (2019a) for more details. Analogous to the multivariate spatial point processes, the marked spectra can be transformed into partial versions using the results of Theorem 8.3.1 of Brillinger (1981).

As of some recent development, Eckardt and Mateu (2019a), using a re-scaled version of the partial spectral coherence, defined a spatial dependence graph model for multivariate point processes with, potentially, real-valued marks, according to which different components are represented as nodes and the conditional independence structure among the k components is reflected by missing edges.

3.4. OBJECT-VALUED MARKS

We now discuss mark summary characteristics for an observed point pattern $\mathbf{x} = \{x_i, \sigma(x_i)\}_{i=1}^n$ in which each point x_i is augmented by a non-scalar quantity $\sigma(x_i)$. Apart from the function-valued mark settings (Comas et al. 2008, 2011, 2013; Ghorbani et al. 2021; Eckardt et al. 2023), this newly introduced class of marked spatial point processes further includes the cases where marks are constrained arrays or inherently structured quantities (Eckardt et al. 2023).

Focusing explicitly on function-valued mark scenarios, where for each point x_i the corresponding mark $\sigma(x_i)$ is a function-valued quantity $g(x_i)$ on $\mathbb{F}(\mathcal{T})$ with $\mathcal{T} = (a, b)$, $-\infty \leq$

Table 2. Pointwise test functions for point processes with function-valued marks

Name of the function	Symbol	Test function (t_f)	Normalizing factor (c_{t_f})
Mark variogram	$\gamma_{gg}(r, t)$	$0.5[g_x(t) - g_y(t)]^2$	$\sigma_g^2(t)$
Stoyan's mark correlation function	$\kappa_{gg}(r, t)$	$g_x(t)g_y(t)$	$\mu_g^2(t)$
r -mark correlation function	$\kappa_{g\bullet}(r, t)$	$g_x(t)$	$\mu_g(t)$
r -mark correlation function	$\kappa_{\bullet g}(r, t)$	$g_y(t)$	$\mu_g(t)$
Beisbart and Kerscher's $\kappa_{gg}(r, t)$	$\kappa_{gg}^{\text{B\&K}}(r, t)$	$g_x(t) + g_y(t)$	$2\mu_g(t)$
Isham's $\kappa_{gg}(r, t)$	$\kappa_{gg}^{\text{Ish}}(r, t)$	$g_x(t)g_y(t) - \mu_g^2(t)$	$\sigma_g^2(t)$
Stoyan's mark covariance function	$\mathbb{C}\text{ov}_{gg}(r, t)$	$g_x(t)g_y(t) - \mu_g^2(t)$	1
Schlather's I	$I_{gg}(r, t)$	$(g_x(t) - \mu_g(r, t))(g_y(t) - \mu_g(r, t))$	$\sigma_g^2(t)$
Shimanti's I	$I_{gg}^{\text{Shi}}(r, t)$	$(g_x(t) - \mu_g(t))(g_y(t) - \mu_g(t))$	$\sigma_g^2(t)$

$a \leq b \leq \infty$ and $g(x_i) : \mathcal{T} \subseteq \mathbb{R} \mapsto \mathbb{R}$, different mark summary characteristics, similarly to Sect. 3.2, can be defined through an extended test function $t_f : \mathbb{F} \times \mathbb{F} \rightarrow \mathbb{R}^+$. Starting with a pointwise specification, Eckardt et al. (2023) proposed a generalized version of the t_f -correlation function in (12), given as

$$\kappa_{t_f}(r, t) = \frac{\mathbb{E} [t_f(g_x(t), g_y(t)) | x, y \in X]}{c_{t_f}(t)}, \quad d(x, y) = r, \quad (15)$$

where $g_x(t)$ is the mark of x at $t \in \mathcal{T}$, and $c_{t_f}(t)$ is, for a fixed $t \in \mathcal{T}$, a pointwise normalizing constant corresponding to the expectation of the test function when r tends to infinity. Denoting the mean and variance of all marks at $t \in \mathcal{T}$ by $\mu_g(t)$ and $\sigma_g^2(t)$, a summary of the extended test functions is presented in Table 2. As an example, normalizing $\mathbb{E} [0.5 (g_x(t) - g_y(t))^2 | x, y \in X]$ by $\sigma_g^2(t)$ gives the pointwise mark variogram $\gamma_{gg}(r, t)$, and, to obtain a pointwise version of Stoyan's mark correlation function, the expectation $\mathbb{E} [g_x(t)g_y(t) | x, y \in X]$ needs to be divided by $\mu_g^2(t)$ (Eckardt et al. 2023).

While all the test functions presented in Table 2, with respect to the argument t , have similar interpretations as those in Table 1, they do not convey any information on the overall pairwise interrelation between the function-valued quantities under study. The desired global mark characteristics, however, can be constructed from their pointwise versions by the integration of the normalized expectation of $t_f(g_x(t), g_y(t))$ over \mathcal{T} . In this respect, e.g., the pointwise mark variogram and mark correlation functions translate into global versions

$$\begin{aligned} \gamma_{gg}(r) &= \int_{\mathcal{T}} \mathbb{E} [0.5 (g_x(t) - g_y(t))^2 | x, y \in X] dt =: \int_{\mathcal{T}} \gamma_{gg}(r, t) dt, \\ \kappa_{gg}(r) &= \int_{\mathcal{T}} \mathbb{E} [g_x(t)g_y(t) | x, y \in X] dt =: \int_{\mathcal{T}} \kappa_{gg}(r, t) dt. \end{aligned}$$

Then, any such global characteristic allows, in a \mathcal{L}_2 sense, for a similar interpretation as discussed for Table 1. In other words, the global mark variogram $\gamma_{gg}(r)$ gives the average variability over distinct pairs of functions for two points with an interpoint distance r . Under mark independence, the average differences among the curves are expected to coincide with

the functional variance such that $\gamma_{gg}(r) = 1$. Likewise, $\kappa_{gg}(r)$ assesses the product of any two function-valued quantities with respect to the interpoint distance of points $x, y \in X$, where $\kappa_{gg}(r) = 1$ for independent marks. Unlike $\kappa_{gg}(r)$, both \mathbf{r} -mark correlation functions $\kappa_{g\bullet}(r)$ and $\kappa_{\bullet g}(r)$ concern the average of the function-valued point attribute of either the first or the second point subject to a distance r . This usually differs from μ_g except under mark independence settings. Similarly to Beisbart and Kerscher's original version, $\kappa_{gg}^{\text{Bei}}(r)$ reflects the average pairwise sum, i.e., the perturbation, of any two functions for a pair of points at a distance r from each other, which is expected to coincide with twice the functional mean μ_g if marks are independent. If two functions of two nearby locations vary strongly from the functional average, the sum of both would also be, on average, different from the expected case, leading to $\kappa_{gg}^{\text{Bei}}(r) \neq 1$. Lastly, in both function-valued versions of Schlather's and Shimanti's I -functions, marks are centered by the conditional and unconditional functional means squared, respectively, and scaled by the functional variance. We note that the same ideas can also be applied to the mark-weighted K -function $K_{t_f}(\cdot)$, given in (13), and the function $U(r)$, given in (14), leading to $K_{t_f}(r) = \int K_{t_f}(r, t)dt$ and $U(r) = \int U(r, t)dt$.

The extensions of the above summary characteristics to multivariate spatial point processes, including *cross-type*, *cross-function* and *cross-type cross-function* mark summary characteristics are discussed by Eckardt et al. (2023). While the cross-type characteristics reveal the interrelation of mark $g(\cdot)$ for points $x \in X_i$ and $y \in X_j$, the cross-function versions are defined for function-valued marks $g_1(\cdot) \neq g_2(\cdot)$ on \mathbb{F}^2 at locations $x, y \in X$. These ideas are also extended to *multi-function* versions for the sets $\{x_i, \mathbf{g}(x_i)\}_{i=1}^n$, with $\mathbf{g}(x_i) \in \mathbb{F}^s$, $s > 2$, see (Eckardt et al. 2023) for details.

4. MARKED SPATIAL POINT PROCESSES ON LINEAR NETWORKS

In certain point process applications, the spatial distribution of events may become confined within an underlying structure such as a linear network. Instances of relevant applications include the study of street crimes, traffic accidents, trees positioned alongside roads, and various other scenarios where spatial locations are restricted to a linear network.

A linear network $\mathbb{L} \subset \mathbb{R}^2$ is considered as a finite union of some line segments, i.e., $\mathbb{L} = \bigcup_{i=1}^k l_i$ where $l_i = [u_i, v_i] = \{tu_i + (1-t)v_i : 0 \leq t \leq 1\}$, $u_i \neq v_i \in \mathbb{R}^2$. The linear network \mathbb{L} is equipped with a regular distance metric $d_{\mathbb{L}}$, with the shortest-path distance being an example of it (Rakshit et al. 2017). Moreover, no over/under-pass is assumed within \mathbb{L} , meaning that each intersection of segments is a node. The total length of \mathbb{L} is denoted by $|\mathbb{L}| = \sum_{i=1}^k |l_i|$ where $|l_i| = d_{\mathbb{L}}(u_i, v_i) = d(u_i, v_i)$. We denote a marked point process on \mathbb{L} by $X^{\mathbb{L}}$ for which an observed point pattern is denoted by $\mathbf{x}^{\mathbb{L}}$, and similarly to (1), if we let $\check{X}^{\mathbb{L}}$ be the unmarked version of $X^{\mathbb{L}}$, then

$$\mathbb{E}[N(\check{X}^{\mathbb{L}} \cap A)] = \int_A \lambda^{\mathbb{L}}(u) d_1 u, \quad A \subset \mathbb{L}, \quad (16)$$

where $\lambda^{\mathbb{L}}(\cdot)$ is the intensity function of $\check{X}^{\mathbb{L}}$, and d_1 stands for integration with respect to arc length on the network. In this case, the intensity function $\lambda^{\mathbb{L}}(u)$ gives the expected number

of points per unit length of the network in the vicinity of a location $u \in \mathbb{L}$ (Baddeley et al. 2021). Similarly to (2) and (3), for any non-negative measurable function $f(\cdot)$ on the product space \mathbb{L}^m ,

$$\mathbb{E} \left[\sum_{x_1, \dots, x_m \in \check{X}^{\mathbb{L}}}^{\neq} f(x_1, \dots, x_m) \right] = \int_{\mathbb{L}^m} f(u_1, \dots, u_m) \lambda_{\mathbb{L}}^{(m)}(u_1, \dots, u_m) d_1 u_1 \cdots d_1 u_m, \quad (17)$$

and

$$g_m^{\mathbb{L}}(u_1, \dots, u_m) = \frac{\lambda_{\mathbb{L}}^{(m)}(u_1, \dots, u_m)}{\lambda_{\mathbb{L}}^{\mathbb{L}}(u_1) \cdots \lambda_{\mathbb{L}}^{\mathbb{L}}(u_m)} = \sum_{j=1}^m \sum_{D_1, \dots, D_j} \xi_{N(D_1)}^{\mathbb{L}}(\{u_i : i \in D_1\}) \cdots \xi_{N(D_j)}^{\mathbb{L}}(\{u_i : i \in D_j\}), \quad (18)$$

where $\lambda_{\mathbb{L}}^{(m)}(\cdot)$ is the m -order product intensity of $X^{\mathbb{L}}$, \sum_{D_1, \dots, D_j} ranges over all partitions $\{D_1, \dots, D_j\}$ of $\{1, \dots, m\}$ into j non-empty and disjoint sets, and $N(D_j)$ is the cardinality of the index set D_j (Cronie et al. 2020). In order to propose marked summary characteristics for point processes on linear networks, similarly to Sect. 3, we need some form of stationarity. However, addressing this challenge has proven to be quite complex (Baddeley et al. 2017), as, currently, there is no transformation that can transform points on a linear network with the guarantee that the shifted point will remain on the same network. Cronie et al. (2020) proposed different notions of stationarity on linear networks, according to which whenever the product intensities $\lambda_{\mathbb{L}}^{(m)}(\cdot)$, $1 \leq m \leq k$, exist, $\bar{\lambda}_{\mathbb{L}} = \inf_{u \in \mathbb{L}} \lambda_{\mathbb{L}}(u) > 0$, and for any $m \in \{2, \dots, k\}$ the correlation function $g_m^{\mathbb{L}}(\cdot)$ satisfies

$$g_m^{\mathbb{L}}(u_1, \dots, u_m) = \bar{g}_m^{\mathbb{L}}(d_{\mathbb{L}}(u, u_1), \dots, d_{\mathbb{L}}(u, u_m)), \quad (19)$$

for any $u \in \mathbb{L}$ and some function $\bar{g}_m^{\mathbb{L}} : [0, \infty)^m \rightarrow [0, \infty)$, they said that $\check{X}^{\mathbb{L}}$ is k -th order intensity-reweighted pseudostationary (with respect to a regular distance $d_{\mathbb{L}}$); $\check{X}^{\mathbb{L}}$ is called intensity reweighted moment pseudostationary (IRMPS) when this holds for any order $k \geq 2$. Moreover, a homogeneous point process $\check{X}^{\mathbb{L}}$, which is also k -th order intensity-reweighted pseudostationary, is called k -th order pseudostationary; if this holds for any order $k \geq 2$, then $\check{X}^{\mathbb{L}}$ is moment pseudostationary. Finally, a moment pseudostationary point process $\check{X}^{\mathbb{L}}$ is considered (strongly) pseudostationary if its moments completely and uniquely characterize its distribution.

4.1. DISCRETE AND INTEGER-VALUED MARKS

We now consider the case where $X^{\mathbb{L}}$ can be decomposed into $1 < k \leq n$ components according to a discrete/integer-valued mark. In this case, $X^{\mathbb{L}}$ is called a multivariate/multitype spatial point process on a linear network \mathbb{L} , and $\mathbf{x}^{\mathbb{L}}$ is denoted as a collection of k distinct point patterns $\{\mathbf{x}_1^{\mathbb{L}}, \dots, \mathbf{x}_k^{\mathbb{L}}\}$, $k \geq 2$. Marked summary characteristics for multivariate/multitype point processes on linear networks can be naturally proposed by mimicking

the underlying principles presented in Sect. 3.1. However, despite the growing availability of multivariate/multitype spatial network-constrained point processes, there exist only a few of such extensions (Spooner et al. 2004; Baddeley et al. 2014; Eckardt and Mateu 2018, 2021). In particular, most attention is paid to the pair correlation function and the K -function, for which, by following (Ang et al. 2012; Baddeley et al. 2014; Rakshit et al. 2017), for a second-order pseudostationary ($d_{\mathbb{L}}$ -correlated in the language of Rakshit et al. (2017)) marked point process $X^{\mathbb{L}}$, we have

$$K_{ij}^{\mathbb{L},\text{inhom}}(u, r_{\mathbb{L}}) = \mathbb{E} \left[\sum_{x \in X_j^{\mathbb{L}}} \frac{\mathbf{1}\{d_{\mathbb{L}}(u, x) \leq r_{\mathbb{L}}\} \nabla\{u, d_{\mathbb{L}}(u, x)\}}{\lambda_j^{\mathbb{L}}(x)} \middle| u \in X_i^{\mathbb{L}} \right],$$

$$i, j = 1, \dots, k, \quad r_{\mathbb{L}} > 0, \quad (20)$$

and

$$K_{i\bullet}^{\mathbb{L},\text{inhom}}(u, r_{\mathbb{L}}) = \mathbb{E} \left[\sum_{\substack{x \in X^{\mathbb{L}} \\ u \neq x}} \frac{\mathbf{1}\{d_{\mathbb{L}}(u, x) \leq r_{\mathbb{L}}\} \nabla\{u, d_{\mathbb{L}}(u, x)\}}{\lambda^{\mathbb{L}}(x)} \middle| u \in X_i^{\mathbb{L}} \right],$$

$$i = 1, \dots, k, \quad r_{\mathbb{L}} > 0, \quad (21)$$

where $\nabla(\cdot)$ acts as an edge correction factor, and $\lambda_j^{\mathbb{L}}(\cdot)$ is the intensity function of the j -th component. If the employed metric is the shortest-path distance, then $\nabla\{u, d_{\mathbb{L}}(u, x)\}$ becomes the reciprocal of the number of points lying exactly $d_{\mathbb{L}}(u, x)$ units from $u \in X_i$. As for some interpretation, if $X_i^{\mathbb{L}}$ does not depend on $X_j^{\mathbb{L}}$, then $K_{ij}^{\mathbb{L},\text{inhom}}(u, r_{\mathbb{L}}) = r_{\mathbb{L}}$, which serves as a criterion to discover independence between processes of different types. If $K_{ij}^{\mathbb{L},\text{inhom}}(u, r_{\mathbb{L}}) > r_{\mathbb{L}}$, one can conclude that the expected number of points of type j around points of type i , within an interpoint distance $r_{\mathbb{L}}$, is more than the expected number of such points under mark independence, indicating a tendency to occur around points of types i . On the contrary, $K_{ij}^{\mathbb{L},\text{inhom}}(u, r_{\mathbb{L}}) < r_{\mathbb{L}}$ means that points of type j tend to maintain an interpoint distance from points of type j . In a similar manner, one could obtain the cross/dot-type versions of the inhomogeneous pair correlation function.

4.1.1. Mark Connection and Mingling Function

For homogeneous point processes on linear networks, Baddeley et al. (2014) presented the network-based versions of the mark connection and mark equality functions as

$$p_{ij}^{\mathbb{L}}(r_{\mathbb{L}}) = \frac{\lambda_i^{\mathbb{L}} \lambda_j^{\mathbb{L}} \rho_{ij}^{\mathbb{L}}(r_{\mathbb{L}})}{[\lambda^{\mathbb{L}}]^2 \rho^{\mathbb{L}}(r_{\mathbb{L}})}, \quad p^{\mathbb{L}}(r_{\mathbb{L}}) = \sum_{i=1}^k p_{ii}^{\mathbb{L}}(r_{\mathbb{L}}), \quad (22)$$

where $\rho_{ij}^{\mathbb{L}}(\cdot)$ is the cross-pair correlation function for the processes with types i, j and $\rho^{\mathbb{L}}(\cdot)$ is the pair correlation of the full process. Given the presence of a pair of points separated by a distance $r_{\mathbb{L}}$, the mark connection function $p_{ij}^{\mathbb{L}}(r_{\mathbb{L}})$ is intuitively the conditional probability

that those points are of types i, j . Similarly, $p^{\mathbb{L}}(r_{\mathbb{L}})$ is the conditional probability that those two points, separated by a distance $r_{\mathbb{L}}$, have the same type.

Next, we propose the counterpart of the normalized mingling function (11) for homogeneous point processes on linear networks as

$$\nu_{\mathbb{L}}(r_{\mathbb{L}}) = \frac{1}{c} \mathbb{E} \left[\sum_{x, y \in X^{\mathbb{L}}}^{\neq} \mathbf{1}\{m(x) \neq m(y)\} \mathbf{1}\{d_{\mathbb{L}}(x, y) \leq r_{\mathbb{L}}\} \right], \quad (23)$$

where c is the same normalizing factor as in (11). Similarly, $\nu_{\mathbb{L}}(\cdot)$ may be used to discover aggregation/repulsion between points of different types subject to an interpoint distance $r_{\mathbb{L}}$. Note that, unlike the cross/dot-type summary characteristics, the mingling functions (11) and (23) quantify the aggregation and/or repulsion tendencies between any given types, rather than focusing on the interaction between two specific types or comparing a single type against all other types.

4.1.2. Higher-Order Marked Inhomogeneous Summary Characteristics

Next, for IRMPS point processes, we extend the inhomogeneous higher-order summary characteristics of Cronie et al. (2020) to multivariate/multitype settings. In particular, we have the following cross-type summary characteristics

$$H_{ij}^{\mathbb{L}, \text{inhom}}(u, r_{\mathbb{L}}) = 1 - \mathbb{E} \left[\prod_{x \in X_j^{\mathbb{L}}} \left(1 - \frac{\bar{\lambda}_j^{\mathbb{L}} \mathbf{1}\{d_{\mathbb{L}}(u, x) \leq r\}}{\lambda_j^{\mathbb{L}}(x)} \nabla\{u, d_{\mathbb{L}}(u, x)\} \right) \middle| u \in X_i^{\mathbb{L}} \right], \quad (24)$$

$$J_{ij}^{\mathbb{L}, \text{inhom}}(u, r_{\mathbb{L}}) = \frac{1 - H_{ij}^{\mathbb{L}, \text{inhom}}(u, r_{\mathbb{L}})}{1 - F_j^{\mathbb{L}, \text{inhom}}(u, r_{\mathbb{L}})}, \quad F_j^{\mathbb{L}, \text{inhom}}(u, r_{\mathbb{L}}) \neq 1, \quad (25)$$

$$F_j^{\mathbb{L}, \text{inhom}}(u, r_{\mathbb{L}}) = 1 - \mathbb{E} \left[\prod_{x \in X_j^{\mathbb{L}}} \left(1 - \frac{\bar{\lambda}_j^{\mathbb{L}} \mathbf{1}\{d_{\mathbb{L}}(u, x) \leq r\}}{\lambda_j^{\mathbb{L}}(x)} \nabla\{u, d_{\mathbb{L}}(u, x)\} \right) \right], \quad (26)$$

and the dot-type ones

$$H_{i\bullet}^{\mathbb{L}, \text{inhom}}(u, r_{\mathbb{L}}) = 1 - \mathbb{E} \left[\prod_{\substack{x \in X^{\mathbb{L}} \\ u \neq x}} \left(1 - \frac{\bar{\lambda}_j^{\mathbb{L}} \mathbf{1}\{d_{\mathbb{L}}(u, x) \leq r\}}{\lambda_j^{\mathbb{L}}(x)} \nabla\{u, d_{\mathbb{L}}(u, x)\} \right) \middle| u \in X_i^{\mathbb{L}} \right], \quad (27)$$

$$J_{i\bullet}^{\mathbb{L}, \text{inhom}}(u, r_{\mathbb{L}}) = \frac{1 - H_{i\bullet}^{\mathbb{L}, \text{inhom}}(u, r_{\mathbb{L}})}{1 - F_{i\bullet}^{\mathbb{L}, \text{inhom}}(u, r_{\mathbb{L}})}, \quad F_{i\bullet}^{\mathbb{L}, \text{inhom}}(u, r_{\mathbb{L}}) \neq 1, \quad (28)$$

where $F_j^{\mathbb{L}, \text{inhom}}(u, r_{\mathbb{L}})$ and $F_j^{\mathbb{L}, \text{inhom}}(u, r_{\mathbb{L}})$ are the inhomogeneous empty space functions of $X^{\mathbb{L}}$ and $X_j^{\mathbb{L}}$, respectively. For any two fixed types i, j , $J_{ij}^{\mathbb{L}, \text{inhom}}(\cdot) = 1$ means points

of type j are randomly distributed around points of type i , $J_{ij}^{\mathbb{L},\text{inhom}}(\cdot) > 1$ means that points of type j tend to happen around points of type i , and $J_{ij}^{\mathbb{L},\text{inhom}}(\cdot) < 1$ indicates that points of type j prefer to maintain a distance from points of type i . Turning to the dot-type J -function, if $J^{\mathbb{L},\text{inhom}}_{i\bullet}(\cdot) = 1$, it implies that points of different types than i exhibit no interaction with points of type i . When $J^{\mathbb{L},\text{inhom}}_{i\bullet}(\cdot) > 1$, it signifies a propensity for other types to appear in proximity to points of type i . Conversely, $J^{\mathbb{L},\text{inhom}}_{i\bullet}(\cdot) < 1$ suggests a preference for other types to hold a distance from points of type i . Note that these higher-order marked summary characteristics are expected to provide a deeper understanding of the spatial patterns and clustering/repulsion behavior among points of different types since they go beyond pairwise interactions, providing information about interactions involving more than two points.

4.2. REAL-VALUED MARKS

We now turn to a situation where points are labelled by real-valued marks. By taking the presented methods in Sect. 3.2 into account, we present novel mark summary characteristics that account for the geometry of the underlying network.

4.2.1. Mark Correlation Functions and Mark-Weighted Summary Characteristics

Generalizing the t_f -correlation functions in (12) to the present setting, the $t_f^{\mathbb{L}}$ -correlation function for point processes on linear networks is of the form

$$\kappa_{t_f^{\mathbb{L}}}(r_{\mathbb{L}}) = \frac{\mathbb{E} \left[t_f^{\mathbb{L}}(m(x), m(y)) \mid x, y \in X^{\mathbb{L}} \right]}{c_{t_f^{\mathbb{L}}}}, \quad d_{\mathbb{L}}(x, y) = r_{\mathbb{L}}, \quad (29)$$

where $c_{t_f^{\mathbb{L}}}$ is a normalizing factor; the superscript \mathbb{L} emphasizes that the t_f -correlation function will be evaluated over the linear network \mathbb{L} . The numerator in (29) is the conditional expectation of a given test function evaluated over a pair of marks whose corresponding spatial locations are $r_{\mathbb{L}}$ distances away, given that the two points belong to the point process $X^{\mathbb{L}}$. The denominator $c_{t_f^{\mathbb{L}}}$ is the expected value of the considered test function under the mark independence assumption. Applying the same principles as in Sect. 3.2, the specific linear-network-based mark characteristic depends on the explicit formulation of the test function. A summary of potential test functions for point processes on linear networks and their notations is given in Table 3.

Taking the geometry of linear networks into account and employing a regular metric (Rakshit et al. 2017), the mark variogram $\gamma_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$ quantifies the variability of marks for pairs of points with an interpoint distance $r_{\mathbb{L}}$. Note that, in contrast to the classic mark variogram $\gamma_{mm}(r)$, given in Table 1, which disregards the underlying geometry of the linear network \mathbb{L} and, thus, may yield biased outcomes and potentially mislead interpretations (see Sect. 4.2.2), the mark variogram $\gamma_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$ takes the specific properties of the linear network into account and provide more accurate outcomes. The large/small local differences, i.e., high/low local variation, between marks give rise to large/small values for the mark variogram function, deviating from the mark independence assumption. Similarly, the geo-

Table 3. Test functions for point processes on linear networks with real-valued marks. The average and variance of all marks are denoted by $\mu_{m,\mathbb{L}}$ and $\sigma_{m,\mathbb{L}}^2$, and $\mu_{m,\mathbb{L}}(r_{\mathbb{L}})$ is the conditional mean of the marks for points with an interpoint distance $r_{\mathbb{L}}$

Name of the function	Symbol	Test function ($t_f^{\mathbb{L}}$)	Normalizing factor ($c_{t_f^{\mathbb{L}}}$)
Mark variogram	$\gamma_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$	$0.5[m(x) - m(y)]^2$	$\sigma_{m,\mathbb{L}}^2$
Stoyan's mark correlation function	$\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$	$m(x)m(y)$	$\mu_{m,\mathbb{L}}^2$
$\mathbf{r}^{\mathbb{L}}$ -mark correlation function	$\kappa_{m\bullet}^{\mathbb{L}}(r_{\mathbb{L}})$	$m(x)$	$\mu_{m,\mathbb{L}}$
$\mathbf{r}^{\mathbb{L}}$ -mark correlation function	$\kappa_{\bullet m}^{\mathbb{L}}(r_{\mathbb{L}})$	$m(y)$	$\mu_{m,\mathbb{L}}$
Beisbart and Kerscher's $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$	$\kappa_{mm}^{\text{Bei},\mathbb{L}}(r_{\mathbb{L}})$	$m(x) + m(y)$	$2\mu_{m,\mathbb{L}}$
Isham's $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$	$\kappa_{mm}^{\text{Ish},\mathbb{L}}(r_{\mathbb{L}})$	$m(x)m(y) - (\mu_{m,\mathbb{L}})^2$	$\sigma_{m,\mathbb{L}}^2$
Stoyan's mark covariance function	$\text{Cov}_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$	$m(x)m(y) - (\mu_{m,\mathbb{L}})^2$	1
Schlather's I	$I_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$	$(m(x) - \mu_{m,\mathbb{L}}(r_{\mathbb{L}}))(m(y) - \mu_{m,\mathbb{L}}(r_{\mathbb{L}}))$	$\sigma_{m,\mathbb{L}}^2$
Shimanti's I	$I_{mm}^{\text{Shi},\mathbb{L}}(r_{\mathbb{L}})$	$(m(x) - \mu_{m,\mathbb{L}})(m(y) - \mu_{m,\mathbb{L}})$	$\sigma_{m,\mathbb{L}}^2$

metrically corrected version of Stoyan's mark correlation $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$ quantifies the average pairwise product of marks for points separated by a distance $r_{\mathbb{L}}$ along the linear network \mathbb{L} . If marks are independent, it coincides with the mark mean squared $\mu_{m,\mathbb{L}}^2$ providing that $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}}) = 1$. If marks for nearby points on a linear network differ from the expected mark under independence, their average would also be different from $\mu_{m,\mathbb{L}}^2$ and thus $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}}) \neq 1$. Similarly, the $\mathbf{r}^{\mathbb{L}}$ -mark correlation functions would only coincide with the mark mean if the marks are independent subject to a distance $r_{\mathbb{L}}$. Consequently, any deviations of $\mathbf{r}^{\mathbb{L}}$ -mark correlation functions from one indicate the existence of a structure in the distribution of marks. The geometrically corrected version of Beisbart and Kerscher's mark correlation $\kappa_{mm}^{\text{Bei},\mathbb{L}}(r_{\mathbb{L}})$ expects that the average pairwise sum of marks is expected to be equal to twice the mark mean under independence. Thus, deviations of $\kappa_{mm}^{\text{Bei},\mathbb{L}}$ from unity highlight local dependencies and the existence of a particular structure among marks. In a similar manner, one may describe other mark characteristics presented in Table 3.

In addition to the above test functions, we further, for a homogeneous point process $X^{\mathbb{L}}$ with constant intensity $\lambda^{\mathbb{L}}$, extend the mark-weighted K -functions $K_{t_f}(r)$, given in (13), to the linear network settings, and propose its counterpart as

$$K_{t_f^{\mathbb{L}}}(r_{\mathbb{L}}) = \mathbb{E} \left[\frac{1}{c_{t_f^{\mathbb{L}}}\lambda^{\mathbb{L}}} \sum_{\substack{x \in X^{\mathbb{L}} \\ u \neq x}} t_f^{\mathbb{L}}(m(x), m(u)) \mathbf{1}\{d_{\mathbb{L}}(x, u) \leq r_{\mathbb{L}}\} \nabla(u, d_{\mathbb{L}}(u, x)) \Big| u \in X^{\mathbb{L}} \right], \quad (30)$$

which becomes

$$K_{t_f^{\mathbb{L}}}^{\text{inhom}}(r_{\mathbb{L}}) = \mathbb{E} \left[\frac{1}{c_{t_f^{\mathbb{L}}}} \sum_{\substack{x \in X^{\mathbb{L}} \\ u \neq x}} t_f^{\mathbb{L}}(m(x), m(u)) \frac{\mathbf{1}\{d_{\mathbb{L}}(x, u) \leq r_{\mathbb{L}}\} \nabla(u, d_{\mathbb{L}}(u, x))}{\lambda^{\mathbb{L}}(x)} \Big| u \in X^{\mathbb{L}} \right] \quad (31)$$

for inhomogeneous point processes. Implementing the test function $t_f^{\mathbb{L}}(\cdot)$ as a weight in the geometrically corrected K -function for spatial point processes on linear networks as in (30) and (31), gives rise to scaled versions of the original K -functions. In particular, since the implemented weight becomes one if marks are independent, the up/down-scaled version of the original K -function highlights some interaction between marks. Under the assumption that points are generated from a Poisson process with independent marks, the above mark-weighted K -functions possess a known value denoted by $r_{\mathbb{L}}$. However, due to the presence of two sources re-scaling the K -function, namely pairwise interaction between points and between marks, it is advisable to compute both the weighted and unweighted versions for a more comprehensive understanding. Note that the weighting idea is applicable to other summary characteristics such as the inhomogeneous empty space function, the inhomogeneous nearest-neighbor distance distribution function, and the inhomogeneous J -function.

Next, for a homogeneous point process $X^{\mathbb{L}}$ with the pair correlation $\rho^{\mathbb{L}}(\cdot)$, we propose an extension of (14) as

$$U^{\mathbb{L}}(r_{\mathbb{L}}) = [\lambda^{\mathbb{L}}]^2 \rho^{\mathbb{L}}(r_{\mathbb{L}}) \kappa_{t_f^{\mathbb{L}}}^{\mathbb{L}}(r_{\mathbb{L}}) |d_1 x| |d_1 y|, \quad (32)$$

where $|d_1 x|$ and $|d_1 y|$ are the sizes of two infinitesimal segments around the points $x, y \in \mathbb{L}$; similar interpretation, as the planar case, holds for $U^{\mathbb{L}}(r_{\mathbb{L}})$.

4.2.2. Numerical Evaluation

This section is devoted to evaluating the performance of the proposed linear-network-based mark characteristics and investigating potential variations between the classic planar mark correlation functions and those that take the underlying linear network into account. In terms of the spatial distribution of points, following the assumptions concerning the definition of mark correlation functions, we consider homogeneous point patterns, with 100 uniformly distributed points, on a dendrite network, which is previously used in [Jammalamadaka et al. \(2013\)](#); [Baddeley et al. \(2014\)](#). Next, to convert the simulated point patterns into marked point patterns, we generate real-valued quantities according to three distinct scenarios, called models I, II, and III, and assign the simulated marks to points. In model I, marks follow the function $f(x, y) = (x + y)/5000$, having a trend from bottom-left to top-right of the network; in model II, marks are the shortest-path distance from each point to the dendrite's border, and in model III marks are the number of points which have a distance less than 80 units to a target point, i.e., the number of nearest neighbor points. Figure 3 shows examples of the considered models. Note that within all three considered models, there is a clear structure for the marks on the dendrite network.

From each model, we simulate 199 patterns, and then, for each simulated point pattern, we calculate the Stoyan's mark correlation function $\kappa_{mm}(r)$ and our proposed network-based mark correlation function $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$; we let $d_{\mathbb{L}}$ be the shortest-path distance. We also let $r_{\mathbb{L}} = 1.25r$ ([Rakshit et al. 2019](#)), leading to $\kappa_{mm}(r)$ and $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$ being evaluated over distances $r \in [0, 200]$ and $r_{\mathbb{L}} \in [0, 250]$, respectively. Based on the 199 estimated mark correlation functions, we obtain 95% pointwise critical envelopes to compare the general




Figure 3. Examples of simulated marked point patterns. From left to right: Models I, II, and III. Colours show mark values.




Figure 4. 95% pointwise critical envelopes for the Stoyan's mark correlation function $\kappa_{mm}(r)$ and our proposed network-based mark correlation function $\kappa_{mm}^L(r_L)$ based on 199 simulated patterns from Models I, II, and III. From left to right: $\kappa_{mm}(r)$, $\kappa_{mm}^L(r_L)$, and their average. From top to bottom: Model I, Model II, and Model III.

behaviors of $\kappa_{mm}(r)$ and $\kappa_{mm}^L(r_L)$ when the data are generated on a dendrite network. Figure 4 shows the estimated Stoyan's mark correlation functions for all models together with their 95% pointwise envelopes and average behavior. Looking at the top row of Fig. 4, show-

ing the results for the model I, while the structure of the marks is not revealed by Stoyan's mark correlation function $\kappa_{mm}(r)$ (left panel), indicating mark independence, our proposed linear-network-based Stoyan's mark correlation functions $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$ (central panel) clearly reveals a positive mark correlation for shortest-path distances up to around 175 units. For larger distances, the mark correlation becomes less than one, which means that the average product of marks for points that are apart by a distance higher than 175 units is smaller than the squared mark mean value. To better compare the average behavior of $\kappa_{mm}(r)$ and $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$, the averaged mark correlation functions are displayed in the right panel. Similarly, model II (middle row) and model III (bottom row) show apparent dissimilarities between the planar and the linear network mark correlation functions. Looking at the middle row and the presented results for model II, one can see a similar tendency of having a high mark correlation for smaller distances, which turns out to be lower than the correlation value under mark independence for larger distances. However, the clear difference between the two is that $\kappa_{mm}(r)$ turns from positive to negative quickly, around $r = 100$, whereas $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$ maintain values higher than the expected value under mark independence up to $r_{\mathbb{L}} = 200$. Note that as the shortest-path distances from the points at the border to the central point on the dendrite network increase in value, central points are surrounded mainly by higher mark values. In contrast, very small marks appeared only very close to the dendrite's border. Even though both $\kappa_{mm}(r)$ and $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$ may show a positive correlation for smaller distances and a negative correlation for larger distances, their main difference is in the degree of positiveness/negativeness and further on the turning point from positive to negative. This impression also holds for the obtained results for model III displayed in the bottom row. While the $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$ clearly show the structure of the marks for small distances, increasing until $r_{\mathbb{L}} = 50$ and then decreasing, $\kappa_{mm}(r)$ entirely show a decreasing trend from small to large distances. Note also the difference between the turning points.

Upon further comparison between $\kappa_{mm}(r)$ and $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$, we proceed by selecting a single realization from each model. We then compare Stoyan's mark correlation function of these realizations to a 95% pointwise envelope under random labelling. This envelope is constructed using 199 point patterns with randomly allocated marks derived from the original pattern. Looking at Fig. 5, a clear difference between the envelopes for $\kappa_{mm}(r)$ and $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$ is that $\kappa_{mm}(r)$ shows a quite higher variation for larger distances compared to $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$, especially for model II displayed in the middle column. Another significant distinction is the overall fluctuation pattern of $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$ compared to $\kappa_{mm}(r)$, particularly with respect to their containment within the envelopes, indicating different conclusions. By joining the findings from both Figs. 4 and 5, it becomes evident that disregarding the impact of underlying linear networks could potentially lead to inaccurate conclusions.

5. REAL DATA ANALYSIS

We now study the mark structure for two real data presented in Sect. 2. For the case of influenza virus proteins, we further let the data points be labelled by the intensities so that we can make use of both cross-type and mark correlation functions in our analysis. Intensities




Figure 5. 95% pointwise critical envelopes for Stoyan’s mark correlation function $\kappa_{mm}(r)$ and our proposed network-based mark correlation function $\kappa_{mm}^L(r_L)$, under random labelling, based on 199 simulated patterns. The solid curves show $\kappa_{mm}(r)$ and $\kappa_{mm}^L(r_L)$. From left to right: model I, II, and III .

are, separately for each type, estimated via the non-parametric Jones–Diggle kernel-based estimators (Jones 1993) after employing Scott’s rule of thumb for bandwidth selection (Scott 2015). In the case of public street trees in Vancouver, Canada, we only make use of our proposed version of Stoyan’s mark correlation function for point processes on linear networks and make an additional comparison with its counterpart for point processes in \mathbb{R}^2 in the case of real datasets. We point out that all two real datasets exhibit inhomogeneity, which is expected when dealing with real phenomena. However, we believe mark correlation functions can serve as good indicators for discovering interactions between marks, although they are developed for stationary and homogeneous point processes.

5.1. INFLUENZA VIRUS PROTEINS

Figure 6 shows the cross-type summary characteristics and Stoyan’s mark correlation function concerning the locations of two different proteins on the membranes of cells infected with the influenza virus, namely M2 and HA. In the cross-type inhomogeneous K -function case, we only use its estimator with an isotropic edge correction. For small distances ($r \leq 400$), the cross-type inhomogeneous K -function moves around its theoretical value for Poisson processes, meaning that the two types of proteins HA and M2 are uncorrelated, which in turn may mean a diverse and widespread attachment of the virus to host cell surfaces. However, for larger distances, the cross-type inhomogeneous K -function lies below its theoretical value for Poisson processes, indicating that, at those distances, the number of proteins of type HA around those of type M2 is smaller than that of complete spatial




Figure 6. Mark summary characteristics for locations of two different proteins on the membranes of cells infected with the influenza virus, namely M2 and HA. From left to right, cross-type inhomogeneous K -function (values are multiplied by 0.01 for graphical reasons), cross-type inhomogeneous J -function, Stoyan's mark correlation function for influenza virus proteins HA, and Stoyan's mark correlation function for influenza virus proteins M2. In the case of Stoyan's mark correlation function, envelopes are 95% pointwise critical envelopes based on 199 simulated patterns under random labelling .

randomness. The reduced presence of HA proteins around M2 at these distances might signify a spatial rearrangement during a later phase, highlighting the dynamic interplay between these viral proteins throughout the infection process. A similar message is conveyed by the cross-type inhomogeneous J -function.

Turning to Stoyan's mark correlation function $\kappa_{mm}(r)$, we first obtain 199 point patterns for each type of Protein by randomly allocating marks. Then, we compare Stoyan's mark correlation function $\kappa_{mm}(r)$ for each type of Protein with their corresponding 95% pointwise critical envelopes obtained from the 199 patterns under random labelling. We can see that regardless of distance r , $\kappa_{mm}(r)$ for HA stays above the envelope, indicating that the average product of intensity is higher than the expected average under random labelling; this behavior is stronger for smaller distances. In the case of M2, $\kappa_{mm}(r)$ similarly remains above the envelope for small to moderate distances and then stays below the envelope for distances larger than approximately 700.

5.2. PUBLIC STREET TREES IN VANCOUVER, CANADA

Now we employ Stoyan's mark correlation function $\kappa_{mm}(r)$ and our proposed network-based version $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$ to investigate the distribution of diameter at breast height (dbh) for the five species of public trees planted along the street network of Vancouver, Canada. Our aim is to uncover potential interactions among dbh values for trees of the same species. Similarly, we rely on 95% pointwise envelopes obtained from 199 point patterns where spatial locations are the same as the original data, but dbh values are randomly allocated. Looking at Fig. 7, one can evidently see that $\kappa_{mm}(r)$ and $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$ give rise to different conclusions, highlighting the importance of taking the underlying network into account. In the case of trees of type Arnold, both $\kappa_{mm}(r)$ and $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$ stay above the envelope but with different degrees and turning points, highlighting that for a pair of nearby trees, at least one of them has a large dbh. Interestingly, for distances around $r = 3\text{km}$, $\kappa_{mm}(r)$ stays below the corresponding envelope, while this is not the case for $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$, pointing to different conclusions. Similar conclusions hold for trees of type Populus; on the one hand $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$ stays above the corresponding envelope for distances around $r_{\mathbb{L}} = 2\text{km}$ while this is not the case for $\kappa_{mm}(r)$, on the other hand, $\kappa_{mm}(r)$ stays below the envelope for distances




Figure 7. Stoyan's mark correlation function for the five species of trees alongside the street network of Vancouver, Canada; the mark is dbh of trees. From left to right: Arnold, Populus, Involucrata, Aquifolium, Bignonioides. The top row considers the network; the bottom row ignores it .

$r = 2.5\text{km}$ while $\kappa_{mm}^{\mathbb{L}}(r_{\mathbb{L}})$ remains inside its envelope. Similar conclusions hold for other species, pointing to the impact of the underlying network. The main difference between the distribution of dbh values for the considered species might be the fact that, for small interpoint distances, mark correlation functions for all species have larger values than the expected value under mark independence, except type Populus, for which mark correlation functions stay below the corresponding envelopes for very small interpoint distances. Note that this means for a pair of trees of the type Populus with very small interpoint distances, the dbh value for at least one of them is pretty small.

6. DISCUSSION

Methods for analyzing spatial point processes, when each point is augmented by some auxiliary quantity, have recently witnessed impressive developments. Apart from some extensions of the current methodological toolboxes to the case of point processes with events on linear networks/graphs (Jammalamadaka et al. 2013; Baddeley et al. 2014), parts of the recent literature proposed generalizations to more challenging scenarios where marks are non-scalar. Extensions to the settings where marks are function-valued were considered, followed by some recent contributions covering functional characteristics, where marks are inherently structured objects including compositions, densities, and graphs (Ghorbani et al. 2021; Eckardt et al. 2023, 2024). In this paper, we have restricted our focus to general marked summary characteristics when points live on either planar spaces or linear networks and marks are integer/real/object-valued. By combining the general marked characteristic ideas with linear network settings, we have not only reviewed the current summary characteristics for point processes in \mathbb{R}^2 but, in particular, introduced novel summary characteristics for marked point processes on linear networks. We point out that the literature further covers some extensions of general point process models to marked settings. We here have not covered models for marked point processes, however, the discussed summary characteristics are valuable tools as initial steps within the model formulation.

Despite all methodological progress, the study of marked inhomogeneous point processes is still in its infancy. Note that (i) it is quite rare to have stationary point processes in practice, and (ii) stationarity for point processes on linear networks is quite limited and challenging (Baddeley et al. 2017; Cronie et al. 2020). In fact, among all developed methods, only the cross/dot-type summary characteristics and mark-weighted summary characteristics are defined for both homogeneous and inhomogeneous point processes. Note further that one may create mark subsets out of the real-values marks and then employ cross/dot-type summary characteristics. These could potentially limit the analysis of marked point processes, especially when marks are real-valued and/or object-valued (Eckardt et al. 2023). Further, the growing availability of complex point patterns calls for suitable methods to investigate the dependencies between different types of object-valued marks, e.g., compositions and shapes, which, in practice, is not available under the present methodological toolbox. Besides, there are various introduced local indicators of spatial association (LISA) in the literature (Anselin 1995; Cressie and Collins 2001), but their extensions for marked point processes have not been established so far; current mark characteristics do not disclose local behaviors. In addition, although still recognized within a small niche of research, spatial frequency domain methods might become useful tools for modern mark scenarios and help avoid burdensome computations. It would also be interesting to account for possible directional behaviors of marks; note the availability of (second-order) directional analysis of spatial point processes on both planar spaces and linear networks (Rajala et al. 2018; Moradi et al. 2021).

Apart from the clear lack of suitable mark summary characteristics for inhomogeneous marked point processes, there is also a need for extensions of any such method to spatio-temporal processes; an application of such a case is trajectories (Moradi 2018) where, depending on the application, spatial locations of points might be network-limited or not. We note that Iftimi et al. (2019) proposed some marked second-order reduced moment measures and K -functions for inhomogeneous point processes with particular focus on space-time settings where spatial locations are in \mathbb{R}^2 .

Further, in the case of function-valued marks, marks might undergo abrupt and/or gradual changes, which may not be accounted for in the current settings (Moradi et al. 2023). Apart from the discussed methods, suitable extensions of point-to-mark tools (Schlather et al. 2004) to more challenging state spaces and/or mark settings have not been established. We further highlight the lack of methods for marked point processes on a sphere. As a final note, it would also be interesting to develop methodologies and test functions that can incorporate negative marks.

ACKNOWLEDGEMENTS

The authors are grateful to the review team for comments that improved this manuscript. The authors also gratefully acknowledge financial support through the German Research Association and the Stochastic Group of the German Mathematical Society. Matthias Eckardt was funded by the Walter Benjamin grant 467634837 from the German Research Foundation. Mehdi Moradi received travel support from the Stochastic Group of the German Mathematical Society. Our datasets, R codes used concerning the mark correlation functions for point processes on linear networks, all R codes used for simulation studies and real data analysis, all R outputs and plots, are publicly available at <https://github.com/Moradii/Marklpp>.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Funding Open access funding provided by Umea University.

Declarations

Conflict of interest The authors have no conflicts of interest to declare.

[Received September 2023. Revised January 2024. Accepted January 2024.]

REFERENCES

- Ang W, Baddeley A, Nair G (2012) Geometrically corrected second order analysis of events on a linear network, with applications to ecology and criminology. *Scand J Stat* 39:591–617
- Anselin L (1995) Local indicators of spatial association - LISA. *Geol Anal* 27(2):93–115
- Baddeley A (2010) Multivariate and marked point processes. In: Gelfand AE, Diggle P, Fuentes M, Guttorp P (eds) *Handbook of spatial statistics*. Chapman & Hall/CRC, pp 371–402
- Baddeley A, Jammalamadaka A, Nair G (2014) Multitype point process analysis of spines on the dendrite network of a neuron. *J Royal Stat Soc Ser C (Appl Stat)* 63(5):673–694
- Baddeley A, Møller J, Waagepetersen R (2000) Non- and semi-parametric estimation of interaction in inhomogeneous point patterns. *Stat Neerl* 54(3):329–350
- Baddeley A, Nair G, Rakshit S, McSwiggan G (2017) Stationary point processes are uncommon on linear networks. *Stat* 6(1):68–78
- Baddeley A, Nair G, Rakshit S, McSwiggan G, Davies TM (2021) Analysing point patterns on networks—a review. *Spat Stat* 42:100435
- Baddeley A, Rubak E, Turner R (2015) *Spatial point patterns: methodology and applications with R*. CRC Press
- Bartlett MS (1964) The spectral analysis of two-dimensional point processes. *Biometrika* 51:299–311
- ayisa FL, Ådahl M, Rydén P, Cronie O (2023) Regularised semi-parametric composite likelihood intensity modelling of a swedish spatial ambulance call point pattern. *J Agric Biol Environ Stat* 1–20
- Brillinger D (1981) *Time series: data analysis and theory*. Holt, Rinehart and Winston, New York
- Capobianco R, Renshaw E (1998) The autocovariance function for marked point processes: a comparison between two different approaches. *Biom J* 40(4):431–446
- Chen BJ, Leser GP, Jackson D, Lamb RA (2008) The influenza virus m2 protein cytoplasmic tail interacts with the m1 protein and influences virus assembly at the site of virus budding. *J Virol* 82(20):10059–10070
- Comas C, Delicado P, Mateu J (2008) Analysing spatial point patterns with associated functional data. In: D. Cocchi, J. Mateu, F. Montes, E. Otranto, E. Porcu, and A. Usai (Eds.), *Statistics for Spatio-temporal modelling. Proceedings of the 4th international workshop on spatio-temporal modelling (METMA-4)*, pp 157–163
- Comas C, Delicado P, Mateu J (2011) A second order approach to analyse spatial point patterns with functional marks. *Test* 20(3):503–523
- Comas C, Mehtätalo L, Miina J (2013) Analysing space-time tree interdependencies based on individual tree growth functions. *Stoch Environ Res Risk Assess* 27(7):1673–1681
- Cressie N (1993) *Statistics for spatial data*. Wiley

- Cressie N, Collins LB (2001) Analysis of spatial point patterns using bundles of product density LISA functions. *J Agric Biol Environ Stat* 6(1):118–135
- Cronie O, Moradi M, Mateu J (2020) Inhomogeneous higher-order summary statistics for point processes on linear networks. *Stat Comput* 30(5):1221–1239
- Cronie O, van Lieshout MNM (2016) Summary statistics for inhomogeneous marked point processes. *Ann Inst Stat Math* 68:905–928
- D'Angelo N, Adelfio G, Mateu J, Cronie O (2023) Local inhomogeneous weighted summary statistics for marked point processes. *J Comput Graph Stat* 1–15. <https://doi.org/10.1080/10618600.2023.2206441>
- Daniel J, Horrocks J, Umphrey GJ (2020) Efficient modelling of presence-only species data via local background sampling. *J Agric Biol Environ Stat* 25:90–111
- Eckardt M, Comas C, Mateu J (2023) Summary characteristics for multivariate function-valued spatial point process attributes. Submitted for publication
- Eckardt M, Ghorbanpour F, Särkkä A (2024) Second-order characteristics for spatial point processes with graph-valued mark. Submitted for publication
- Eckardt M, Gonzáles J, Mateu J (2021) Graphical modelling and partial characteristics for multitype and multivariate-marked spatio-temporal point processes. *Comput Stat Data Anal* 156:107139
- Eckardt M, Greven S, Myllymäki M, (2023) On spatial point processes with composition-valued marks. Submitted for publication
- Eckardt M, Mateu J (2018) Point patterns occurring on complex structures in space and space-time: an alternative network approach. *J Comput Graph Stat* 27(2):312–322
- Eckardt M, Mateu J (2019) Analysing multivariate spatial point processes with continuous marks: a graphical modelling approach. *Int Stat Rev* 87(1):44–67
- Eckardt M, Mateu J (2019) Partial characteristics for marked spatial point processes. *Environmetrics* 30(6):e2565
- Eckardt M, Mateu J (2021) Second-order and local characteristics of network intensity functions. *Test* 30(2):318–340
- Fedriani JM, Wiegand T, Calvo G, Suárez-Esteban A, Jácome M, Zywiec M, Delibes M (2015) Unravelling conflicting density- and distance-dependent effects on plant reproduction using a spatially explicit approach. *J Ecol* 103(5):1344–1353
- Feigelson ED, Babu GJ (2012) *Modern statistical methods for astronomy: with R applications*. Cambridge University Press
- Flagg KA, Hoegh A, Borkowski JJ (2020) Modeling partially surveyed point process data: inferring spatial point intensity of geomagnetic anomalies. *J Agric Biol Environ Stat* 25:186–205
- Gavrikov V, Stoyan D (1995) The use of marked point processes in ecological and environmental forest studies. *Environ Ecol Stat* 2(4):331–344
- Ghorbani M, Cronie O, Mateu J, Yu J (2021) Functional marked point processes: a natural structure to unify spatio-temporal frameworks and to analyse dependent functional data. *Test* 30:529–568
- Guan Y (2006) Tests for independence between marks and points of a marked point process. *Biometrics* 62(1):126–134
- Guan Y, Afshartous DR (2007) Test for independence between marks and points of marked point processes: a subsampling approach. *Environ Ecol Stat* 14:101–111
- Harkness RD, Isham V (1983) A bivariate spatial point pattern of ants' nests. *J Royal Stat Soc Ser C (Appl Stat)* 32(3):293–303
- Hui G, Pommerening A (2014) Analysing tree species and size diversity patterns in multi-species uneven-aged forests of northern China. *For Ecol Manag* 316:125–138
- Iftimi A, Cronie O, Montes F (2019) Second-order analysis of marked inhomogeneous spatiotemporal point processes: applications to earthquake data. *Scand J Stat* 46(3):661–685
- Illian J, Penttinen A, Stoyan H, Stoyan D (2008) *Statistical analysis and modelling of spatial point patterns*. John Wiley & Sons, New York

- Jammalamadaka A, Banerjee S, Manjunath BS, Kosik KS (2013) Statistical analysis of dendritic spine distributions in rat hippocampal cultures. *BMC Bioinf* 14:1–19
- Jones MC (1993) Simple boundary correction for kernel density estimation. *Stat Comput* 3:135–146
- Lewandowski A, Pommerening A (1997) Zur Beschreibung der Waldstruktur – Erwartete und beobachtete Arten-Durchmischung. *Forstwissenschaftliches Centralblatt vereinigt mit Tharandter forstliches Jahrbuch* 116(1):129–139
- Lotwick HW, Silverman BW (1982) Methods for analysing spatial processes of several types of points. *J Royal Stat Soc Ser B (Methodological)* 44(3):406–413
- Mateu J, Moradi M, Cronie O (2020) Spatio-temporal point patterns on linear networks: pseudo-separable intensity estimation. *Spat Stat* 37:100400
- McSwiggan G, Baddeley A, Nair G (2017) Kernel density estimation on a linear network. *Scand J Stat* 44(2):324–345
- Møller J, Waagepetersen R (2003) *Statistical inference and simulation for spatial point processes*. Chapman and Hall/CRC Boca Raton
- Moradi M (2018) *Spatial and spatio-temporal point patterns on linear networks*. PhD dissertation, University Jaume I
- Moradi M, Cronie O, Pérez-Goya U, Mateu J (2023) Hierarchical spatio-temporal change-point detection. *The American statistician*, 1–11
- Moradi M, Cronie O, Rubak E, Lachieze-Rey R, Mateu J, Baddeley A (2019) Resample-smoothing of Voronoi intensity estimators. *Stat Comput* 29(5):995–1010
- Moradi M, Mateu J (2020) First- and second-order characteristics of spatio-temporal point processes on linear networks. *J Comput Graph Stat* 29(3):432–443
- Moradi M, Mateu J, Comas C (2021) Directional analysis for point patterns on linear networks. *Stat* 10(1):e323
- Moradi M, Rodríguez-Cortés F, Mateu J (2018) On kernel-based intensity estimation of spatial point patterns on linear networks. *J Comput Graph Stat* 27(2):302–311
- Mugglestone MA, Renshaw E (1996) The exploratory analysis of bivariate spatial point pattern using cross-spectra. *Environmetrics* 7:361–377
- Okabe A, Yamada I (2001) The K -function method on a network and its computational implementation. *Geograph Anal* 33(3):271–290
- Penttinen A, Stoyan D (1989) Statistical analysis for a class of line segment processes. *Scand J Stat* 16(2):153–168
- Penttinen A, Stoyan D, Henttonen HM (1992) Marked point processes in forest statistics. *For Sci* 38(4):806–824
- Pommerening A, Gonçalves AC, Rodríguez-Soalleiro R (2011) Species mingling and diameter differentiation as second-order characteristics. *Allgemeine Forst- und Jagdzeitung* 182(7/8):115–129
- Pommerening A, Grabarnik P (2019) *Individual-based methods in forest ecology and management*. Springer Cham
- Rajala T, Redenbach C, Särkkä A, Sormani M (2018) A review on anisotropy analysis of spatial point patterns. *Spat Stat* 28:141–168
- Rakshit S, Baddeley A, Nair G (2019) Efficient code for second order analysis of events on a linear network. *J Stat Softw* 90:1–37
- Rakshit S, Davies TM, Moradi M, McSwiggan G, Nair G, Mateu J, Baddeley A (2019) Fast kernel smoothing of point patterns on a large network using two-dimensional convolution. *Int Stat Rev* 87(3):531–556
- Rakshit S, Nair G, Baddeley A (2017) Second-order analysis of point patterns on a network using any distance metric. *Spat Stat* 22:129–154
- Ramsay J, Silverman B (1997) *Functional data analysis*. Springer
- Raventós J, Mujica E, Wiegand T, Bonet A (2011) Analyzing the spatial structure of *broughtonia cubensis* (orchidaceae) populations in the dry forests of Guanahacabibes Cuba. *Biotropica* 43(2):173–182
- Renshaw E (2002) Two-dimensional spectral analysis for marked point processes. *Biom J* 44:718–745
- Renshaw E, Ford E (1984) The description of spatial pattern using two-dimensional spectral analysis. *Vegetatio* 56:75–85

- Renshaw E, Ford ED (1983) The interpretation of process from pattern using two-dimensional spectral analysis: methods and problems of interpretation. *Appl Stat* 32:51–63
- Ripley BD (1976) The second-order analysis of stationary point processes. *J Appl Probab* 13:255–266
- Russell JC, Hanks EM, Haran M (2016) Dynamic models of animal movement with spatial point process interactions. *J Agric Biol Environ Stat* 21:22–40
- Schlather M (2001) On the second-order characteristics of marked point processes. *Bernoulli* 7(1):99–117
- Schlather M, Riberio P, Diggle P (2004) Detecting dependence between marks and locations of marked point processes. *J Royal Stat Soc Ser B (Methodological)* 66:79–93
- Scott DW (2015) *Multivariate density estimation: theory, practice, and visualization*. John Wiley & Sons
- Shimatanı K (2002) Point processes for fine-scale spatial genetics and molecular ecology. *Biom J* 44(3):325–352
- Spooner PG, Lunt ID, Okabe A, Shiode S (2004) Spatial analysis of roadside acacia populations on a road network using the network k-function. *Landsc Ecol* 19(5):491–499
- Stoyan D (1984) Correlations of the marks of marked point processes - statistical inference and simple models. *Elektronische Informationsverarbeitung und Kybernetik* 20(5/6):285–294
- Stoyan D (1987) Statistical analysis of spatial point processes: a soft-core model and cross-correlations of marks. *Biom J* 29(8):971–980
- Stoyan D, Stoyan H (1994) *Fractals, random shapes, and point fields: methods of geometrical statistics*. Wiley, Chichester, New York
- Stoyan D, Walder O (2000) On variograms in point process statistics, ii: models for markings and ecological interpretation. *Biom J* 42:171–187
- van Lieshout MNM (2006) A J-function for marked point patterns. *Ann Inst Stat Math* 58(2):235–259
- van Lieshout MNM (2011) A J-function for inhomogeneous point processes. *Stat Neerl* 65(2):183–201
- van Lieshout MNM, Baddeley A (1996) A nonparametric measure of spatial interaction in point patterns. *Stat Neerl* 50(3):344–361
- van Lieshout MNM, Baddeley A (1999) Indices of dependence between types in multivariate point patterns. *Scand J Stat* 26(4):511–532
- Walder O, Stoyan D (1996) On variograms in point process statistics. *Biom J* 38(8):895–905
- Wiegand T, Moloney KA (2013) *Handbook of spatial point-pattern analysis in ecology*. Chapman and Hall/CRC
- Xie Z, Yan J (2008) Kernel density estimation of traffic accidents in a network space. *Comput Environ Urban Syst* 32(5):396–406
- Yazigi A, Penttinen A, Ylitalo A-K, Maltamo M, Packalen P, Mehtatalo L (2021) Modeling forest tree data using sequential spatial point processes. *J Agric Biol Environ Stat* 27:1–21

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.