



Correction to: Nitric oxide-caused rabbit chondrocyte apoptosis is linked to cytoskeletal protein proteolysis anomaly through intracellular JNK and ERK signal pathways

Qun Chen¹ · Xibin Kao² · Yan Gao³ · Jinghong Chen¹ · Zhaoheng Dong⁴ · Chen Chen⁵

Published online: 3 August 2022

© The Author(s) under exclusive licence to The Korean Society of Toxicogenomics and Toxicoproteomics 2022

Correction to: Molecular & Cellular Toxicology
<https://doi.org/10.1007/s13273-022-00241-1>

In this article Fig. 5 has been updated and should be read as follows (Fig. 5):

The original article can be found online at <https://doi.org/10.1007/s13273-022-00241-1>.

✉ Qun Chen
chenqun2003076@163.com

Xibin Kao
xbkao1980@163.com

Yan Gao
gaoyan2017@yeah.net

Jinghong Chen
896939179@qq.com

Zhaoheng Dong
dongzhaoheng@sohu.com

Chen Chen
chen.chen@uq.edu.au

Commission of the People's Republic of China, Key Laboratory for Disease Prevention and Control and Health Promotion of Shaanxi Province, Xi'an, Jiaotong University Health Science Center, Xi'an 710061, Shaanxi, People's Republic of China

² Institute for Hygiene of Ordnance Industry, Xi'an 710061, Shaanxi, People's Republic of China

³ Institute of Health Supervision, Beilin District, Xi'an 710003, Shaanxi, People's Republic of China

⁴ Shandong Shenghua Electronic New Materials Co., Ltd., Yantai, Shandong, China

⁵ Endocrinology, School of Biomedical Sciences, Faculty of Medicine, University of Queensland, Brisbane, Australia

¹ Institute of Endemic Diseases, Key Laboratory of Trace Elements and Endemic Diseases, National Health




Fig. 5 The effects of PD98059 and SP600125 on cytoskeletal proteins and mRNA expression in SNP-treated chondrocytes. **A** The cells were treated with different concentrations of SNP for 30 min and the levels of actin, tubulin, vimentin was analyzed by Western blot. The results were obtained from 3 independent experiments. **B** The cells were treated with different concentrations of PD98059. **C** The cells were treated with different concentrations of SP600125. **D** Representative blots are shown. **E** The mRNA expression of tubulin

and vimentin were calculated in response to different concentrations of SNP. **F** The mRNA expression of tubulin and vimentin in response to PD98059 in presence or absence of 1.0 mM SNP. **G** The mRNA expression of tubulin and vimentin in response to SP600125 in presence or absence of 1.0 mM SNP. * $P < 0.05$ indicated statistically significant different to that of the Control group. # $P < 0.05$ indicated statistically significant difference to the 1.0 mM SNP alone group

The original article has been corrected.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.