



RESEARCH ARTICLE

# Modeling place-based nature-based solutions to promote urban carbon neutrality

Cong Cong, Haozhi Pan , Jessica Page, Stephan Barthel, Zahra Kalantari

Received: 25 July 2022 / Revised: 12 January 2023 / Accepted: 13 April 2023 / Published online: 15 May 2023

**Abstract** Nature-based solutions (NbS) are recognized as widely available and cost-effective mechanisms for sequestering carbon and offsetting carbon emissions. Realistic NbS implementations for carbon neutrality need to be effective at the global level and also appropriate for the socio-economic and physical conditions prevailing at the local level. This paper presents a framework that can help stakeholders identify demands, locations, and types of NbS interventions that could maximize NbS benefits at the local scale. Key processes in the framework include (1) interpolating carbon emissions data at larger spatial scales to high-resolution cells, using land use and socio-economic data; (2) assessing NbS effects on carbon reduction and their location-related suitability, through qualitative literature review, and (3) spatially allocating and coupling multiple NbS interventions to land use cells. The system was tested in Stockholm, Sweden. The findings show that the urban center should be allocated with combinations of improving access to green spaces and streetscapes, while the rural and suburban areas should prioritize preserving and utilizing natural areas. Our proposed method framework can help planners better select target locations for intended risk/hazard-mitigating interventions.

**Keywords** Carbon emissions · Carbon neutrality · Ecosystem services · Land use · Nature-based solutions

## INTRODUCTION

Urban areas contribute more than 60% of global greenhouse gas (GHG) emissions, through residential, commercial, and transportation activities (Lanau et al. 2021). Carbon sequestration by urban vegetation and natural

amenities, such as parks, gardens, and residential lawns, can be a cost-effective mechanism for offsetting carbon emissions (Kalantari et al. 2019; Li and Wang 2021). Nature-based solutions (NbS) are increasingly being adopted by cities worldwide to enhance natural capital and preserve ecosystem service values (Laforteza et al. 2018). NbS are defined by the European Commission as “solutions that are inspired and supported by nature, which are cost-effective, simultaneously provide environmental, social and economic benefits and help build resilience” (European Environment Agency 2021, p. 17).

A ‘toolbox’ developed for deploying NbS in urban settings are combinations of ecosystem service or green infrastructure approaches, including ecological restoration (Nesshöver et al. 2017; Dorst et al. 2019), soil-vegetation and landscape (Keesstra et al. 2018), biodiversity preservation (Kabisch et al. 2016), water management (Krauze and Wagner 2019; Gómez Martín et al. 2020), and integrated planning (Albert et al. 2019; Pan et al. 2021). Well-designed NbS can help mitigate carbon emissions and provide other social and economic co-benefits, including enhancing habitat quality, reducing the costs of artificial infrastructure, promoting participation and equitable access to amenities, creating jobs in the green sector, and improving recreational opportunities and health (Raymond et al. 2017; Giordano et al. 2020; Ruangpan et al. 2021).

Despite prior emphases on better NbS planning, a continuing absence of systematic approaches results in either (a) one-off and case-specific placement of infrastructure not entirely trusted by all stakeholders (Raymond et al. 2017); or (b) highly conceptualized models with low spatially detailed suitability (Connop et al. 2016; Kuller et al. 2019). Realistic NbS implementation plans toward carbon neutrality need to be both effective in mitigating carbon emissions at the global level and suitable for the socio-

economic and physical conditions at the local level. Prioritizing suitable sites and solutions can enhance the long-term viability of NbS and its ability to provide multiple ecosystem services. To achieve this, it is necessary to fully consider the spatial details that are crucial for the successful implementation of NbS, including designing interventions that target specific emission sources. Fine-scale emissions analysis can help identify localized emissions driven by human behaviors, which may not be apparent when looking at emissions data on a larger scale.

In response to these challenges, this paper presents a novel systematic framework for prioritizing NbS types and locations in an integrative manner, to maximize their potential for mitigating carbon emissions. The method involves (1) fine-scale (30 × 30 m) spatial accounting and mapping of carbon emissions from transportation, residential, and industrial activities taking place within a study region; (2) a meta-analysis identifying relevant NbS strategies and their carbon emission reduction benefits in different places and cases, where NbS strategies include both direct carbon mitigation, such as carbon storage and sequestration, and indirect carbon mitigation, through climate regulation and interventions influencing human behaviors; and (3) prioritized NbS interventions allocated across the study region based on both their potential for carbon emission mitigation and their local suitability. Using the case of Stockholm County, Sweden, we tested the capacity of this framework for tailoring and targeting emission reduction approaches and visually communicating the carbon mitigation impacts of NbS strategies.

This paper addresses two main research questions: Which NbS strategies are most effective in mitigating urban carbon emissions? And where should selected types of NbS be deployed to maximize their carbon emissions saving potential? Acknowledging that the ability of NbS to mitigate carbon emissions goes beyond the carbon sequestration potential of green infrastructure and natural amenities, this paper focuses on the potential of NbS to offset human activity-related emissions including transportation, residential, and industrial emissions, and aims to provide a finer-scale assessment of the benefits of NbS.

The remainder of this paper is organized as follows: “[Related works](#)” section summarizes previous work on NbS and carbon emissions mitigation and identifies research gaps. “[Materials and methods](#)” section introduces our modeling methods, the case study area, and data sources. “[Results](#)” section presents the results obtained in the analysis and proposes an operational framework for place-based NbS. “[Discussion](#)” section discusses potential applications of the results and policy implications. “[Conclusions](#)” section presents some conclusions and discusses limitations of the study.

## RELATED WORKS

### Adopting NbS for offsetting carbon emissions

Many studies have shown that urban NbS can provide multiple ecosystem services with environmental, social, and economic co-benefits, including climate change mitigation and adaptation (Chen 2015; De la Sota et al. 2019; Choi et al. 2021). For example, green infrastructure (GI) or green and blue infrastructure (GBI), as opposed to artificial (gray) infrastructure, can efficiently reduce urban carbon emissions. In a study by Anderson and Gough (2020), an average carbon dioxide (CO<sub>2</sub>) reduction of 6% was achieved by the application of GI in the built environment. According to Ren et al. (2019), the average annual increase in carbon storage by urban GI offsets 3.9% of the increase in urban carbon emissions in China. A study by Tomalty (2012) found that forestland, wetlands, and agricultural land in Ontario’s Greenbelt around Toronto can store 86.6 million tons of carbon and sequester 200 000 tons of carbon annually.

NbS can reduce carbon emission levels through different direct and indirect pathways. Direct pathways typically refer to the natural growth of vegetation, during which plants remove atmospheric CO<sub>2</sub> and store it in their biomass (Nowak and Crane 2002). Several studies have quantified CO<sub>2</sub> storage and sequestration by urban forests, e.g., Zhao et al. (2010) report carbon sequestration by urban forests in Hangzhou, China, of over 1.3 million metric tons (MMT) C/year, offsetting 18.6% of annual industrial carbon emissions with carbon storage equivalent to 1.75 times the annual industrial amount emitted. A later study by Chen (2015) estimated total carbon sequestration of 1.90 MMT from GI, with an average rate of 2.16 t/ha/year, in 35 major cities of China, offsetting up to 22.5% of carbon emissions from fossil fuel combustion.

NbS also reduce carbon emissions through indirect pathways, with the most prominent evidence being demonstrated in the following aspects. First, vegetation decreases building cooling demand by shading and evapotranspiration, thereby avoiding carbon emissions associated with fossil fuel use in energy production (McPherson 1998). Tsoka et al. (2021) report that adding trees can reduce buildings’ cooling energy demand by up to 54%, with foliage density and planting pattern as significant factors of the energy-saving efficacy. Green roofs generally reduce CO<sub>2</sub> emissions ranging between 1.703 and 1.889 kg/m<sup>2</sup>/year (Kuronuma et al. 2018), but the site selection of green building, as well as facilities accessibility, surrounding environments and residents’ behavior account for 11–50% of the variation in green buildings’ energy preservation (Gill et al. 2010; Gou and Lau 2014).

Second, micro-scale features in streets, such as amenities and esthetics, affect human perceptions and can encourage people to engage in more walking, biking, and other pro-environment behaviors (Smardon 1988; Sarkar et al. 2015). Streetscape factors significantly contribute to explaining walking mode choice as well as the associated reduction of automobile carbon emissions. A high odd ratio (1.680–2.070) is found for street-level urban greening to increase walking modal choices (Koo et al. 2022). Third, greenbelt policies were found effective in limiting urban growth and promoting a more compact land-use development pattern, preventing land use-associated emissions and carbon stock losses (Han et al. 2022).

A fine-scale analysis of emissions can reveal localized socio-economic drivers of emissions that may not be evident when examining emissions data on a larger scale. This information can be valuable in designing interventions that target specific emission sources, such as reducing emissions from vehicles in urban areas, to achieve maximum emissions mitigation potential. For example, NbS interventions need to be deliberately allocated to areas with major emission sources such as roads with high traffic volume (Gromke and Blocken 2015), buildings with high cooling and heating demand (Xie et al. 2017), and natural carbon stocks near urban development (Ferreira et al. 2018). Similarly, although direct emission mitigation implementations, such as carbon sequestration, can be effective in any location, their benefits can be increased substantially by an optimal land management (Sha et al. 2022). For instance, by coupling eco-environmental measures and relocating significant industrial emission sources, there is a noticeable trend of improvement in removing industrial emissions (Song et al. 2023).

### Challenges to systematically scaling up NbS

There have been recent attempts by many cities to scale up NbS through systematic design, in the hope of acquiring accrued benefits when a number of projects are implemented. Such projects may include developing new NbS and extending or linking existing NbS. Implementing NbS at a large scale requires coordination of multiple sectors, from energy and transport to land-use planning and health (Raymond et al. 2017).

Many projects have demonstrated the potential for moving from siting NbS at single locations to city- or regional-scale deployment and the difficulties in doing so, e.g., the regulatory framework, business models, innovative governance models, and social acceptance that need to be in place before expanding network and knowledge beyond single-case demonstrations (Nesshöver et al. 2017; Bradfer-Lawrence et al. 2021; Cortinovis et al. 2022). Fastenrath et al. (2020) concluded that scaling up of NbS

requires interdisciplinary expertise to address ecological, institutional, and socio-cultural challenges. In a participatory analysis of six Swedish municipalities, Wamsler et al. (2020) identified science–policy integration as a key strategy for the success of NbS implementation from a city-to-city learning lab of urban development projects.

In spatial terms, implementing NbS at a larger scale requires a better understanding of the options that fit local environments. The location and size of each NbS affect the benefits it can produce (Cortinovis and Geneletti 2018; Andersson et al. 2020). For example, planting trees can be more easily integrated into spatial plans in less built-up areas than in the urban core, where the possibilities are constrained by existing land uses and spaces (Pataki et al. 2021). Areas of high priority for stormwater abatement are generally not best suited for maximizing landscape connectivity (Meerow and Newell 2017). A more realistic plan for scaling up NbS includes investigating the areas available for different NbS types, identifying appropriate spatial scales of implementation, and evaluating benefits that can be expected from various approaches (Nesshöver et al. 2017; Bradfer-Lawrence et al. 2021; Cortinovis et al. 2022).

Among previous studies on implementing NbS on a regional scale, Midgley et al. (2021) compiled an inventory of water-related ecological infrastructure intervention projects in South Africa and established a range of typologies with specific benefits for landscape actors; Meerow and Newell (2017) developed a GIS-based approach for systematically prioritizing urban green infrastructure based on a wider range of socio-economic and environmental benefits in a heterogeneous landscape; and Cortinovis et al. (2022) showed that characteristics of urban form (i.e., density, impervious area, land use types) affect the potential and benefits of NbS. However, as spatial resolution is important in terms of estimating benefits, finer-scale analysis in such studies would provide better support for implementation. A spatially detailed, systematic, and replicable strategy for selecting and allocating NbS, and for better communicating their benefits, is still lacking.

### Spatially explicit NbS implementation for carbon emission reduction

Spatial modeling specializes in combining and synthesizing different sources of information to assist local communities, planners, and agencies in identifying “hotspots” associated with infrastructure siting, assessing potential spatial tradeoffs, and ultimately enabling more informed plans based on stakeholder input (Geertman and Stillwell 2003). Modeling the spatial interactions between NbS and various systems (e.g., land use, social, environmental) is

key in revealing the drivers, feedback, and interactions of the benefits from the large-scale implementation of NbS (Bierwagen et al. 2010; Goldenberg et al. 2017). A study by Madureira and Andresen (2014) identified spatial priority areas for green infrastructure based on two criteria, namely proximity to public green spaces and the potential to improve local temperature regulation. Norton et al. (2015) developed a multi-scale hierarchical model to prioritize GI placement and type, although they mainly focused on the cooling benefits and did not include weighting protocols. Pan et al. (2020) developed a spatially explicit land use model that couples human processes (socio-economic and land use policies) and ecological processes (GHG emissions associated with human activities that have global climate impacts) to understand GHG emissions associated with urbanization and human-driven land-use changes.

Integrating spatial information with social-technology synergy can facilitate the collaborative process in NbS planning. Sarabi et al. (2022) introduced and tested an NbS planning support system (NbS-PSS) that allows users to interact at different stages of the NbS planning process to ensure the fulfillment of societal needs and equitable distribution of ecosystem services. Venter et al. (2021) created a tool for engaging stakeholders in spatial prioritization of green roof retrofitting in Oslo, Norway, and found high spatial correlation in ecosystem services deficits.

There are two key limitations to the current approach of spatially explicit and systematic PSS models: (1) Most urban NbS assessments focus on a limited set of choices, most often street trees or green infrastructure, while little is known about how to select the most effective NbS in different urban contexts; and (2) most models identify neighborhoods for prioritized NbS intervention, while site and solution suitability assessment is not incorporated. To overcome these limitations, this paper presents a modeling framework that integrates the prioritization of NbS measures and the assessment of site suitability in various urban contexts. The

framework is sufficiently flexible and replicable to be adopted by cities with access to different types of data.


## MATERIALS AND METHODS

### Overview of methodology

The construction of a spatially explicit model to identify demands, locations, and types of NbS interventions involved three main steps. First, we studied disaggregated carbon emission maps in the global database for three different sectors (residential, road transportation, industrial), to identify neighborhoods that are likely to cause high levels of carbon emissions and are thus mitigation priority areas. Second, we reviewed the available literature on NbS to compile and assess evidence on the effects of different NbS approaches in offsetting carbon emissions and appropriate siting options. In this step, a set of solutions for the case study area was identified. Third, we spatially deployed NbS interventions based on their expected effects and fine-scale spatial suitability. Carbon reduction benefits are an important criterion when NbS are implemented at a larger scale, but spatial variations in emissions exposure and local conditions can help to identify smaller-scale mitigation priorities. A flowchart of the model is presented in Fig. 1.

### Step 1: Spatially-explicit identification of carbon emission sources

Allocating NbS implementations to intervene with human behaviors and socio-economic dimensions requires a more accurate understanding of the locations and levels of major urban carbon emission sources. Fine-scale emissions analysis can help identify localized socio-economic drivers of emissions, which may not be apparent when looking at emissions data at a larger scale. In order to reveal carbon



**Fig. 1** Model framework for spatially explicit deployment of nature-based solutions (NbS)

emission ‘hotspots’ that require policy attention at the local level, we applied a systematic approach to identify carbon emission sources at a  $30 \times 30$  m scale, which is the finest resolution available from land cover maps, from established spatial carbon emission data.

### Transportation emissions

The literature suggests a positive correlation between pollution concentrations and road traffic (Reynolds and Broderick 2000), and annual average daily traffic (AADT) on road segments (Fu et al. 2017). As traffic volume data at the scale of Stockholm County are not available, we used road classification as a proxy for volume of traffic on its roads. In general, roads of higher classes are designed for higher traffic volume and higher speed, which result in higher emissions production and concentration. In this study, we first used binary dasymetric mapping (Mennis and Hultgren 2006) and set road = 1 and non-road = 0 to identify emissions to linear sources (road), and then adjusted the emission sources in each road segment cell by road class and population density:

$$E_t = \sum_{s=1}^S \frac{C_t \times E_s}{C_s} \quad (1)$$

$E_t$ : emission source level in target (small) zone t;  $E_s$ : emissions in origin (large) zone s;  $C_t$ : count of road cells in target zone t.  $C_t = 1$  in our case, as the target zone is the smallest scale ( $30 \times 30$  m) to which data were disaggregated;  $C_s$ : count of road cells in source zone s. This includes motorways, primary roads, and secondary roads. We calculated the value using zonal statistics in ArcGIS:

$$e_t = E_t \times W_{\text{class}} \times W_{\text{pop}} \quad (2)$$

$e_t$ : transportation emission in kt CO<sub>2</sub> eq per cell;  $W_{\text{class}}$ : weight of road classes. We used a standardized posted speed on each road segment as a proxy and normalized the values to 0–1;  $W_{\text{pop}}$ : weight of population density, normalized to 0–1.

### Residential emissions

We attempted to construct a relationship between residential carbon emissions ( $e_r$ ) and population and building density. As percentage of urbanized area is the main driver of carbon emissions (Vaccari et al. 2013), as model inputs we used the population per cell from EuroStat and the urban fabric density classifications (11 100–11 300) in the Urban Atlas land use database (European Commission 2017):

$$e_r = f(Den_{\text{pop}}, Den_{\text{building}}) \quad (3)$$

$e_r$ : residential emission in kt CO<sub>2</sub> eq per cell;  $Den_{\text{pop}}$ : population per cell;  $Den_{\text{building}}$ : building density per cell, normalized to 0–1.

We used a generic function ( $f$ ) in Eq. 3) instead of a combination of linear parameters because the relationship between urbanization and carbon emission level is often non-linear (Hong 2017; Ahmed et al. 2019). We applied a random forest algorithm to predict emission levels of sources using population counts and building density as fine-scale covariates. Similar approaches have been used by Sorichetta et al. (2015) and Stevens et al. (2015) for predicting population density from aggregated counts to small-area counts. In our model, we used 70% observations as training data and 30% as test data. The model attempts to use 500 trees and the square root of the number of columns at each split as the default parameters. We looked for a better-performing choice of parameters by using a grid search to adjust the number of trees and the number of features with the goal of reducing the root mean square error (RMSE).

### Industrial emissions

As in the method used for transport data, we applied a binary dasymetric mapping model (Mennis and Hultgren 2006) to identify emission sources from industrial and commercial complexes. To represent the industrial category, we used the land cover type 12 100 in the Urban Atlas land use database, which contains sites of industrial activities, major commercial sites, energy plants, sewage treatment plants, public, military, and private units, etc.

$$e_t = \sum_{s=1}^S \frac{C_t \times E_s}{C_s} \quad (4)$$

$e_t$ : emissions in target (small) zone t;  $E_s$ : emissions in origin (large) zone s;  $C_t$ : count of industry land use cells in target zone t.  $C_t = 1$  in our case, as the target zone was the smallest scale ( $30 \times 30$  m) to which data were disaggregated;  $C_s$ : count of industry land use cells at source zone s. We calculated the value using zonal statistics in ArcGIS.

## Step 2: Selection of suitable NBS for direct carbon sequestration and indirect carbon mitigation based on a meta-analysis

Progress in benchmarking the carbon offsetting efforts of NbS is typically constrained by insufficient synthesis of results. To better communicate the effect of NbS, we integrated the findings in existing studies that acknowledge the levels of benefits of different types of NbS toward carbon neutrality.



We applied the qualitative meta-summary techniques proposed by Sandelowski and Barroso (2007) to summarize the mechanisms proposed in the literature. Meta-summary techniques were particularly useful for our purposes, as they synthesize a combination of qualitative and quantitative research findings. Researchers have approached the subject of carbon emission mitigation through a variety of analytical methods that include statistical modeling, simulation, case studies, surveys, and historical data analysis. We used meta-analysis to (1) extract relevant statements on findings from each article; (2) reduce these statements to abstract findings that included the direction and intensity of carbon mitigation effect and the local social and economic conditions in which these methods are applied; and (3) thematize and categorize findings into key NbS strategies that could be used in our subsequent analysis.

To begin, we performed a literature search in Web of Science database on July 16, 2022, using multiple search queries combining keywords associated with NbS and carbon emission issues (see Table 1). After exploring titles, abstracts, and keywords, we limited the search to the document type “articles”, published between 2010 and 2020, and written in English.

From the list of retrieved papers (578 articles), we used different inclusion (IC) and exclusion (EC) criteria (PRISMA, 2020) to exclude studies not relevant to answer the research questions. Articles were excluded after title and abstract screening.

IC1: The paper focuses on the pathways toward zero carbon emissions.

EC1: The paper focuses on estimating the gross volumes of carbon emissions, rather than identifying the mechanisms or pathways toward carbon neutrality.

IC2: The article assesses or quantifies the efficacy of carbon emission reduction strategies.

EC2: The article proposes policy suggestions, but does not examine the quantitative relationship between carbon reduction measures and effects.

**Table 1** Combination of keywords used to design the search queries

NbS keywords	Linking boolean	Carbon emission keywords
Nature-based solutions OR NbS;	AND	Carbon emissions;
Green roofs;		Climate change;
Urban parks OR public parks;		Carbon sequestration
Street trees;		
Green infrastructure OR GI;		
Preserved habitats;		
Urban agriculture		

IC3: The article investigates carbon mitigation strategies that can be identified as NbS.

EC3: The article investigates strategies that do not fall into NbS categories (e.g., non-fossil fuel energy sources, low-carbon subsidy policies, etc.).

IC4: The article produces transferable metrics (such as elasticity, percentage change, value per unit) that could be applied to other places.

EC4: The article produces case-specific numerical values (such as total carbon emission reduction) that cannot be used to evaluate carbon reduction effects in other places.

The full text of the remaining 54 articles was reviewed against the research questions. In this selection phase, we looked for NbS interventions that involve direct mitigation (e.g., carbon sink) or indirect mitigation (e.g., interventions that could influence human behavior toward low-carbon travel), aiming to include cases representing different NbS approaches to the greatest extent possible. We also intentionally covered research conducted in multiple countries and regions of the world and included a variety of study designs. In total, 20 articles were included in the final review. Table 2 summarizes the basic characteristics of these 20 articles, which focused on cases in 10 countries, including and not restricted to Asian countries (China:  $n = 5$ , Japan:  $n = 1$ , South Korea:  $n = 2$ ), USA ( $n = 3$ ), Canada ( $n = 2$ ), European countries (Spain:  $n = 1$ , Italy:  $n = 2$ , U.K.:  $n = 1$ ). Fifteen of the 20 articles adopted a case study design where methods included primary data analysis (e.g., field survey) and secondary data analysis (e.g., GIS). The remainder ( $n = 5$ ) conducted energy model simulations or applied cross-sectional analyses based on statistical regression models.

The following information was extracted from the selected papers:

- Statements indicating the relationship between studied NbS and carbon reduction.
- The approach and indicators used to describe the impact of NbS on carbon emissions.
- Location- or environment-related variables for NbS design.

Our goal in this step was to characterize the functionality of each candidate NbS in its applied urban setting. The result was a matrix-like multi-dimensional overview of the prioritization of different types of NbS approaches based on the spatial characteristics of the study area.

### Step 3: Identify priority areas and NbS intervention types

We combined the high-emission maps produced in the first step and the NbS candidates derived from the second step

**Table 2** Basic characteristics of the articles ( $n = 20$ ) included in the review

Study	Country	Cities	Study design	Study period	
1	Anderson and Gough (2020)	Canada	Ontario	Field survey	2017
2	Chen (2015)	China	Multiple Chinese cities	Case study	2010
3	Zhao et al. (2010)	China	Hangzhou	Case study	2000–2002
4	De la Sota et al. (2019)	Spain	Lugo	Case study	40-year horizon
5	Teo et al. (2021)	Global cities		Modeling	2015
6	McPherson et al. (2011)	USA	Los Angeles,	Case study	2007–2010
7	Cai et al. (2019)	China	Wuxi	Experiment	2017–2018
8	Ismail et al. (2012)	Malaysia	Teluk Bahang	Experiment	2008
9	Kuronuma et al. (2018)	Japan		Modeling	45-year horizon
10	Russo et al. (2015)	Italy	Bolzano	Case study	2011
11	Tang et al. (2016)	China	Beijing	Case study	2012–2014
12	Jo et al. (2020)	South Korea	Seoul, Daejeon, Daegu, Chuncheon, Suncheon	Case study	2012–2019
13	Sarkar et al. (2015)	U.K	London	Survey	2005–2010
14	Lindsay et al. (2011)	New Zealand		Case study	2003–2006
15	Vaccari et al. (2013)	Italy	Florence	Case study	2006–2011
16	Escobedo et al. (2010)	USA	Gainesville and Miami-Dade	Modeling	2008
17	Jo et al. (2019)	South Korea	Seoul	Case study	2017
18	Ye et al. (2015)	China	Xiamen	Case study	2009
19	Tomalty (2012)	Canada	Ontario	Case study	2011–2031
20	Han et al. (2022)	USA	Six metropolitan counties	Case study	2006–2016

to explore the question of where each type of NbS can be implemented. We characterized land surfaces across the area based on two groups of indicators: *challenges* and *spatial suitability*, where challenges refer to emissions that every location is dealing with, while *spatial suitability* includes indicators assessing the location's land use and physical conditions for implementing certain types of NbS. Table 3 lists the spatial data used for the challenges and spatial suitability indicators. A similar classification of indicators is proposed by Kuller et al. (2019) and Sarabi et al. (2022).

We considered each NbS candidate separately. First, we produced spatial suitability maps by masking out locations where the selected NbS would not be applicable. For

example, if the selected NbS was green buildings, only the building footprint area was included in the analysis. Second, we combined the challenges and spatial suitability maps to generate neighborhoods for prioritizing the selected NbS. There are several methods for combining criteria in site selection problems, such as weighted linear combination (WLC) (Malczewski 2006), and the analytic hierarchy process (AHP) method (Parry et al. 2018). In this study, we chose thresholds for high percentile values to identify locations, where implementation was most effective in each map and combined the identified locations with equal weights. Ideally, an inclusive social-technological modeling approach would provide the option for stakeholders to define thresholds (i.e., how large an

**Table 3** Challenges and spatial suitability indicators

	Indicators	Datasets	Data source
Challenges			
Emissions	CO <sub>2</sub> emissions by sector	Emissions	Global Carbon Grid (2019)
Spatial suitability			
Urban setting	Population density	Census data	EuroStat (2018)
	Building density	Urban fabric density classifications	Urban Atlas (2018)
	Existing development	Land use data	Urban Atlas (2018)
Physical requirements	Roads	Street network	Tillväxtoch regionplaneförvaltningen (TRF), 2017
	Buildings	Building footprints	

implementation area should be considered) and choose weights for indicators from both the opportunity and challenge sides.

### Case study site and data sources

Stockholm is the political capital and commercial center of Sweden. In 2020, the estimated population of Stockholm County was 2.3 million and the population within city boundaries was 975 000. The region is expected to continue to grow, with an estimated population of 3.5 million living in Stockholm County by 2050 (TRF 2017). In order to provide infrastructure for this growth and ensure a pleasant city environment, city and county development plans emphasize increasing sustainability through protecting green and blue areas, supporting biodiversity, and improving nature recreation opportunities for residents. The climate policy framework's long-term climate goal is that by 2045 at the latest, Sweden will have zero net emissions of greenhouse gases to the atmosphere, with negative emissions thereafter. The location and land uses of the study site are shown in Fig. 2.


In our analysis, we used Global Carbon Grid data (Qin et al. 2022) from the Global Infrastructure Emissions

Database (<http://gidmodel.org.cn/>). The Global Carbon Grid provides  $0.1^\circ \times 0.1^\circ$  CO<sub>2</sub> emission maps (year 2019) for six source sectors (power, industry, residential, transport, shipping, and aviation), built upon a framework that integrates multiple data flows, including point sources, country-level sectoral activities and emissions, and transport emissions and distributions. All of these are updated on an annual basis to provide the most up-to-date global emission maps. The location-based estimates lay the foundation for building accurate high-resolution emission maps.

## RESULTS


### Spatial distribution of major carbon emission sources by sector

By identifying CO<sub>2</sub> emission sources on a  $30 \times 30$  m scale, we first identified major transportation emission sources in the study region (Fig. 3). Most of these major emission sources are in the existing urban center of Stockholm city and in Solna, due in part to high road



**Fig. 2** Maps showing (left) the location of the case study area in south-east Sweden and (right) land uses in Stockholm County and location of Stockholm city





**Fig. 3** Map of Stockholm County showing the intensity of annual road transportation carbon emissions

density and more human mobility and activities. Medium-to-high emission areas occur around major transportation corridors extending from the urban center toward emerging sub-centers, e.g., in Huddinge, Sundbyberg, Täby, and the Arlanda Airport region. High-travel speeds along highways can be a major contributing factor to the major emission sources in these areas. Rural-to-urban development also increases the inter-city car commute, which means more vehicle-kilometers traveled, more transportation energy use, and hence more emissions.

Major residential emission sources are in highly populated areas in Stockholm County, either within the urban center or scattered in peripheral areas than in sparsely populated areas (Fig. 4). This influence is moderated by building density, e.g., single-family developments in north Solna and Sundbyberg, despite lower population density, exhibit almost the same level of emissions as multi-family developments in south Stockholm City. This confirms the previous findings that low-density residential developments are more likely to introduce higher building energy emissions (Pan et al. 2019), a factor that is sometimes overlooked in residential carbon emission forecasts and in urban and environmental planning.


Industrial and commercial emission sources are usually identified from the combustion of fossil fuels for energy and heat, the use of certain products that contain

greenhouse gases, and waste handling (EPA 2020). In our maps for Stockholm County, the distribution of industrial and commercial emission sources showed less obvious patterns associated with urban development and was instead influenced by major energy consumers. Sites of construction companies, building materials suppliers, the automobile industry, and recycling centers displayed higher emissions, even in peripheral or rural areas of the county (Fig. 5).


### Scope for NbS to reduce urban carbon emissions

The mechanisms for achieving zero carbon emissions proposed in our reviewed articles fell within several key NbS strategies. We selected the following five most common types of NbS implemented in urban areas for further study: (1) green and blue infrastructure (GBI), (2) green buildings, (3) street trees, (4) urban green areas, and (5) greenbelt.

The objective of this part of the analysis was to analyze the efficacy of using NbS to mitigate carbon emissions. Developing an efficacy standard is difficult due to the vast differences in green area and structure, plant species, climate conditions, land use, and methodologies used in different studies (Aguaron and McPherson 2012; Strohbach and Haase 2012; Weissert et al. 2014). However, several patterns emerged from our literature review.



**Fig. 4** Map of Stockholm County showing the intensity of annual residential carbon emissions



**Fig. 5** Map of Stockholm County showing the intensity of annual industrial carbon emissions

- (1) The potential for trees, urban open spaces, and green facades to offset climate change impacts relates not only to the type and scale of these facilities, but also to their location (Zolch et al. 2018; Sarabi et al. 2022). For example, Vaccaci et al. (2013) found that continuous green areas have higher carbon offsetting rates than fragmented vegetation in densely built-up areas, while De la Sota et al. (2019) highlight the accrued benefits of spatially connecting green infrastructures in carbon mitigation and Ye et al. (2015) advocate locating more accessible green spaces and water bodies in residential areas.
- (2) Some NbS are more effective in mitigating carbon emissions from one sector than another. For example, green roofs can significantly reduce carbon emissions from the residential sector due to saved building energy (Kuronuma et al. 2018; Cai et al. 2019), while their capacity for capturing transportation-related carbon emissions is negligible and less studied.
- (3) Although direct comparison between different studies can be problematic, relative metrics such as carbon sequestration rate, carbon emission reduction rate, and percentage of carbon offset from total emissions can indicate the range of carbon mitigation capabilities of each NbS.

Based on Escobedo et al. (2010) and Sarabi et al. (2022), we rated the level of impact of NbS on a scale from 1 to 5 to represent the relative efficacy of NbS in carbon emission reduction, where a value of 5 indicates a relatively high potential to address the challenge and a value of 1 indicates low potential (Table 4). As the impact of each type of NbS is influenced by the social and physical conditions under which it is applied, we further characterized these conditions by urban setting, emission type, and study scale in a multi-dimensional assessment. At this point of the analysis, we roughly categorized urban settings as high-density urban core, peri-urban areas, and rural areas. More detailed urban setting characterization was performed in step 3.

### Priority areas and types of NbS interventions

Challenge and spatial suitability maps were produced in this step for each NbS, using the criteria listed in Table 5. The site selection analysis was performed on 30-m grid cells considering the spatial resolution of the utilized datasets. To facilitate the adoption of the system, we mainly used data that are easily accessible for cities (publicly accessible data from European, national, or municipal databases).

The final map for NbS prioritized areas revealed several opportunities for the inclusion of NbS in planning and policy to facilitate reductions in overall carbon emissions (Fig. 6, left panel). Increasing street trees emerged as the leading opportunity for the densely developed urban center, while GBI preservation appeared to be an effective approach throughout suburban and rural areas of Stockholm County, due to the city's forested environment and its high vegetation sequestration potential. Spatial co-location also appeared to be relevant to achieving better cost-effectiveness of urban NbS implementation, e.g., access to green space should be integrated in GBI preservation projects, as some parts of high-quality natural areas can be designed as green spaces to improve green access to residents and workers in the urban fringe.

The large-scale image for suburban Stockholm near Sundbyberg (Fig. 6, right panel) displayed how potential NbS interventions might play out locally. This area mainly includes multi-family residential use and commercial use where green buildings can be expected to achieve high environmental and energy returns. We also identified concentrated areas for street trees, due to the potential for improving the gray infrastructure environment in this medium-to-high density area. GBI interventions should primarily be applied to areas around natural reserves, to prevent future development encroaching into natural environments.


**Table 4** Nature-based solution (NbS) strategies for different spatial settings and their carbon emission mitigation effects on different sectors. Scores (1–5) represent implementation priority (1 for the lowest and 5 for the highest)

		GBI	Green buildings	Street trees	Urban green areas	Greenbelts
Urban core	Transportation	1	2	4	1	1
	Residential	1	4	3	5	2
	Industry	2	4	3	4	2
Peri-urban areas	Transportation	4	1	3	1	5
	Residential	3	3	2	3	4
	Industry	2	2	1	3	3
Rural areas	Transportation	4	1	2	1	3
	Residential	4	1	1	2	2
	Industry	3	1	1	2	2



**Table 5** Threshold values used for prioritizing nature-based solution (NbS) intervention areas

	Green–blue infrastructure	Green buildings	Street trees	Urban green areas	Greenbelts
Emissions (challenge)					
Transportation emissions			Top 50th percentile		
Residential emissions		Top 75th percentile			
Industrial emissions		Top 75th percentile			
Urban setting and demographic characters (spatial suitability)					
Population density			Top 90th percentile	Top 90th percentile	Bottom 25th percentile
Building density		Continuous urban fabric			
Physical environment (spatial suitability)					
Building rooftops		Yes			
Existing land use	Forest and semi-natural areas (shrub and/or herbaceous vegetation, open spaces with little or no vegetation)		Road transportation	Open spaces with little or no vegetation	Preserved natural resources, no growth areas, forests, wetlands



**Fig. 6** (Left) Areas in Stockholm County prioritized for different nature-based solution (NbS) interventions according to our approach, visualized with a satellite imagery background, and (right) a zoomed-in image of the Sundbyberg area in suburban Stockholm

**DISCUSSION**

Our results have three major takeaways for practice and policy-making. The first concerns the *spatial distribution of*

*proposed NbS interventions*. While NbS can be deployed throughout urban and rural areas, prioritization of NbS should cater to the specific location, e.g., in urban centers, the most welcome and efficient measures combine artificial

and natural green amenities, which includes improving access to green spaces and street trees simultaneously (Samuelsson 2021). Implementation of these measures requires policy on urban natural amenities, but also social policies such as targeting disadvantaged areas, identifying disparities, removing physical barriers (gated green spaces), partnership, and participation (Gupta et al. 2016; Biernacka and Kronenberg 2018; Kronenberg et al. 2020). For rural and suburban areas, the most important NbS measures relate to preserving and utilizing natural areas, which include greenbelt, grass strips, and other forms of green infrastructure. Natural features, including urban forests, transitional ecosystems, and local natural reserves, can provide ecosystem services for urban areas, save infrastructure costs, and prevent urban sprawl (Gavrilidis et al. 2019).

The second practical takeaway from our findings is that *NbS measures can benefit more than one carbon emissions sector (transportation, residential, industry) and generate co-benefits at places where multiple NbS interventions are prioritized.* Our spatially detailed analysis revealed the potential for the co-existence of multiple NbS approaches. Urban centers and sub-centers with high road density and high population density can benefit from approaches ranging from green infrastructures to street environments, quality-of-life amenities, accessible recreational opportunities, and green roofs. Implementation of these NbS interventions would generate co-benefits for the residential, transportation, and industrial sectors.

The third practical takeaway concerns *the behavioral influences of implementing NbS.* In this study, we assessed both direct and indirect pathways for NbS to achieve carbon savings goals. Direct effects such as preserving green spaces are widely reported in the literature (Li and Wang 2021; Page et al. 2021; Ramchunder and Ziegler 2021), but using NbS as a potential tool for nudging and influencing human behaviors is a significant, but often overlooked, opportunity in policy-making to achieve carbon neutrality (Linder et al. 2022). For example, improving streetscapes in the urban center not only contributes to transportation emission reduction but is also beneficial for fostering a walking and biking culture and building an environment-aware civil society (Ewing et al. 2015; Cain et al. 2017; Liu et al. 2020). The socio-psychological aspects of change were highlighted by Seyfang and Haxeltine (2012), aiming at a shift in the behavioral norms and societal shifts in values and beliefs (Westley et al. 2011). We thus see an urgent need to include behavioral and resilience building interventions, such as green education, new green jobs, participatory place-based learning and experimentation, and inviting civic organizations into NbS management, in a wider urban NbS toolbox to achieve urban carbon neutrality.

NbS has many other benefits beyond climate actions, such as enhancing habitat quality, promoting recreational

opportunities and human health, increasing equitable access to amenities, and creating jobs in the green sector (Raymond et al. 2017; Giordano et al. 2020; Ruangpan et al. 2021). Due to space restrictions, we focused our investigation on carbon emission effects, but future studies should acknowledge the multifunctionality and co-benefits of NbS in all environmental and socio-ecological dimensions.

## CONCLUSIONS

This paper presents a novel systematic framework for identifying the need for NbS interventions at the local scale and for prioritizing sites and strategies for NbS interventions. By moving from the global to the neighborhood level, we identified NbS strategies that can maximize carbon emissions saving potential given local conditions, and sites where they should be implemented. In our case study, the city center and surrounding areas of Stockholm were shown to require different types of NbS to maximize the carbon-saving benefits needed to transform Stockholm into a carbon-neutral city. In the urban center, the most welcome and efficient measures combine improving access to green spaces and streetscapes. In rural and suburban areas, the most important measures related to preserving and utilizing natural areas, which include greenbelt and green infrastructure. Natural features, including urban forests, transitional ecosystems, and local nature reserves, can provide ecosystem services to urban areas, saving infrastructure costs and preventing urban sprawl.

The framework developed in this study can be applied to address a variety of urban sustainability and resilience challenges, such as heat stress, stormwater management, air quality, landscape connectivity, and spatial equity, each of which will require a specific set of mitigating approaches. In a typical application, one could first rank neighborhoods in terms of the level of exposure and/or vulnerability to a set of challenges, then prioritize hazard-mitigating approaches based on their benefits and impacts, and finally, conduct rule-based suitability analysis to overlay social criteria and physical conditions and generate spatially detailed maps. Our proposed method can help planners better select target locations for intended risk/hazard-mitigating interventions.

One methodological limitation of this study was the lack of empirical data and such analysis can be enhanced as further evidence becomes available. Land cover type data from Urban Atlas are useful but could be complemented with additional socio-economic variables to better account for local-scale emissions. Further, our analysis only captured the physical environment where NbS can be applied, so in future studies implementation of NbS should be



coordinated with other environmental and social benefits across social groups.

Another limitation is that our study did not focus on iterating solutions for a more collaborative spatial planning process. While our methods provide spatially detailed solutions for ensuring that NbS actions align with existing and/or proposed urban planning strategies and governance processes, further studies can integrate the analysis into a collaborative planning process to facilitate greater stakeholder inputs in terms of identifying suitable intervention areas and solutions. Practitioners need to integrate diverse types and systems of knowledge and values for NbS design and implementation to ensure that plans are socially comprehensible and acceptable.

**Acknowledgements** The work is supported by the National Natural Science Foundation of China [NSFC 52000130], Swedish funding agency [Formas Grant 2021-00293] and [Formas Grant 2021-00416], the program FAIRTRANS financed by Formas and Mistra [DIA 2019/28], and the program FUTURE PROOF CITIES financed by KK-Stiftelsen.

**Author contributions** CC: Data curation, Methodology, Software, Visualization, and Investigation; HP: Writing—Original draft preparation, Conceptualization, and Validation. JP: Data curation and Methodology; SB: Supervision, Writing—Reviewing and Editing; ZK: Writing—Reviewing and Editing, Conceptualization, and Supervision.

**Funding** Open access funding provided by Royal Institute of Technology.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## REFERENCES

- Aguaron, E., and E.G. McPherson. 2012. Comparison of methods for estimating carbon dioxide storage by Sacramento's urban forest. In *Carbon sequestration in urban ecosystems*, ed. R. Lal and B. Augustin, 43–71. The Netherlands: Springer.
- Ahmed, Z., Z. Wang, and S. Ali. 2019. Investigating the non-linear relationship between urbanization and CO<sub>2</sub> emissions: An empirical analysis. *Air Quality Atmosphere and Health* 12: 945–953.
- Albert, C., B. Schröter, D. Haase, M. Brillinger, J. Henze, S. Herrmann, and B. Matzdorf. 2019. Addressing societal challenges through nature-based solutions: How can landscape planning and governance research contribute? *Landscape and Urban Planning* 182: 12–21.
- Anderson, V., and W. Gough. 2020. Evaluating the potential of nature-based solutions to reduce ozone, nitrogen dioxide, and carbon dioxide through a multi-type green infrastructure study in Ontario. *Canada. City and Environment Interactions* 6: 100043.
- Andersson, E., D. Haase, S. Scheuer, and T. Wellmann. 2020. Neighbourhood character affects the spatial extent and magnitude of the functional footprint of urban green infrastructure. *Landscape Ecology* 35: 1605–1618.
- Biernacka, M., and J. Kronenberg. 2018. Classification of institutional barriers affecting the availability, accessibility and attractiveness of urban green spaces. *Urban Forestry & Urban Greening* 36: 22–33.
- Bierwagen, B.G., D.M. Theobald, C.R. Pyke, A. Choate, P. Groth, J.V. Thomas, and P. Morefield. 2010. National housing and impervious surface scenarios for integrated climate impact assessments. *Proceedings of the National Academy of Sciences* 107: 20887–20892.
- Bradfer-Lawrence, T., T. Finch, R.B. Bradbury, G.M. Buchanan, A. Millgley, and R.H. Field. 2021. The potential contribution of terrestrial nature-based solutions to a national 'net zero' climate target. *Journal of Applied Ecology* 58: 2349–2360.
- Cai, L., X.P. Feng, J.Y. Yu, Q.C. Xiang, and R. Chen. 2019. Reduction in carbon dioxide emission and energy savings obtained by using a green roof. *Aerosol Air Quality Research* 19: 2432–2445.
- Cain, K.L., K.A. Gavand, T.L. Conway, C.M. Geremia, R.A. Millstein, L.D. Frank, B.E. Saelens, M.A. Adams, et al. 2017. Developing and validating an abbreviated version of the Microscale Audit for Pedestrian Streetscapes (MAPS-Abbreviated). *Journal of Transport & Health* 5: 84–96.
- Chen, W.Y. 2015. The role of urban green infrastructure in offsetting carbon emissions in 35 major Chinese cities: A nationwide estimate. *Cities* 44: 112–120.
- Choi, C., P. Berry, and A. Smith. 2021. The climate benefits, co-benefits, and trade-offs of green infrastructure: A systematic literature review. *Journal of Environmental Management* 291: 112583.
- Connop, S., P. Vandergert, B. Eisenberg, M. Collier, C. Nash, J. Clough, and D. Newport. 2016. Renaturing cities using a regionally-focused biodiversity-led multifunctional benefits approach to urban green infrastructure. *Environmental Science and Policy* 62: 99–111.
- Cortinovis, C., and D. Geneletti. 2018. Ecosystem services in urban plans: What is there, and what is still needed for better decisions. *Land Use Policy* 70: 298–312.
- Cortinovis, C., P. Olsson, N. Boke-Olén, and K. Hedlund. 2022. Scaling up nature-based solutions for climate-change adaptation: Potential and benefits in three European cities. *Urban Forestry & Urban Greening* 67: 127450.
- De la Sota, C., V.J. Ruffato-Ferreira, L. Ruiz-García, and S. Alvarez. 2019. Urban green infrastructure as a strategy of climate change mitigation. A case study in northern Spain. *Urban Forestry & Urban Greening* 40: 145–151.
- Dorst, H., A. van der Jagt, R. Raven, and H. Runhaar. 2019. Urban greening through nature-based solutions—Key characteristics of an emerging concept. *Sustainable Cities and Society* 49: 101620.
- EPA. 2020. Sources of Greenhouse Gas Emissions. <https://www.epa.gov/ghgemissions/sources-greenhouse-gas-emissions>. Accessed 29 Dec 2022.
- Escobedo, F., S. Varela, M. Zhao, J.E. Wagner, and W. Zipperer. 2010. Analyzing the efficacy of subtropical urban forests in offsetting carbon emissions from cities. *Environmental Science & Policy* 13: 362–372.

- European Commission. 2017. Eurostat population database; <https://ec.europa.eu/eurostat/web/main/data/database>. Accessed 11 May 2022.
- European Environment Agency. 2021. Nature-based solutions in Europe: Policy, knowledge and practice for climate change adaptation and disaster risk reduction. <https://op.europa.eu/en/publication-detail/-/publication/da65d478-a24d-11eb-b85c-01aa75ed71a1/language-en/format-PDF/source-210308869>. Accessed 29 Dec 2022.
- Ewing, R., A. Hajrasouliha, K.M. Neckerman, M. Purciel-Hill, and W. Greene. 2015. Streetscape features related to pedestrian activity. *Journal of Planning Education and Research* 36: 5–15.
- Fastenrath, S., J. Bush, and L. Coenen. 2020. Scaling-up nature-based solutions. Lessons from the Living Melbourne strategy. *Geoforum* 116: 63–72.
- Ferreira, A., R. Guilherme, C. Ferreira, and M. Oliveira. 2018. Urban agriculture, a tool towards more resilient urban communities? *Current Opinion in Environmental Science & Health* 5: 93–97.
- Fu, M., J. Kelly, and J. Clinch. 2017. Estimating annual average daily traffic and transport emissions for a national road network: A bottom-up methodology for both nationally-aggregated and spatially-disaggregated results. *Journal of Transport Geography* 58: 186–195.
- Gavrilidis, A.A., M.R. Nişã, D.A. Onose, D.L. Badiu, and I.I. Nãstase. 2019. Methodological framework for urban sprawl control through sustainable planning of urban green infrastructure. *Ecological Indicators* 96: 67–78.
- Geertman, S., and J. Stillwell. 2003. Interactive support systems for participatory planning. In *Planning support systems in practice. Advances in spatial science*, ed. S. Geertman and J. Stillwell. Berlin: Springer.
- Gill, Z.M., M.J. Tierney, I.M. Pegg, and N. Allan. 2010. Low-energy dwellings: The contribution of behaviors to actual performance. *Building Research & Information* 38: 491–508.
- Giordano, R., I. Pluchinotta, A. Pagano, A. Scricciu, and F. Nanu. 2020. Enhancing nature-based solutions acceptance through stakeholders' engagement in co-benefits identification and trade-offs analysis. *Science of the Total Environment* 713: 136552.
- Goldenberg, R., Z. Kalantari, V. Cvetkovic, U. Mörberg, B. Deal, and G. Destouni. 2017. Distinction, quantification and mapping of potential and realized supply-demand of flow-dependent ecosystem services. *Science of the Total Environment* 593–594: 599–609.
- Gómez Martín, E., M. Máñez Costa, and K. Schwerdtner Máñez. 2020. An operationalized classification of Nature Based Solutions for water-related hazards: From theory to practice. *Ecological Economics* 167: 106460.
- Gou, Z., and S.S.-Y. Lau. 2014. Contextualizing green building rating systems: Case study of Hong Kong. *Habitat International* 44: 282–289.
- Gromke, C., and B. Blocken. 2015. Influence of avenue-trees on air quality at the urban neighborhood scale. Part II: Traffic pollutant concentrations at pedestrian level. *Environmental Pollution* 196: 176–184.
- Gupta, K., A. Roy, K. Luthra, and S. Maithani. 2016. GIS based analysis for assessing the accessibility at hierarchical levels of urban green spaces. *Urban Forestry & Urban Greening* 18: 198–211.
- Han, A., T. Daniels, and C. Kim. 2022. Managing urban growth in the wake of climate change: Revisiting greenbelt policy in the US. *Land Use Policy* 112: 105867.
- Hong, J. 2017. Non-linear influences of the built environment on transportation emissions: Focusing on densities. *Journal of Transport and Land Use* 10: 229–240.
- Ismail, A., M. Samad, A. Rahman, and F. Yeok. 2012. Cooling potentials and CO<sub>2</sub> uptake of Ipomoea Pes-caprae installed on the flat roof of a single storey residential building in Malaysia. *Procedia—Social and Behavioral Sciences* 35: 361–368.
- Jo, H.-K., J.-Y. Kim, and H.-M. Park. 2019. Carbon reduction and planning strategies for urban parks in Seoul. *Urban Forestry & Urban Greening* 41: 48–54.
- Jo, H.-K., J.-Y. Kim, and H.-M. Park. 2020. Carbon and PM<sub>2.5</sub> reduction and design guidelines for street trees in Korea. *Sustainability* 12: 10414.
- Kabisch, N., N. Frantzeskaki, S. Pauleit, S. Naumann, M. Davis, M. Artmann, D. Haase, S. Knapp, et al. 2016. Nature-based solutions to climate change mitigation and adaptation in urban areas: perspectives on indicators, knowledge gaps, barriers, and opportunities for action. *Ecology and Society* 21.
- Kalantari, Z., C.S. Santos Ferreira, J. Page, R. Goldenberg, J. Olsson, and G. Destouni. 2019. Meeting sustainable development challenges in growing cities: Coupled social-ecological systems modeling of land use and water changes. *Journal of Environmental Management* 245: 471–480.
- Keesstra, S., J. Nunes, A. Novara, D. Finger, D. Avelar, Z. Kalantari, and A. Cerdà. 2018. The superior effect of nature based solutions in land management for enhancing ecosystem services. *Science of the Total Environment* 610–611: 997–1009.
- Koo, B.W., S. Guhathakurta, and N. Botchwey. 2022. How are neighborhood and street-level walkability factors associated with walking behaviors? A big data approach using street view images. *Environment and Behavior* 54: 211–241.
- Krauze, K., and I. Wagner. 2019. From classical water-ecosystem theories to nature-based solutions—Contextualizing nature-based solutions for sustainable city. *Science of the Total Environment* 655: 697–706.
- Kronenberg, J., A. Haase, E. Łaszkiwicz, A. Antal, A. Baravikova, M. Biernacka, D. Dushkova, R. Filčak, et al. 2020. Environmental justice in the context of urban green space availability, accessibility, and attractiveness in postsocialist cities. *Cities* 106: 102862.
- Kuller, M., P.M. Bach, S. Roberts, D. Browne, and A. Deletic. 2019. A planning-support tool for spatial suitability assessment of green urban stormwater infrastructure. *Science of the Total Environment* 686: 856–868.
- Kuronuma, T., H. Watanabe, T. Ishihara, D. Kou, K. Tushima, M. Ando, and S. Shindo. 2018. CO<sub>2</sub> payoff of extensive green roofs with different vegetation species. *Sustainability* 10: 2256.
- Laforteza, R., J. van den Chen, C. Bosch, and T. Randrup. 2018. Nature-based solutions for resilient landscapes and cities. *Environmental Research* 165: 431–441.
- Lanau, M., L. Herbert, and G. Liu. 2021. Extending urban stocks and flows analysis to urban greenhouse gas emission accounting: A case of Odense, Denmark. *Journal of Industrial Ecology* 25: 961–978.
- Li, P., and Z.H. Wang. 2021. Environmental co-benefits of urban greening for mitigating heat and carbon emissions. *Journal of Environmental Management* 293: 112963.
- Linder, N., M. Giusti, K. Samuelsson, and S. Barthel. 2022. Pro-environmental habits: An underexplored research agenda in sustainability science. *Ambio* 51: 546–556. <https://doi.org/10.1007/s13280-021-01619-6>.
- Lindsay, G., A. Macmillan, and A. Woodward. 2011. Moving urban trips from cars to bicycles: Impact on health and emissions. *Australian and New Zealand Journal of Public Health* 35: 54.
- Liu, Y., D. Yang, H.J.P. Timmermans, and B. de Vries. 2020. Analysis of the impact of street-scale built environment design near metro stations on pedestrian and cyclist road segment choice: A stated choice experiment. *Journal of Transport Geography* 82: 102570.

- Madureira, H., and T. Andresen. 2014. Planning for multifunctional urban green infrastructures: Promises and challenges. *Urban Design International* 19: 38–49.
- Malczewski, J. 2006. A GIS-based multi-criteria decision analysis: A survey of the literature. *International Journal of Geographical Information Science* 20: 703–726.
- McPherson, E.G. 1998. Atmospheric carbon dioxide reduction by Sacramento's urban forest. *Journal of Arboriculture* 24: 215–223.
- McPherson, E.G., J. Simpson, Q. Xiao, and C. Wu. 2011. Million trees Los Angeles canopy cover and benefit assessment. *Landscape and Urban Planning* 99: 40–50.
- Meerow, S., and J. Newell. 2017. Spatial planning for multifunctional green infrastructure: Growing resilience in Detroit. *Landscape and Urban Planning* 159: 62–75.
- Mennis, J., and T. Hultgren. 2006. Intelligent dasymetric mapping and its application to areal interpolation. *Cartography and Geographical Information Science* 33: 179–194.
- Midgley, S., K. Esler, P. Holden, A. Rebelo, S. Stuart-Hill, J. Cullis, and N. Methner. 2021. Typologies of collaborative governance for scalable nature-based solutions in two strategic South African river systems. *Ambio* 50: 1587–1609. <https://doi.org/10.1007/s13280-021-01531-z>.
- Nesshöver, C., T. Assmuth, K.N. Irvine, G.M. Rusch, K.A. Waylen, B. Delbaere, D. Haase, L. Jones-Walters, et al. 2017. The science, policy and practice of nature-based solutions: An interdisciplinary perspective. *Science of the Total Environment* 579: 1215–1227.
- Norton, B., A. Coutts, S. Livesley, R. Harris, A. Hunter, and N. Williams. 2015. Planning for cooler cities: A framework to prioritise green infrastructure to mitigate high temperatures in urban landscapes. *Landscape and Urban Planning* 134: 127–138.
- Nowak, D., and D. Crane. 2002. Carbon storage and sequestration by urban trees in the USA. *Environmental Pollution* 116: 381–389.
- Page, J., E. Käresdotter, G. Destouni, H. Pan, and Z. Kalantari. 2021. A more complete accounting of greenhouse gas emissions and sequestration in urban landscapes. *Anthropocene* 34: 100296.
- Pan, H., J. Page, C. Cong, S. Barthel, and Z. Kalantari. 2021. How ecosystems services drive urban growth: Integrating nature-based solutions. *Anthropocene* 35: 100297.
- Pan, H., J. Page, L. Zhang, S. Chen, C. Cong, G. Destouni, Z. Kalantari, and B. Deal. 2019. Using comparative socio-ecological modeling to support Climate Action Planning (CAP). *Journal of Cleaner Production* 232: 30–42.
- Pan, H., J. Page, L. Zhang, C. Cong, C. Ferreira, E. Jonsson, H. Näsström, G. Destouni, et al. 2020. Understanding interactions between urban development policies and GHG emissions: A case study in Stockholm Region. *Ambio* 49: 1313–1327. <https://doi.org/10.1007/s13280-019-01290-y>.
- Parry, J., S. Ganaie, and M. Bhat. 2018. GIS based land suitability analysis using AHP model for urban services planning in Srinagar and Jammu urban centers of J&K, India. *Journal of Urban Management* 7: 46–56.
- Pataki, D., M. Alberti, M. Cadenasso, A. Felson, M. McDonnell, S. Pincetl, R.V. Pouyat, and H. Setälä. 2021. The benefits and limits of urban tree planting for environmental and human health. *Frontiers in Ecology and Evolution* 9: 155.
- Qin, X., D. Tong, F. Liu, R. Wu, B. Zheng, Y. Zheng, J. Liu, R. Xu, et al. 2022. Global and regional drivers of power plant CO<sub>2</sub> emissions over the last three decades revealed from unit-based database. *Earth's Future* 10: e2022EF002657.
- Ramchunder, S.J., and A.D. Ziegler. 2021. Promoting sustainability education through hands-on approaches: A tree carbon sequestration exercise in a Singapore green space. *Sustainability Science* 16: 1045–1059.
- Raymond, C.M., N. Frantzeskaki, N. Kabisch, P. Berry, M. Breil, M.R. Nita, D. Geneletti, and C. Calfapietra. 2017. A framework for assessing and implementing the co-benefits of nature-based solutions in urban areas. *Environmental Science & Policy* 77: 15–24.
- Ren, Z., H. Zheng, X. He, D. Zhang, G. Shen, and C. Zhai. 2019. Changes in spatio-temporal patterns of urban forest and its above-ground carbon storage: Implication for urban CO<sub>2</sub> emissions mitigation under China's rapid urban expansion and greening. *Environment International* 129: 438–450.
- Reynolds, W., and B. Broderick. 2000. Development of an emissions inventory model for mobile sources. *Transportation Research Part d: Transport and Environment* 5: 77–101.
- Ruangpan, L., Z. Vojinovic, J. Plavšić, D.-J. Doong, T. Bahlmann, A. Alves, L.-H. Tseng, A. Randelović, et al. 2021. Incorporating stakeholders' preferences into a multi-criteria framework for planning large-scale Nature-Based Solutions. *Ambio* 50: 1514–1531. <https://doi.org/10.1007/s13280-020-01419-4>.
- Russo, A., F. Escobedo, N. Timilsina, and S. Zerbe. 2015. Transportation carbon dioxide emission offsets by public urban trees: A case study in Bolzano, Italy. *Urban Forestry and Urban Greening* 14: 398–403.
- Samuelsson, K. 2021. The topodiverse City: Urban form for subjective well-being. *Frontiers in Built Environment* 7: 735221.
- Sandelowski, M., and J. Barroso. 2007. *Handbook for synthesizing qualitative research*. New York: Springer.
- Sarabi, S., Q. Han, B. Vries, and A. Romme. 2022. The nature-based solutions planning support system: A playground for site and solution prioritization. *Sustainable Cities and Society* 78: 103608.
- Sarkar, C., C. Webster, M. Pryor, D. Tang, S. Melbourne, X. Zhang, and L. Jianzheng. 2015. Exploring associations between urban green, street design and walking: Results from the greater London boroughs. *Landscape and Urban Planning* 143: 112–125.
- Seyfang, G., and A. Haxeltine. 2012. Growing grassroots innovations: Exploring the role of community-based initiatives in governing sustainable energy transitions. *Environment and Planning C: Government and Policy* 30: 381–400.
- Sha, Z., Y. Bai, R. Li, H. Lan, X. Zhang, J. Li, X. Liu, S. Chang, et al. 2022. The global carbon sink potential of terrestrial vegetation can be increased substantially by optimal land management. *Communication Earth and Environment* 3: 8.
- Smardon, R.C. 1988. Perception and aesthetics of the urban environment: Review of the role of vegetation. *Landscape and Urban Planning* 15: 85–106.
- Song, M., Y. Zhao, J. Liang, and F. Li. 2023. Spatial-temporal variability of carbon emission and sequestration and coupling coordination degree in Beijing district territory. *Cleaner Environmental Systems* 8: 100102.
- Sorichetta, A., G.M. Hornby, F.R. Stevens, A.E. Gaughan, C. Linard, and A.J. Tatem. 2015. High-resolution gridded population datasets for Latin America and the Caribbean in 2010, 2015, and 2020. *Scientific Data* 2: 150045.
- Stevens, F.R., A.E. Gaughan, C. Linard, and A.J. Tatem. 2015. Disaggregating census data for population mapping using random forests with remotely-sensed and ancillary data. *PLoS ONE* 10: e0107042.
- Strohbach, M.W., and D. Haase. 2012. Above-ground carbon storage by urban trees in Leipzig, Germany: Analysis of patterns in a European city. *Landscape and Urban Planning* 104: 95–104.
- Tang, Y., A. Chen, and S. Zhao. 2016. Carbon storage and sequestration of urban street trees in Beijing, China. *Frontiers in Ecology and Evolution* 4: 53.
- Teo, H., Y. Zeng, T. Sarira, T. Fung, Q. Zheng, X. Song, K. Chong, and L. Koh. 2021. Global urban reforestation can be an

- important natural climate solution. *Environmental Research Letters* 16: 034059.
- Tomalty, R. 2012. Carbon in the Bank—Ontario’s Greenbelt and Its Role in Mitigating Climate Change. Vancouver: David Suzuki Foundation. <https://david Suzuki.org/wp-content/uploads/2012/08/carbon-bank-ontario-greenbelt-role-mitigating-climate-change.pdf>. Accessed 29 Dec 2022.
- TRF—Tillväxt-och Regionplaneförvaltningen. 2017. Regional Utvecklingsplan FÖr Stockholmsregionen (RUFs) 2050. <http://www.rufs.se/publikationer/2018/rufs-2050/>
- Tsoka, S., T. Leduc, and A. Rodler. 2021. Assessing the effects of urban street trees on building cooling energy needs: The role of foliage density and planting pattern. *Sustainable Cities and Society* 65: 102633.
- Vaccari, F., B. Gioli, P. Toscano, and C. Perrone. 2013. Carbon dioxide balance assessment of the city of Florence (Italy), and implications for urban planning. *Landscape and Urban Planning* 120: 138–146.
- Venter, Z.S., D.N. Barton, L. Martinez-Izquierdo, J. Langemeyer, F. Baró, and T. McPhearson. 2021. Interactive spatial planning of urban green infrastructure—Retrofitting green roofs where ecosystem services are most needed in Oslo. *Ecosystem Services* 50: 101314.
- Wamsler, C., B. Wickenberg, H. Hanson, J. Alkan Olsson, S. Stålhammar, H. Björn, H. Falck, D. Gerell, et al. 2020. Environmental and climate policy integration: Targeted strategies for overcoming barriers to nature-based solutions and climate change adaptation. *Journal of Cleaner Production* 247: 119154.
- Weissert, L.F., J.A. Salmond, and L. Schwendenmann. 2014. A review of the current progress in quantifying the potential of urban forests to mitigate urban CO<sub>2</sub> emissions. *Urban Climate* 8: 100–125.
- Westley, F., P. Olsson, C. Folke, T. Homer-Dixon, H. Vredenburg, D. Loorbach, J. Thompson, M. Nilsson, et al. 2011. Tipping toward sustainability: Emerging pathways of transformation. *Ambio* 40: 762–780. <https://doi.org/10.1007/s13280-011-0186-9>.
- Xie, X., Y. Lu, and Z. Gou. 2017. Green building pro-environment behaviors: Are green users also green buyers? *Sustainability* 9: 1703.
- Ye, H., X. He, Y. Song, X. Li, G. Zhang, T. Lin, and L. Xiao. 2015. A sustainable urban form: The challenges of compactness from the viewpoint of energy consumption and carbon emission. *Energy and Buildings* 93: 90–98.
- Zhao, M., Z.H. Kong, F.J. Escobedo, and J. Gao. 2010. Impacts of urban forests on offsetting carbon emissions from industrial energy use in Hangzhou, China. *Journal of Environmental Management* 91: 807–813.
- Zolch, T., C. Wamsler, and S. Pauleit. 2018. Integrating the ecosystem-based approach into municipal climate adaptation strategies: The case of Germany. *Journal of Cleaner Production* 170: 966–977.

**Publisher’s Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## AUTHOR BIOGRAPHIES

**Cong Cong** is a Lecturer at MIT Department of Urban Studies and Planning, Massachusetts Institute of Technology, USA. Her research interests include computer science and urban planning.

*Address:* MIT Department of Urban Studies and Planning, Massachusetts Institute of Technology, Cambridge, MA, USA.

*Address:* Department of Urban and Regional Planning, University of Illinois at Urbana-Champaign, Champaign, IL, USA.

e-mail: ccong2@mit.edu; ccong2@illinois.edu

**Haozhi Pan** (✉) is an Associate Professor at School of International and Public Affairs; China Institute for Urban Governance, Shanghai Jiao Tong University, China. His research interests include urban sustainability, urban governance, and regional economics.

*Address:* School of International and Public Affairs, China Institute for Urban Governance, Shanghai Jiao Tong University, 1954 Huashan Rd., Shanghai 200030, China.

e-mail: panhaozhi@sjtu.edu.cn

**Jessica Page** is a Ph.D. candidate at Department of Physical Geography, Stockholm University, Sweden. Her research interests include Regional and global hydrology and water resources issues and Sustainable urban and rural development and adaptive land-use planning for decision support.

*Address:* Department of Physical Geography, Stockholm University, Stockholm, Sweden.

e-mail: jessica.page@natgeo.su.se

**Stephan Barthel** is a Professor at the department of Building Engineering, Energy Systems and Sustainability Science, University of Gävle; Stockholm Resilience Centre, Stockholm University. His research interests include environmental issues in metropolitan landscapes.

*Address:* Department of Building Engineering, Energy Systems and Sustainability Science, University of Gävle, Gävle, Sweden.

*Address:* Stockholm Resilience Centre, Stockholm University, Stockholm, Sweden.

e-mail: stephan.barthel@hig.se

**Zahra Kalantari** (✉) is an Associate Professor at Department of Sustainable Development, Environmental Science and Engineering (SEED), KTH Royal Institute of Technology, Stockholm, Sweden. Her research interests include Regional and global hydrology and water resources issues and Sustainable urban and rural development and adaptive land-use planning for decision support.

*Address:* Department of Sustainable Development, Environmental Science and Engineering (SEED), KTH Royal Institute of Technology, Stockholm, Sweden.

e-mail: zahrak@kth.se; zahra.kalantari@natgeo.su.se