ERRATUM

Erratum to: Efficacy of Additional Canagliflozin Administration to Type 2 Diabetes Patients Receiving Insulin Therapy: Examination of Diurnal Glycemic Patterns Using Continuous Glucose Monitoring (CGM)

Mihoko Matsumura · Yuki Nakatani · Seiichi Tanka ·

Chie Aoki · Masaaki Sagara · Kazunori Yanagi · Kunihiro Suzuki ·

Yoshimasa Aso

Published online: October 23, 2017

© The Author(s) 2017. This article is an open access publication

Erratum to: Diabetes Ther (2017) 8:821–827
DOI 10.1007/s13300-017-0274-3

Unfortunately, the original publication of this paper contained errors in the presentation of Fig. 2. The corrected Fig. 2 is given below.

Fig. 2 The total amount of insulin in all patients was divided into basal and bolus regimens to show changes from days 1 to 7. The number of basal insulin units decreased from 128 to 76 and that of bolus insulin from 266 to 154

The online version of the original article can be found under doi:[10.1007/s13300-017-0274-3](https://doi.org/10.1007/s13300-017-0274-3).

M. Matsumura (✉) · Y. Nakatani · S. Tanka ·
C. Aoki · M. Sagara · K. Yanagi · K. Suzuki · Y. Aso
Department of Endocrinology and Metabolism,
Dokkyo Medical University, Tochigi, Japan
e-mail: m-mihoko@dokkyomed.ac.jp

Y. Nakatani
Department of Diabetes and Endocrinology, Nikko
Medical Center, Dokkyo Medical University,
Tochigi, Japan

Open Access. This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits any non-commercial use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.