

CORRECTION

## Correction to: Bedtime-to-Morning Glucose Difference and iGlarLixi in Type 2 Diabetes: Post Hoc Analysis of LixiLan-L

Ariel Zisman · Terry Dex · Michelle Roberts · Aramesh Saremi ·  
Jason Chao · Vanita R. Aroda

Published online: December 4, 2018  
© The Author(s) 2018

Correction to: Diabetes Ther (2018) 9:2155–2162  
<https://doi.org/10.1007/s13300-018-0507-0>

In the original publication, the text in abstract section under the ‘Results’ section is incorrectly published as ‘higher proportion of patients reached a BeAM value < 55 mg/dL. The correct text should read as ‘higher proportion of patients reached a BeAM value < 50 mg/dL’.

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits any non-commercial use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

---

The original article can be found online at <https://doi.org/10.1007/s13300-018-0507-0>.

---

A. Zisman (✉)  
Endocrine Center of Aventura, Miami, FL, USA  
e-mail: [azisman@myendocrine.com](mailto:azisman@myendocrine.com)

T. Dex · M. Roberts · A. Saremi  
Sanofi US, Inc, Bridgewater, NJ, USA

J. Chao  
Xinyi, Inc, Bridgewater, NJ, USA

V. R. Aroda  
Brigham and Women’s Hospital, Harvard Medical  
School, Boston, MA, USA