CORRECTION

Correction to: The Association Between Phosphorylated Neurofilament Heavy Chain (pNF-H) and Small Fiber Neuropathy (SFN) in Patients with Impaired Glucose Tolerance

Yu-peng Li · Zhong-qing Yan · Li-ping Han · Ai-li Yin ·
Jin-yong Xu · Ya-ran Zhai · Sai Hao · Lin Zhang · Yun Xie

Published online: December 21, 2019
© The Author(s) 2019

Correction to:
Diabetes Ther
<https://doi.org/10.1007/s13300-019-00716-w>

In the original article, there was some error in Table 2. The correct Table 2 is given below.

The original article can be found online at <https://doi.org/10.1007/s13300-019-00716-w>.

Y. Li (✉) · Z. Yan · L. Han · A. Yin · J. Xu · Y. Zhai ·
S. Hao · L. Zhang · Y. Xie (✉)
Tianjin Medical University Metabolic Diseases
Hospital & Tianjin Medical University Chu Hsien-I
Memorial Hospital, Tianjin, China
e-mail: dr_liyupeng@126.com Y. Xie
e-mail: xieyuntj@126.com

Y. Li · Z. Yan · L. Han · A. Yin · J. Xu · Y. Zhai ·
S. Hao · L. Zhang · Y. Xie
Tianjin Institute of Endocrinology, Tianjin, China

Y. Li · Z. Yan · L. Han · A. Yin · J. Xu · Y. Zhai ·
S. Hao · L. Zhang · Y. Xie
NHC Key Laboratory of Hormones and
Development (Tianjin Medical University), Tianjin
Key Laboratory of Metabolic Diseases, Tianjin,
China

Table 2 Basic demographical and clinical parameters among three groups

	CON (<i>N</i> = 33)	IGT (<i>N</i> = 61) (exclude 17 cases large fiber neuropathy)		<i>p</i> value
		Non-SFN control	SFN	
<i>N</i>	33	20	24	
M/F	10/23	9/11	11/23	0.521
Age (years)	50.5 ± 8.7	53.6 ± 7.8	57.7 ± 8.8*	0.010
BMI (kg/m ²)	23.9 ± 2.7	25.22 ± 2.53	25.13 ± 3.49	0.175
WHR	0.87 ± 0.07	0.9 ± 0.07	0.9 ± 0.08	0.125
HbA1c (%)	5.6 ± 0.3	5.98 ± 0.39*	5.93 ± 0.39*	0.002
FPG (mmol/l)	5.02 ± 0.32	5.92 ± 0.53*	5.89 ± 0.53*	0.000
2 hPG (mmol/l)	6.02 ± 1.11	8.87 ± 0.75*	9.52 ± 0.92*,#	0.000
TG (mmol/l)	1.52 ± 0.57	1.62 ± 0.45	1.60 ± 0.99	0.881
TC (mmol/l)	5.59 ± 0.73	5.15 ± 1.01	5.00 ± 0.99	0.070
HDL-c (mmol/l)	1.58 ± 0.45	1.51 ± 0.35	1.39 ± 0.31	0.230
LDL-c (mmol/l)	3.56 ± 0.66	3.11 ± 0.88	3.10 ± 0.96	0.134
Cr (mmol/l)	71.47 ± 15.25	69.25 ± 10.90	70.24 ± 17.3	0.883
BUN (mmol/l)	5.04 ± 1.49	5.56 ± 0.58	5.68 ± 1.13	0.132
ACR (mg/mmol)	0.76 (0.49, 1.50)	1.12 ± 0.68	0.71 (0.57, 1.88)	0.891

Continuous, normally distributed variables are expressed as means ± SD and non-normally distributed variables are expressed as medians (25% and 75% quartiles)

M male, *F* female, *BMI* body mass index, *WHR* waist to hip ratio, *HbA1c* hemoglobin A1c, *FPG* fasting plasma glucose, *2hPG* 2h postprandial plasma glucose, *TG* triglycerides, *TC* total cholesterol, *HDL-c* HDL-cholesterol, *LDL-c* LDL-cholesterol, *Cr* creatinine, *BUN* blood urea nitrogen, *ACR* urinary albumin to urinary creatinine ratio

**p* < 0.05 compared to CTRL, #*p* < 0.05 compared to IGT, based on one-way ANOVA with SNK analysis, Kruskal–Wallis H test or v2 test, as appropriate

OPEN ACCESS

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>), which

permits any noncommercial use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.