

Diaphragmatic Myoclonus

We diagnosed abdominal myoclonus in an 8-year-old boy with terminal stage of neuronal ceroid lipofuscinosis (NCL). He was normal up to two years of age when he had onset of myoclonic seizures. Over time his seizures worsened. He lost speech and mobility by six years. Recently he was admitted with recurrent seizures and frequent jerking of the abdominal muscles. His three siblings had died of similar illness. Examination revealed choreathetoid movements of limbs, multifocal myoclonus, and diaphragmatic myoclonus (*Web Video I*). Surface electromyography (EMG) electrodes applied on the abdomen wall revealed around 120 contractions per minute. Electroencephalography (EEG) showed bihemispherical periodic discharges at rate of 1.5 to 2 per second, left side more than right. Intravenous bolus of phenytoin followed by midazolam infusion controlled his abdominal (diaphragmatic) contractions.

Diaphragmatic myoclonus (flutter) is an abnormal movement of diaphragm resulting in inward and outward movements of abdominal wall synchronous with diaphragm contractions. In case other respiratory muscles are involved, it is called respiratory myoclonus. Other names like moving umbilicus syndrome, and belly



FIG. 1 Diaphragmatic myoclonus in a child (see Video at website).

dancer's dyskinesia have also been given. The diagnosis of diaphragmatic movements can be confirmed by EMG of diaphragm, or on fluoroscopy. The origin of the myoclonus is believed to be the brainstem, basal ganglia or cortex.

*ROSHAN KOUL AND AMNA ALFUTAISI

Department of Child Health (Neurology),
Sultan Qaboos University Hospital and
College of Medicine and Health Sciences,
Muscat, Oman.

*roshankoul@hotmail.com

NOTICE

Call for Submission of 'Clinical Videos'

Under this section, *Indian Pediatrics* publishes videos depicting an intricate technique or an interesting clinical manifestation, which are difficult to describe clearly in text or by pictures. A video file submitted for consideration for publication should be of high resolution and should be edited by the author in final publishable format. MPEG or MP4 formats are acceptable. The maximum size of file should be 20 MB. The file should not have been published elsewhere, and will be a copyright of *Indian Pediatrics*, if published. For this section, there should be a write-up of up to 250 words discussing the condition and its differential diagnoses. The write up should also be accompanied by a thumbnail image for publication in the print version and PDF. Submit videos as separate Supplementary files with your main manuscript. A maximum of three authors (not more than two from a single department) are permissible for this section. In case the video shows a patient, he/she should not be identifiable. In case the identification is unavoidable, or even otherwise, each video must be accompanied by written permission of parent/guardian, as applicable. Authors are responsible for obtaining participant consent-to-disclose forms for any videos of identifiable participants, and should edit out any names mentioned in the recording. The consent form should indicate its purpose (publication in the journal in print and online, with the understanding that it will have public access) and the signed consent of the parent/legal guardian. The copy of the consent form must be sent as supplementary file along with the write-up, and original form should be retained by the author. A sample consent form is available at our website www.indianpediatrics.net.