

Hydroxyzine-induced Vertical Nystagmus

Nystagmus is an involuntary movement in which the eye moves both rapidly and slowly on a vertical or horizontal axis. Acquired nystagmus can result from a range of neurological disorders, such as cerebellar disease, multiple sclerosis, vestibular disease, stroke, tumors, and trauma; or as an adverse effect of drugs (sedatives, anticonvulsants and alcohol). Horizontal nystagmus can be of peripheral-vestibular or central origin, while vertical nystagmus most often results from a lesion of the ocular motor pathways or centered within the brainstem. Drugs can cause both vertical and horizontal nystagmus *via* the dopaminergic-cholinergic blockade; for example, hydroxyzine is an H1-receptor blocker that can have anticholinergic effects.

An 8-year-old girl presented with fever and nuchal rigidity, and a fluid examination confirmed the diagnosis of meningitis. Vancomycin and ceftriaxone were administered; however, due to an allergic rash that appeared after the infusion of the ceftriaxone, oral hydroxyzine was started (2 mg/kg/day divided into 3 doses). On the 3rd day of treatment, the patient complained that her eye ball were moving up and down. Upon physical examination, a medium degree of vertical nystagmus was detected in both eyes (*Fig. 1* and *Web Video 1*); there were no additional neurological findings. Her brain magnetic resonance imaging was normal, and no alternative causes of the vertical nystagmus were



FIG.1 Vertical nystagmus

found. Hydroxyzine was discontinued, the nystagmus disappeared within 24 hours, and it did not relapse. After that, the meningitis therapy was completed, and the patient was discharged without any sequelae.

The most common side effects of hydroxyzine are drowsiness and decreased alertness. Voltage-gated sodium channels are expressed in the cerebellar Purkinje neurons, and sodium channel blocking is considered to be the cause of nystagmus in patients with antihistamine poisoning.

**GÜLSÜM ALKAN¹, MELIKE EMİROLU¹
AND AYSE KARTAL²**

*Departments of ¹Pediatric Infectious Diseases and
²Pediatric Neurology; Selcuk University Faculty of Medicine,
Konya, Turkey.
¹galkan-85@hotmail.com*

NOTICE

Call for Submission of 'Clinical Videos'

Under this section, *Indian Pediatrics* publishes videos depicting an intricate technique or an interesting clinical manifestation, which are difficult to describe clearly in text or by pictures. A video file submitted for consideration for publication should be of high resolution and should be edited by the author in final publishable format. MPEG or MP4 formats are acceptable. The maximum size of file should be 20 MB. The file should not have been published elsewhere, and will be a copyright of *Indian Pediatrics*, if published. For this section, there should be a write-up of up to 250 words discussing the condition and its differential diagnoses. The write-up should also be accompanied by a thumbnail image for publication in the print version and PDF. Submit videos as separate Supplementary files with your main manuscript. A maximum of three authors (not more than two from a single department) are permissible for this section. In case the video shows a patient, he/she should not be identifiable. In case the identification is unavoidable, or even otherwise, each video must be accompanied by written permission of parent/guardian, as applicable. Authors are responsible for obtaining participant consent-to-disclose forms for any videos of identifiable participants, and should edit out any names mentioned in the recording. The consent form should indicate its purpose (publication in the journal in print and online, with the understanding that it will have public access) and the signed consent of the parent/legal guardian. The copy of the consent form must be sent as supplementary file along with the write-up, and original form should be retained by the author. A sample consent form is available at our website www.indianpediatrics.net.