



Correction to: Chasing myrtle rust in New Zealand: host range and distribution over the first year after invasion

M. Toome-Heller¹  · W. W. H. Ho¹ · R. J. Ganley^{2,3} · C. E. A. Elliott¹ · B. Quinn¹ · H. G. Pearson⁴ · B. J. R. Alexander¹

Published online: 28 March 2020

© Australasian Plant Pathology Society Inc. 2020

Correction to: Australasian Plant Pathology

<https://doi.org/10.1007/s13313-020-00694-9>

The original version of this article unfortunately contained a mistake. The image order in Fig. 2 was incorrect. The corrected figure is given below.

The online version of the original article can be found at <https://doi.org/10.1007/s13313-020-00694-9>

✉ M. Toome-Heller
merje.toome@mpi.govt.nz

¹ Plant Health and Environment Laboratory, Ministry for Primary Industries, Auckland, New Zealand

² Scion, Rotorua, New Zealand

³ Present address: The New Zealand Institute for Plant and Food Research Limited, Te Puke, New Zealand

⁴ Biosecurity Surveillance and Incursion Investigation Plant Health, Ministry for Primary Industries, Wellington, New Zealand



Fig. 2 Symptoms on new and most common hosts of *Austropuccinia psidii* in New Zealand. New host records are marked with an asterisk. (a) *Acca sellowiana*; (b) *Eucalyptus globoidea*; (c) *Lophomyrtus bullata*; (d) *Lophomyrtus obcordata*; (e) *L. bullata/ L. obcordata* hybrid; (f)

Melaleuca salignus; (g) *Melaleuca viminalis*; (h) *Metrosideros bartlettii**; (i) *Metrosideros diffusa**; (j) *Metrosideros excelsa*; (k) *Metrosideros fulgens**; (l) *Metrosideros perforata**; (m) *Metrosideros robusta**; (n) *Syzygium australe*; (o) *Syzygium maire**