First report of *Neoscytalidium novaehollandiae* on common sage (*Salvia officinalis*)

Sibel Derviş¹ · İnci Güler Güney¹ · İslim Koşar² · Tuğba Bozoğlu³ · Göksel Özer³

Received: 22 June 2021 / Accepted: 27 July 2021 / Published online: 31 July 2021
© Australasian Plant Pathology Society Inc. 2021

Abstract

In June 2020, many plants exhibited symptoms of root rot and foliar blight in the experimental field of common sage in Şanlıurfa province, Turkey. The pathogen was identified as *Neoscytalidium novaehollandiae* based on morphological characteristics and phylogenetic analysis of partial sequence of the transcription elongation factor 1- α gene and the internal transcribed spacer of rDNA. Koch's postulates were fulfilled by successful re-isolation of the pathogen from inoculated plants in the pathogenicity assay. To our best knowledge, this is the first report of *N. novaehollandiae* causing root rot and foliar blight of common sage worldwide.

Keywords *Salvia officinalis* · *Neoscytalidium novaehollandiae* · Foliar blight · Root rot

The genus *Salvia* L. belongs to the mint family Lamiaceae, and comprises about 900 species, spread throughout the world (Delamare et al. 2007). Some of the species have great economic value since they are used as spices and flavouring agents by cosmetic and perfumery industries. *Salvia* species are odorous small shrubs native to the Mediterranean region (Albania, Croatia, Greece, Italy, Turkey, etc.) and cultivated worldwide (Raaijmakers et al. 2007). In Turkey, the genus *Salvia* is represented by over 100 taxa, half of which are endemic, and 1,271 tons of sage production is conducted on a cultivation area of over 660 ha in 2020 (TURKSTAT 2021). Common sage (*Salvia officinalis* L.) is a perennial herb or sub-shrub and consumed for aromatic and medicinal species credited with a long list of medicinal use (Newall et al. 1996).

Several fungal and oomycetous species were reported on *S. officinalis* including *Golovinomyces biocellatus* (Radisek et al. 2012) and *G. neosalviae* (Venegas-Portilla et al. 2020) causing powdery mildew, *Sclerotinia sclerotiorum* causing

white mold (Garibaldi et al. 2004) *Peronospora lamii* causing downy mildew (Humphreys-Jones et al. 2008), and *Phytophthora cryptogea* causing root rot (Garibaldi et al. 2015; Çakır et al. 2017). In June 2020, symptoms including yellowing and shedding of leaves, foliar blight, and root rot prior to plant death were observed on two-year-old *S. officinalis* cv. Elif plants in the sage fields of Koruklu village (36°42' N; 38°58' E, 410 m altitude), Akçatepe district of Şanlıurfa province (Fig. 1a, b). The incidence of diseased plants was approximately 25%.

To isolate the pathogen, 20 diseased plants were sampled from the field. Symptomatic roots of plants were cleaned thoroughly from soil residues under tap water. Portions (about 3–5 mm) of necrotic root tissues were surface disinfested in 1% sodium hypochlorite solution for 60 s and rinsed twice in sterile water, air dried, and placed onto potato dextrose agar (PDA) amended with 0.2 g/l streptomycin sulphate. After a five-day incubation period at 26 °C in the dark, fungal colonies with similar morphological characteristics were consistently isolated and transferred to new PDA plates using the hyphal tip method. The 7-day-old hyphae from the representative isolate (Nn-01Sf) were also transferred to 1.5% water agar containing sterile pine needles to promote the formation of conidiomata.

Colonies were initially powdery white with dense, hairy aerial mycelium, which becoming dark gray to black within ten days (Fig. 2a, b). Arthroconidia were dark brown, thick-walled, cylindrical to oblong, 0 to 1 septate, 6.7 × 4.2 µm

✉ Göksel Özer
gokozer@gmail.com

¹ Department of Plant and Animal Production, Mardin Artuklu University, Vocational School of Kızıltepe, Mardin 47000, Turkey

² GAP Agricultural Research Institute, Şanlıurfa 63040, Turkey

³ Faculty of Agriculture, Department of Plant Protection, Bolu Abant İzzet Baysal University, Bolu 14020, Turkey

Fig.1 Symptoms on *Salvia officinalis* plants; **a** foliar blight in the field, **b** root rot in the field, **c** foliar blight and root rot symptoms on five healthy 2-year-old common sage cv. Elif plants after 14 days of inoculation

Fig.2 Characteristics of *Neoscytalidium novaehollandiae* isolate Nn01Sf; **a**, **b** colony morphologies on PDA, **c** arthroconidia, **d** pycnidia, **e** conidia formed in pycnidia

($n=30$; Fig. 2c), formed both singly and in arthric chains by hyphal fragmentation. Pycnidia observed on pine needles on water agar were stromatic, semi-immersed, and black with a mean diameter of $310\ \mu\text{m}$ (Fig. 2d). Conidia formed in pycnidia were hyaline, oblong to globose, apices rounded, 0- to 1-septate, and $11.2 \times 4.6\ \mu\text{m}$ in length ($n=30$; Fig. 2e). These morphological characteristics were consistent with the description of *Neoscytalidium* spp. by Phillips et al. (2013).

For further precise identification, genomic DNA of the representative isolate was extracted using DNeasy Blood and Tissue Kit (Qiagen GmbH, Hilden, Germany) according to the manufacturer's protocol. The sequences of the translation elongation factor 1- α gene (EF1- α) and the internal transcribed spacer region (ITS) of rDNA were amplified using primers EF1-728 F and EF1-986R (Carbone and Kohn 1999) and ITS1 and ITS4 (White et al. 1990), respectively, and both strands of the PCR products were sequenced at the MacroGen Inc. Sequencing Service (Seoul, Korea).

All sequences of the isolate were read and edited checked manually and nucleotide arrangements at ambiguous positions were clarified using both primer directions using MEGA X software (Kumar et al. 2018). Multiple sequence alignment was constructed using the online sequence alignment tool, MAFFT (<https://mafft.cbrc.jp/alignment/server/large.html>, Katoh et al. 2019). Phylogenetic analysis on the combined dataset was done using PAUP (Phylogenetic Analysis Using Parsimony) version 4.0b10 (Swofford 2003) for Maximum Likelihood analyses and visualized TREE-INDER version of March 2011 (Jobb 2011) and MEGA X. The isolate was clustered closely with the isolates of *N. novaehollandiae* in the phylogenetic tree constructed (Fig. 3). The resultant sequences were deposited in GenBank with Accessions Nos. MZ392224 and MZ389710 for the EF1- α and ITS loci, respectively, and compared with the sequences registered in GenBank based on nucleotide similarity. A BLASTn search of the resultant sequences showed 99.30 and 100.00% identity with the EF1- α (Accessions No. EF585581) and ITS (Accessions No. EF585535) sequences of *N. novaehollandiae* CBS122071 strain, respectively. Based on morphological and molecular identification, the isolates were identified as *Neoscytalidium novaehollandiae* Pavlic, T.I. Burgess & M.J. Wingf. The isolate Nn-01Sf was deposited in Erciyes University Culture Collection (EUCC—WDCM 1202) with the accession number EUCC- 21,068 M.

Pathogenicity tests performed were by spraying conidial suspension (1×10^6 conidia/mL) of *N. novaehollandiae* Nn-01Sf strain on leaves and stems of five healthy 2-year-old *S. officinalis* cv. Elif plants, till run off. Five control plants were treated with sterile water. After inoculation, the plants were maintained in a moist chamber constructed using plastic bags to maintain moisture for 24 h. These bags were removed, and all plants were transferred to a growth chamber at $25\ ^\circ\text{C}$ with a 16-h light/dark photoperiod. Two

Fig. 3 Phylogenetic tree inferred by using the Maximum Likelihood method was reconstructed using the EF1- α and ITS sequences of *Neoscytalidium novaehollandiae* isolate Nn01Sf. The bootstrap value (percentage, based on 1,000 replications) is shown on the branch

weeks after inoculation, root rot and foliar blight symptoms (Fig. 1c) were observed on all plants inoculated, whereas all control plants remained symptomless. The pathogen was re-isolated from all inoculated plants and identified by conidial morphology, fulfilling Koch's postulates.

Neoscytalidium species cause canker, dieback, stem blight and root rot diseases on several hosts of agronomic importance. *N. novaehollandiae* has been reported as a destructive pathogen in Turkey causing stem canker and branch dieback of almond trees (Ören et al. 2020), stem blight on tomato (Derviş et al. 2020), and wood canker caused by on grapevine (Akgül et al. 2019). However, there is no available data on *N. novaehollandiae* associated with common sage. To our knowledge, this is the first report of *Neoscytalidium novaehollandiae* on common sage in Turkey and worldwide (Farr and Rossman 2021).

Declarations

Research involving human participants and/or animal The authors declare that no human participants and animals were involved in this study.

Informed consent This manuscript is new and not being considered elsewhere. All authors have approved the submission of this manuscript.

Conflict of interest The authors declare that they have no conflict of interest.

References

- Akgül DS, Savaş NG, Özarslandan M (2019) First report of wood canker caused by *Lasiodiplodia exigua* and *Neoscytalidium novaehollandiae* on grapevine in Turkey. *Plant Dis* 103(5):1036. <https://doi.org/10.1094/PDIS-11-18-1938-PDN>
- Carbone I, Kohn LM (1999) A method for designing primer sets for speciation studies in filamentous ascomycetes. *Mycologia* 91:553–556. <https://doi.org/10.2307/3761358>
- Çakır E, Bahtiyarca Bağdat R, Katırcıoğlu YZ, Maden S (2017) Occurrence of root rot caused by *Phytophthora cryptogea* on common sage (*Salvia officinalis*) in Turkey. *J Agric Sci Technol A* 7:401–406. <https://doi.org/10.17265/2161-6256/2017.06.005>
- Delamare APL, Moschen-Pistorello IT, Artico L, Atti-Serafini L, Echeverrigaray S (2007) Antibacterial activity of the essential oils of *Salvia officinalis* L. and *Salvia triloba* L. cultivated in South Brazil. *Food Chem* 100(2):603–608. <https://doi.org/10.1016/j.foodchem.2005.09.078>
- Derviş S, Özer G, Türkölmez Ş (2020) First report of *Neoscytalidium novaehollandiae* causing stem blight on tomato in Turkey. *Eur J Plant Pathol* 102(4):1339–1340. <https://doi.org/10.1007/s42161-020-00627-x>
- Farr DF, Rossman AY (2021) Fungal Databases, U.S. National Fungus Collections, ARS, USDA. <https://nt.ars-grin.gov/fungaldatabases/>. Accessed 03 June 2021
- Garibaldi A, Minuto A, Gullino ML (2004) First report of white mold caused by *Sclerotinia sclerotiorum* on Common Sage (*Salvia officinalis*) in Italy. *Plant Dis* 88(9):1044–1044. <https://doi.org/10.1094/PDIS.2004.88.9.1044B>
- Garibaldi A, Bertetti D, Pensa P, Ortu G, Gullino ML (2015) *Phytophthora cryptogea* on common sage (*Salvia officinalis* L.) in Italy. *Plant Dis* 99(1):161–161. <https://doi.org/10.1094/PDIS-09-14-0902-PDN>
- Humphreys-Jones DR, Barnes AV, Lane CR (2008) First report of the downy mildew *Peronospora lamii* on *Salvia officinalis* and *Rosmarinus officinalis* in the UK. *Plant Pathol* 57(2). <https://doi.org/10.1111/j.1365-3059.2007.01654.x>
- Jobb G (2011) TREEFINDER version of March 2011. Munich, Germany (<http://www.treefinder.de>).
- Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. *Brief Bioinform* 20(4):1160–1166. <https://doi.org/10.1093/bib/bbx108>
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: Molecular Evolutionary Genetics Analysis across computing platforms. *Mol Biol Evol* 35:1547–1549. <https://doi.org/10.1093/molbev/msy096>
- Newall CA, Anderson LA, Phillipson JD (1996) Herbal Medicines. The Pharmaceutical Press, London, A Guide for Healthcare Professionals, p 231
- Ören E, Koca G, Gencer R, Bayraktar H (2020) First report of *Neoscytalidium novaehollandiae* associated with stem canker and branch dieback of almond trees. *Australas Plant Dis Notes* 15(1):1–3. <https://doi.org/10.1007/s13314-020-00386-9>
- Phillips AJ, Alves A, Abdollahzadeh J, Slippers B, Wingfield MJ, Groenewald JZ, Crous PW (2013) The Botryosphaeriaceae: genera and species known from culture. *Stud Mycol* 76:51–167. <https://doi.org/10.3114/sim0021>
- Raal A, Orav A, Arak E (2007) Composition of the essential oil of *Salvia officinalis* L. from various European countries. *Nat Prod Res* 21(5):406–411. <https://doi.org/10.1080/14786410500528478>
- Radisek S, Ferant N, Jakse J, Javornik B (2012) First report of powdery mildew caused by *Golovinomyces biocellatus* on common sage (*Salvia officinalis*) in Slovenia. *Plant Dis* 96(7):1065–1065. <https://doi.org/10.1094/PDIS-01-12-0010-PDN>
- Swofford DL (2003) PAUP* ver 4.0. b10. Phylogenetic Analysis Using Parsimony and Other Methods. Sunderland, MA: Sinauer Associates, Sunderland
- TURKSTAT (2021) Turkish Statistical Institute. Crop Production Statistics. <https://biruni.tuik.gov.tr/medas/?kn=92&locale=tr>. Accessed 05 June 2021
- Venegas-Portilla A, Solano-Báez AR, Leyva-Mir SG, Camacho-Tapia M, Lima NB, Tovar-Pedraza JM, Márquez-Licona G (2020) First Report of *Golovinomyces neosalviae* Causing Powdery Mildew on Common Sage (*Salvia officinalis*) in Mexico. *Plant Dis* 104(5):1546. <https://doi.org/10.1094/PDIS-11-19-2320-PDN>
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (eds) PCR Protocols. A Guide to Methods and Applications, Academic Press, San Diego, CA, USA, pp 315–322