



Correction to: Effects and Mechanism of Action of *Panax notoginseng* Saponins on the Pharmacokinetics of Warfarin

Jiafen Qian¹ · Wenjun Chen¹ · Jianmei Wu¹ · Meina Lv¹ · Shaojun Jiang¹ · Zhiwei Zeng¹ · Zongwei Fang¹ · Mingrong Chen¹ · Jinhua Zhang¹

Published online: 22 February 2022
© Springer Nature Switzerland AG 2022

Correction to:
European Journal of Drug Metabolism and Pharmacokinetics
<https://doi.org/10.1007/s13318-022-00753-0>

The original version of this article unfortunately contained a few typos in the abstract.

Abstract, First section heading, which previously read:

Background

should read:

Background

Abstract, Results: 3rd sentence, which previously read:

Compared with the warfarin alone group, the clearance rate (CL/F) of warfarin in the warfarin + low-dose PNS, warfarin

+ medium-dose PNS, and warfarin + high-dose PNS groups was 10% ($P > 0.05$), 23% ($P < 0.05$), and 33% ($P < 0.05$) lower, respectively, while the systemic exposure (area under the concentration-time curve, AUC_{0-t}) increased by 106% ($P > 0.05$), 119% ($P < 0.05$), and 134% ($P < 0.05$), respectively, and the blood concentration of warfarin increased by 112%, 113%, and 114%, respectively (all $P > 0.05$).

Should read:

Compared with the warfarin alone group, the clearance rate (CL/F) of warfarin in the warfarin + low-dose PNS, warfarin + medium-dose PNS, and warfarin + high-dose PNS groups was 10% ($P > 0.05$), 23% ($P < 0.05$), and 33% ($P < 0.05$) lower, respectively, while the systemic exposure (area under the concentration-time curve, AUC_{0-t}) increased by 106% ($P > 0.05$), 119% ($P < 0.05$), and 134% ($P < 0.05$), respectively, and the blood concentration of warfarin increased by 112%, 113%, and 114%, respectively (all $P > 0.05$).

The original article can be found online at <https://doi.org/10.1007/s13318-022-00753-0>.

✉ Jinhua Zhang
pollyzhang2006@126.com

¹ Department of Pharmacy, Fujian Medical University Union Hospital, #29 Xinquan Road, Fuzhou 350001, Fujian, China