



Correction to: On Borg's method for non-selfadjoint Sturm–Liouville operators

Sergey Buterin¹ · Maria Kuznetsova¹

Published online: 18 June 2019
© Springer Nature Switzerland AG 2019

Correction to: Analysis and Mathematical Physics

<https://doi.org/10.1007/s13324-019-00307-9>

The original version of the article unfortunately contained few misprints under Preliminaries section. The corrected text is given below.

$$\mathcal{D}_j = \begin{bmatrix} d_{n_{1j}} & & \\ & d_{n_{2j}} & \\ & & \ddots \end{bmatrix}$$

with the square cells d_{n_j} determined in (13), where $\{n_{kj}\}_{k \geq 1}$ is the increasing sequence of all elements of the set \mathcal{S}_j . The other elements of \mathcal{D}_j (i.e. not included in d_{n_j} 's) are equal to zero. Note that, according to the asymptotics (2), $d_{n_{\nu j}} = [1]$ for sufficiently large ν .

The Original Article was corrected.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s13324-019-00307-9>.

✉ Sergey Buterin
buterinsa@info.sgu.ru

Maria Kuznetsova
kuznetsovama@info.sgu.ru

¹ Department of Mathematics, Saratov State University, Astrakhanskaya 83, Saratov, Russia 410012