



Correction to: Severe hypoglycemia caused by a small dose of repaglinide and concurrent use of nilotinib and febuxostat in a patient with type 2 diabetes

Hitomi Komatsu¹ · Mariko Enomoto¹ · Hisashi Shiraishi¹ · Yasuyo Morita¹ · Daisuke Hashimoto² · Shuichi Nakayama² · Shogo Funakoshi² · Seiki Hirano² · Yoshio Terada² · Mitsuhiro Miyamura¹ · Shimpei Fujimoto²

Published online: 23 April 2020
© The Japan Diabetes Society 2020

Correction to: *Diabetology International*
<https://doi.org/10.1007/s13340-020-00434-w>

In the original publication, the unit of C-peptide immunoreactivity (CPR) in Table 1 was published incorrectly. The correct unit is ng/ml.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s13340-020-00434-w>.

✉ Shimpei Fujimoto
fujimoto@kochi-u.ac.jp

¹ Department of Pharmacy, Kochi Medical School Hospital, Nankoku, Japan

² Department of Endocrinology, Metabolism, and Nephrology, Kochi Medical School, Kochi University, Kohasu, Oko-cho, Nankoku, Kochi 783-8505, Japan