Correction to: Decline in the estimated glomerular filtration rate (eGFR) following metabolic control and its relationship with baseline eGFR in type 2 diabetes with microalbuminuria or macroalbuminuria

Shoichi Akazawa¹ · Eiji Sadashima² · Yasunori Sera¹ · Nobuhiko Koga³

Published online: 27 November 2021
© The Japan Diabetes Society 2021

Correction to: *Diabetology International*
<https://doi.org/10.1007/s13340-021-00517-2>

In the original publication of the article, in Table 3, under Macroalbuminuria ($n = 81$), on the row “HbA1c”, the value of “Correlation coefficient and Regression coefficient” of Multivariate was incorrectly published as “ -0.205 and -3.34 ”.

The correct coefficient values should read as “ 0.205 and 3.34 ”.

The correct table 3 is provided below.

The original article can be found online at <https://doi.org/10.1007/s13340-021-00517-2>.

✉ Shoichi Akazawa
akazawa@tenjinkai.or.jp

¹ Center of Diabetes, Koseikai Hospital, Hayama 1-3-2, Nagasaki, Japan

² Life Science Institute, Saga-Ken Medical Center Koseikan, Saga, Japan

³ Shin-Koga Hospital, Kurume, Japan

Table 3 Univariate and multivariate regression analyses for baseline eGFR in patients with type 2 diabetes with microalbuminuria and macroalbuminuria

	Univariate				Multivariate			
	Correlation coefficient (r)	Regression coefficient	Standard error	p value	Correlation coefficient (r)	Regression coefficient	Standard error	p value
Microalbuminuria (n = 124)								
Sex (male)	0.000	0.009	1.879	0.996				
BMI	0.007	0.394	0.394	0.386				
Duration of diabetes	− 0.148	− 0.352	0.219	0.111				
Mean blood pressure	0.213	0.389	0.159	0.017	0.088	0.158	0.145	0.279
HbA1c	0.404	6.714	1.374	<0.001	0.39	6.350	1.282	<0.001
Urinary albumin excretion	− 0.054	− 0.013	0.022	0.551				
Total cholesterol	0.144	0.088	0.055	0.113				
HDL-cholesterol	0.121	0.187	0.139	0.193				
Triglyceride	− 0.007	− 0.001	0.016	0.929				
Plaque in carotid artery	− 0.355	− 1.046	0.254	<0.001	− 0.319	− 0.942	0.238	0.010
$(R^2=0.270, p<0.001)$								
Macroalbuminuria (n = 81)								
Sex (male)	0.026	0.609	2.537	0.811				
BMI	0.168	0.712	0.470	0.134				
Duration of diabetes	− 0.287	− 0.662	0.265	0.015	− 0.258	− 0.605	0.264	0.025
Mean blood pressure	0.111	0.155	0.156	0.323				
HbA1c	0.205	3.833	2.053	0.065	0.205	3.34	1.796	0.067
Urinary protein excretion	− 0.358	− 6.226	1.862	0.001	− 0.214	− 3.909	2.081	0.065
Total cholesterol	− 0.140	− 0.011	0.009	0.256				
HDL-cholesterol	− 0.244	− 0.049	0.036	0.185				
Triglyceride	− 0.100	− 0.003	0.002	0.175				
Plaque in carotid artery	− 0.355	− 1.046	0.254	<0.001	− 0.241	− 0.768	0.37	0.042
$(R^2=0.227, p<0.001)$								

Independent variables comprise the baseline value of eGFR (before the start of diabetes dialysis prevention). Mean blood pressure = diastolic pressure + systolic pressure − diastolic pressure/3. Plaque in the carotid artery is the sum of the plaque thickness in the carotid artery

BMI body mass index, *eGFR* estimated glomerular filtration rate, *HDL* high-density lipoprotein, *UAC* urinary albumin-to-creatinine excretion rate, *UPC* urinary protein-to-creatinine excretion rate

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.