




## Correction: Time course change of the insulin requirements during the perioperative period in hepatectomy and pancreatectomy by using an artificial pancreas STG-55

Sanae Teshigawara<sup>1</sup> · Atsuhito Tone<sup>1</sup>  · Akihiro Katayama<sup>2</sup> · Yusuke Imai<sup>1</sup> · Toshihisa Tahara<sup>1</sup> · Mayumi Senoo<sup>1</sup> · Satoko Watanabe<sup>1</sup> · Mitsuhiro Kaneto<sup>1</sup> · Yasuyuki Shimomura<sup>1</sup> · Chiaki Yagi<sup>3</sup> · Hiroki Kajioka<sup>3</sup> · Toru Kojima<sup>3</sup> · Takefumi Niguma<sup>3</sup> · Tatsuaki Nakatou<sup>1</sup>

Published online: 28 April 2023  
© The Japan Diabetes Society 2023

**Correction: Diabetology International**  
<https://doi.org/10.1007/s13340-023-00623-3>

In the original publication of the article, funding information was published incorrectly. The correct funding information should read as “None”.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

---

The original article can be found online at <https://doi.org/10.1007/s13340-023-00623-3>.

---

✉ Atsuhito Tone  
aitone@ms1.megaegg.ne.jp

<sup>1</sup> Department of Internal Medicine, Diabetes Center, Okayama Saiseikai General Hospital, 2-25 Kokutaicho, Kita-Ku, Okayama 700-8511, Japan

<sup>2</sup> Department of Diabetology and Metabolism, National Hospital Organization, Okayama Medical Center, Okayama, Japan

<sup>3</sup> Department of Surgery, Okayama Saiseikai General Hospital, Okayama, Japan