Correction to: Molecular quantification of spider preying on springtails in winter-spring fallow waterlogged paddy fields

Jiaqi Sun¹ · Shuai Pang¹ · Xueqin Wang^{2,3} · Guanghua Wang^{2,3} · Zengrong Zhu³ · Jiaan Cheng³ · Yaoyu Bai¹

Published online: 30 January 2021

© The Japanese Society of Applied Entomology and Zoology 2021

Correction to: Applied Entomology and Zoology
<https://doi.org/10.1007/s13355-020-00712-8>

In the original publication of the article, on the 3rd paragraph of results section, the following sentence “They were five dominant species in rice paddy ecosystem, including 187 *P. subpiraticus*, 74 (Sundevall) (Araneida: Linyphiidae); (Bösenberg and Strand) (Araneida: Linyphiidae) (Bösenberg and Strand) (Araneida: Clubionidae); (Karsch) (Araneida: Salticidae).” was published incorrectly.

The correct sentence should read as below,

They were five dominant species in rice paddy ecosystem, including 187 *P. subpiraticus*, 74 *Erigonidium graminicolum* (Sundevall) (Araneida: Linyphiidae), 52 *Oedothorax insec-ticeps* (Bösenberg and Strand) (Araneida: Erigonidae), 23

Clubiona japonicola (Bösenberg and Strand) (Araneida: Clubionidae) and 38 *Marpissa magister* (Karsch) (Araneida: Salticidae).

The original article has been corrected.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s13355-020-00712-8>.

✉ Yaoyu Bai
yybai711@163.com

¹ College of Plant Protection, Southwest University,
2 # Tiansheng Road, Beibei 400715, Chongqing,
People's Republic of China

² Institute of Plant Protection, Henan Academy of Agricultural
Sciences, Zhengzhou 450002, China

³ Institute of Insect Sciences, Zhejiang University,
Hangzhou 310058, China