Correction to: Use of several host plants with different seed maturation times by the chestnut weevil, *Curculio sikkimensis* (Coleoptera: Curculionidae), in mixed oak and chestnut coppice stands

Morio Higaki¹

Published online: 15 May 2021

© The Japanese Society of Applied Entomology and Zoology 2021

Correction to: Applied Entomology and Zoology
<https://doi.org/10.1007/s13355-021-00740-y>

In the original publication of the article, the Fig. 4 was published incorrectly. The correction Fig. 4 is provided below.

The original article was updated.

The original article can be found online at <https://doi.org/10.1007/s13355-021-00740-y>.

✉ Morio Higaki
scarabee@affrc.go.jp

¹ Institute of Fruit Tree and Tea Science, National Agriculture and Food Research Organization, Tsukuba, Ibaraki 305-8605, Japan

Fig. 4 **a** Seasonal change in the number of *Curculio sikimensis* larvae per seed of three host species (*Castanea crenata*, *Quercus serrata*, and *Quercus myrsinifolia*) in 2004, which is shown for each 5- or 6-day period within which the sample was collected. **b** Comparison between the probability density functions (PDFs) of the estimated total amount of oviposition in 2004 ($\hat{\mu}_t$) and the PDF of the amount of available seeds of the three host species. The quantity is shown as the probability density for each 5- or 6-day period within which the sample was collected