Correction to: Susceptibility of nymphs and adults of *Blattella germanica* (Dictyoptera: Blattellidae) to *Aspergillus terreus* (Eurotiales: Trichocomaceae) infested via oral ingestion, contact, and bait methods: a comparative study

Yamina Djouhri^{1,2} · Mahdia Saidi-Touati² · Amina Benelmouffok³ · Mohamed Nadji Bouchtout⁴

Published online: 26 February 2022

© The Author(s) under exclusive licence to The Japanese Society of Applied Entomology and Zoology 2022

Correction to: Applied Entomology and Zoology
<https://doi.org/10.1007/s13355-022-00770-0>

In the original publication of the article, the values of “ α β γ δ ” were placed incorrectly in tables 1 and 2.

The correct Tables 1 and 2 are provided in this correction article.

The original article has been corrected.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s13355-022-00770-0>.

✉ Yamina Djouhri
djouhri.yamina@gmail.com

¹ Laboratory of Biodiversity and Environment: Interactions and Genomes, Faculty of Biological Sciences, University of Science and Technology Houari Boumediene, BP 32, El Alia Bab Ezzouar, CP 16000 Algiers, Algeria

² Laboratory of Dynamics and Biodiversity, Team of Arthropods Biodiversity, Faculty of Biological Sciences, University of Science and Technology Houari Boumediene, BP 32, El Alia Bab Ezzouar, CP 16000 Algiers, Algeria

³ Mycology Department, Pasteur Institute of Algeria, Staouéli, Dély-Brahim, Algiers, Algeria

⁴ Laboratory of Cellular and Molecular Biology, Cytokine and NO Synthase Team, University of Science and Technology Houari Boumediene, BP 32, El Alia Bab Ezzouar, CP 16000 Algiers, Algeria

Contamination mode	<i>A. terreus</i> concentration (spores/mL)	Days post-inoculation									
		Day 3		Day 6		Day 9		Day 12		Day 15	
		$\alpha\beta\gamma\delta$		$\alpha\beta\gamma\delta$		$\alpha\beta\gamma\delta$		$\alpha\beta\gamma\delta$		$\alpha\beta\gamma\delta$	
Ingestion	Control	0±0	a	0±0	a	0±0	a	3.33±3.33	a	3.33±3.33	a
	1×10 ⁶	3.33±3.33	a a a	13.33±3.33	a a a	16.67±3.33	a a a	16.67±8.81 ^s	a a a	16.67±8.82 ^s	a a a
	1×10 ⁸	13.33±3.33	a a a	33.33±3.33	b a a	40±5.77	b a a	37.78±6.19 ^s	b a a	37.78±6.19 ^s	b a a
Direct contact	Control	0±0	a	0±0	a	3.33±3.33	a	3.33±3.33	a	3.33±3.33	a
	1×10 ⁶	3.33±3.33	a a a	6.67±6.67	a a a	13.70±3.17 ^s	a a a	20.37±5.46 ^s	b a a	27.41±2.59 ^s	b a a
	1×10 ⁸	10±0	b a a	20±0	b a b	35.44±2.29 ^s	b b a	38.15±4.27 ^s	b a a	38.15±4.27 ^s	b a a
Bait	Control	0±0	ns	0±0	ns	0±0	ns	0±0	ns	0±0	ns
	1×10 ⁶	0±0	ns	0±0	ns	3.33±3.33	ns	6.67±3.33	ns	6.67±3.33	ns
	1×10 ⁸	0±0	ns	0±0	ns	3.33±3.33	ns	6.67±3.33	ns	10±0	ns

Means with the symbol § are means corrected by Abbott's formula

Table 2 Cumulative mortality in percentages (mean $\pm$ standard error) of *Blattella germanica* adults exposed to two different concentrations of *Aspergillus terreus* strain IPAREG0319 administered by three different modes

Contamination mode	<i>A. terreus</i> concentration (spores/mL)	Days post-inoculation							
		Day 3		Day 6		Day 9		Day 12	
		α β γ δ		α β γ δ		α β γ δ		α β γ δ	
Ingestion	Control	3.33 $\pm$ 3.33	a	6.67 $\pm$ 3.33	a	6.67 $\pm$ 3.33	a	6.67 $\pm$ 3.33	a
	1 $\times$ 10 ⁶	7.03 $\pm$ 3.53 [§]	a a a	28.52 $\pm$ 3.29 [§]	a a a	60.74 $\pm$ 3.22 [§]	b a a	71.48 $\pm$ 3.29 [§]	b a a
Direct contact	1 $\times$ 10 ⁸	13.70 $\pm$ 3.16 [§]	a a a	38.89 $\pm$ 5.56 [§]	b a a	75.18 $\pm$ 2.59 [§]	b a a	82.22 $\pm$ 3.39 [§]	b a a
	Control	3.33 $\pm$ 3.33	a	6.67 $\pm$ 3.33	a	6.67 $\pm$ 3.33	a	6.67 $\pm$ 3.33	a
Bait	1 $\times$ 10 ⁶	10 $\pm$ 5.77 [§]	a a a	28.15 $\pm$ 39.63 [§]	a a a	42.46 $\pm$ 1.48 [§]	b a a	57.03 $\pm$ 1.48 [§]	b a a
	1 $\times$ 10 ⁸	17.04 $\pm$ 2.96 [§]	b a a	39.63 $\pm$ 8.02 [§]	b a a	53.70 $\pm$ 6.68 [§]	b b a	79.25 $\pm$ 9.63 [§]	b b b
	Control	0 $\pm$ 0	ns	3.33 $\pm$ 3.33	ns	3.33 $\pm$ 3.33	ns	3.33 $\pm$ 3.33	ns
	1 $\times$ 10 ⁶	0 $\pm$ 0	ns	0 $\pm$ 0 [§]	ns	0 $\pm$ 0 [§]	ns	6.67 $\pm$ 3.33 [§]	ns
	1 $\times$ 10 ⁸	0 $\pm$ 0	ns	3.33 $\pm$ 3.33 [§]	ns	3.33 $\pm$ 3.33 [§]	ns	10 $\pm$ 5.77 [§]	ns

Data presented are means of three replicates (10 insects per replicate)

For each formulation, the mean is followed by a letter in at least one of four attributed subcolumns (α , β , γ and δ)

Letters in subcolumn α represent statistical significance between control and treated groups of the same contamination mode

Letters in subcolumn β represent statistical significance between concentrations 1 $\times$ 10⁶ and 1 $\times$ 10⁸ of the same contamination mode

Letters in subcolumns γ and δ represent statistical significance between different modes for the 1 $\times$ 10⁶ and the 1 $\times$ 10⁸ concentrations, respectively

Different letters on the same subcolumn indicate significant statistical differences (Tukey's test $p < 0.05$); ns: not significant

Means with the symbol § are means corrected by Abbott's formula