Correction to: CSF inflammatory markers and neurocognitive function after addition of maraviroc to monotherapy darunavir/ritonavir in stable HIV patients: the CINAMMON study

T. J. Barber^{1,2} · A. Imaz³ · M. Boffito^{1,2} · J. Niubó³ · A. Pozniak¹ · R. Fortuny³ · J. Alonso⁴ · N. Davies^{1,2} · S. Mandalia^{1,2} · D. Podzamczek³ · B. Gazzard^{1,2}

Published online: 9 January 2018
© Journal of NeuroVirology, Inc. 2018

Correction to: J. Neurovirol.

<https://doi.org/10.1007/s13365-017-0600-6>

Due to a production error the bottom portion of Figure 1 was omitted. The corrected figure is given below.

The online version of the original article can be found at <https://doi.org/10.1007/s13365-017-0600-6>

✉ T. J. Barber
t.barber@nhs.net

¹ Chelsea and Westminster NHS Foundation Trust and St Stephen's AIDS Trust 4th Floor, St Stephen's Centre, Chelsea and Westminster Hospital, 369 Fulham Road, London SW10 9NH, UK

² Imperial College London, London, UK

³ Bellvitge University Hospital, Barcelona, Spain

⁴ Hospital Universitari Vall d'Hebron, Barcelona, Spain

Point estimates shown are median

Units and normal ranges:

- Neopterin: 4.2 +/- 1.0 nmol/l
- S100 B: <740pg/mL (shown here x10⁻²)
- Ferritin: <16ng/mL
- NFH: <473pg/mL (shown here x10⁻¹)