Correction to: Effect of fluoroscopic X-rays combined with iodinated radiographic contrast media on human hematological parameters

Benjamaporn Supawat^{1,2} · Sakornniya Wattanapongpitak^{1,2} · Singkome Tima³ · Suchart Kothan^{1,2} · Montree Tungjai^{1,2}

Published online: 29 June 2021

© Korean Society of Environmental Risk Assessment and Health Science 2021

Correction to:

Toxicology and Environmental Health Sciences
<https://doi.org/10.1007/s13530-021-00093-7>

Due to unfortunate oversight during the correction process
Figs. 1 and 2 have been interchanged. They should read:

The original article can be found online at <https://doi.org/10.1007/s13530-021-00093-7>.

✉ Montree Tungjai
mtungjai@gmail.com

¹ Department of Radiologic Technology, Faculty of Associated Medical Sciences, Chiang Mai University, Chiang Mai 50200, Thailand

² Center of Radiation Research and Medical Imaging, Department of Radiologic Technology, Faculty of Associated Medical Sciences, Chiang Mai University, Chiang Mai 50200, Thailand

³ Division of Clinical Microscopy, Department of Medical Technology, Faculty of Associated Medical Sciences, Chiang Mai University, Chiang Mai 50200, Thailand

Fig. 1 Morphology of red blood cells in blood from male subjects that was incubated with or without 5 mg/mL of each in three iodinated radiographic contrast media (iopromid, iohexol, and iodixanol) combined with irradiation to fluoroscopic x-ray for 10, 30, and 60 min and in the control blood groups. Scale bar = 10 μ m

Scale bar = 10 μ m

A
Fluoroscopic x-ray (70 kV, 0.8 mA) at dose rate 0.25 mGy/s

- Hematological parameters examination
- Morphology observation

B

Fig. 2 **a** The experimental design. **b** Irradiation system

The original article has been corrected.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.