



Correction to: Fatty acid homeostasis in honey bees (*Apis mellifera*) fed commercial diet supplements

Vanessa CORBY-HARRIS¹, Meghan M. BENNETT¹, Megan E. DEETER^{1,2},
 Lucy SNYDER¹, Charlotte MEADOR¹, Ashley C. WELCHERT^{1,2}, Amelia HOFFMAN¹,
 Bethany T. OBERNESSER^{1,2}, and Mark J. CARROLL¹

¹ Carl Hayden Bee Research Center, USDA-ARS, Tucson, AZ 85719, USA

² Department of Entomology, University of Arizona, Tucson, AZ 85721, USA

Received 4 January 2022 – Revised 5 January 2022 – Accepted 19 January 2022

Correction to: APIDOLOGIE (2021) 52:1195–1209

<https://doi.org/10.1007/s13592-021-00896-0>

The original article was published while updates to Table IV in the online resources were still being processed. Provided below is the finalized and updated table. The table along with the article's other online resources can be found here:

[https://static-content.springer.com/esm/
 art%3A10.1007%2Fs13592-021-00896-0/
 MediaObjects/13592_2021_896_MOESM1_
 ESM.pdf](https://static-content.springer.com/esm/art%3A10.1007%2Fs13592-021-00896-0/MediaObjects/13592_2021_896_MOESM1_ESM.pdf)

Life stage	Tissue	Diet ¹	N	$\omega 6_{(\min, \max)}$	$\omega 3_{(\min, \max)}$	$\omega 6:\omega 3$
Queens	Whole	Out-door	3	0.254	4.996	0.05
			(0.108, 0.268)	(1.828, 10.259)		
	Low	High	5	0.074	1.927	0.04
			(0.023, 0.081)	(0.829, 2.191)		
Larvae	Whole	Out-door	4	0.100	1.824	0.05
			(0.085, 0.113)	(1.722, 2.320)		
		Not measured	4	0.052	1.030	0.05
			(0.018, 8.221)	(0.340, 2.464)		
Pupae	Head	Low	4	0.040	1.456	0.03
			(0.001, 0.056)	(0.054, 1.944)		
		High	3	0.050	0.769	0.06
			(0.016, 0.186)	(0.191, 3.296)		
Abdo-men	Low	Out-door	5	0.068	2.346	0.03
			(0.004, 0.211)	(0.282, 16.057)		
		High	4	0.020	1.824	0.01
			(0.010, 0.108)	(1.575, 3.582)		

Corresponding author: V. Corby-Harris,
vanessa.corby@usda.gov

Manuscript editor: James Nieh

Life stage	Tissue	Diet ¹	N	$\omega 6_{(\min, \max)}$	$\omega 3_{(\min, \max)}$	$\omega 6:\omega 3$
Queens	Whole	Out-door	3	0.254	4.996	0.05
			(0.108, 0.268)	(1.828, 10.259)		
		High	5	0.074	1.927	0.04
			(0.023, 0.081)	(0.829, 2.191)		
Larvae	Whole	Out-door	4	0.100	1.824	0.05
			(0.085, 0.113)	(1.722, 2.320)		
		Not measured	4	0.052	1.030	0.05
			(0.018, 8.221)	(0.340, 2.464)		
Pupae	Head	Low	4	0.040	1.456	0.03
			(0.001, 0.056)	(0.054, 1.944)		
		High	3	0.050	0.769	0.06
			(0.016, 0.186)	(0.191, 3.296)		
Abdo-men	Low	Out-door	5	0.068	2.346	0.03
			(0.004, 0.211)	(0.282, 16.057)		
		High	4	0.020	1.824	0.01
			(0.010, 0.108)	(1.575, 3.582)		

Life stage	Tissue	Diet ¹	N	ω_6 _(min, max)	ω_3 _(min, max)	$\omega_6:\omega_3$	Life stage	Tissue	Diet ¹	N	ω_6 _(min, max)	ω_3 _(min, max)	$\omega_6:\omega_3$
NEB	Head	Low	3	0.141 (0.029, 0.145)	2.116 (2.042, 2.934)	0.07	Forager	High	Out-door	4	0.875 (0.381, 0.943)	0.851 (0.302, 0.948)	1.03
				0.047 (0.006, 0.165)	0.658 (0.115, 0.910)	0.07				3	0.858 (0.435, 1.256)	1.280 (0.615, 1.612)	0.67
				0.014 (0.001, 0.048)	0.237 (0.022, 0.692)	0.06				4	0.084 (0.021, 0.261)	0.454 (0.183, 0.270)	0.18
				0.041 (0.028, 0.184)	0.446 (0.210, 1.409)	0.09				4	0.109 (0.052, 0.283)	0.415 (0.113, 0.176)	0.26
			5	0.012 (0.007, 0.364)	0.586 (0.207, 0.979)	0.02				3	0.109 (0.041, 0.387)	0.450 (0.412, 1.356)	0.24
	Abdo-men	Low	4	0.013 (0.004, 0.044)	0.445 (0.176, 0.763)	0.03	Abdo-men	High	Out-door	5	0.129 (0.012, 0.287)	0.302 (0.091, 0.355)	0.43
				0.242 (0.102, 0.540)	2.075 (1.222, 4.442)	0.12				4	0.311 (0.012, 0.652)	0.355 (0.013, 0.515)	0.88
			5	0.189 (0.058, 0.304)	0.662 (0.234, 1.700)	0.28				3	0.625 (0.336, 0.790)	0.603 (0.582, 0.895)	1.04
		Out-door	4	0.145 (0.069, 0.410)	0.351 (0.182, 1.518)	0.41				3			
				0.073 (0.067, 0.142)	0.302 (0.227, 0.821)	0.24				4			
	Abdo-men	Low	5	0.762 (0.218, 0.959)	0.884 (0.308, 1.164)	0.86				5			

¹Low = low-EFA diet, High = high-EFA diet

The original article has been corrected.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.