



Treatment of Binge-Eating Disorder Across the Lifespan: An Updated Review of the Literature and Considerations for Future Research

Kathryn E. Smith¹ · Andrea B. Goldschmidt²

Accepted: 8 February 2024
© The Author(s) 2024

Abstract

Purpose of Review The present review describes the recent literature on treatment for binge-eating disorder (BED) in adults and youth, with a particular focus on research gaps, emerging treatments, and future research directions.

Recent Findings Evidence supports the efficacy of several treatment modalities in adults, including self-help treatment, clinician-led psychotherapy, and pharmacotherapy; the largest effect sizes have been found for psychotherapies, most of which were cognitive-behavioral in orientation. Adapted psychotherapies for youth also show promise but lack a robust body of evidence. Predictors, moderators, and mediators of treatment outcome remain poorly understood; individuals with BED continue to experience significant barriers to treatment; and research is needed to address suboptimal treatment response. Recent work has highlighted the potential of adaptive interventions and investigation of novel mechanisms to address these gaps.

Summary Research on BED treatment continues to grow, though critical questions must be answered to improve treatment efficacy across the lifespan.

Keywords Binge-eating disorder · Treatment · Psychotherapy · Pharmacotherapy · Review · Efficacy

Introduction

Binge-eating disorder (BED) is characterized by recurrent episodes of binge eating (i.e., consuming an objectively large amount of food while experiencing a sense of loss of control over eating) in the absence of regular use of inappropriate compensatory behaviors intended to influence body weight [1]. Recent point prevalence estimates indicate BED affects ~1–4% of adults [2] and ~1% of youth [3] and is strongly associated with elevated body mass index (BMI; kg/m² [4]), excess weight gain [5–7], and physical [8, 9] and psychiatric comorbidities [10], including components of metabolic syndrome, gastrointestinal symptoms, mood and anxiety disorders, and sleep- and pain-related conditions. As such, BED presents significant burden to the individual and to society, costing consumers nearly \$20 billion

each year in direct healthcare expenditures, lost wages, and decreased productivity [11]. There are several efficacious interventions for BED in adults, including psychological, behavioral, pharmacological, and surgical treatment options [12], with fewer well-established (but nevertheless promising) treatment options for children and adolescents [13]. The purpose of this article is to provide an updated review on the treatment of BED across the lifespan, with special attention to new and emerging interventions and areas for future research.

Adult Treatment Approaches

Psychological Interventions

Several empirically supported treatment frameworks are available for adults with BED. First-line interventions for those with mild-to-moderate symptom severity include cognitive-behavioral therapy (CBT), which focuses primarily on establishing regular eating patterns and addressing maladaptive cognitions related to body shape and weight, the putative maintaining factors of binge-eating behaviors [14], and interpersonal psychotherapy (IPT), which focuses

✉ Kathryn E. Smith
ksmith41@usc.edu

¹ Department of Psychiatry and Behavioral Sciences, University of Southern California, 2250 Alcazar St #2200, Los Angeles, CA 90033, USA

² Department of Psychiatry, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA

on improving problematic social factors that drive negative affect and subsequent binge eating. Emerging research also supports the efficacy of integrative cognitive-affective therapy (ICAT), which targets momentary cognitive, behavioral, and emotional antecedents to binge eating, and adapted third-wave interventions such as dialectical behavior therapy (DBT), acceptance and commitment therapy (ACT), mindfulness-based interventions, and compassion-focused therapy [15•, 16–18]. While these interventions are typically delivered in individual or group-based formats, several approaches have been adapted into self-help and/or digital formats to improve access and reduce costs.

To date, CBT has been the most commonly studied intervention for BED with the largest body of supporting evidence, and as such, it forms the foundation for most currently available self-help interventions. The efficacy of self-help treatments has been supported by randomized controlled trials (RCTs) showing reductions in binge-eating symptoms and eating-related psychopathology compared to wait-list controls, with effects maintained up to 12-months post-treatment [15•, 19•]. Similarly, with respect to clinician-led psychotherapy, recent meta-analyses showed that compared to inactive control conditions, psychotherapy (consisting of mostly CBT) showed large effect sizes in reducing binge-eating episodes and eating disorder psychopathology at end-of-treatment and through 12-months post-treatment [15•, 18]. Notably, thus far, there remains limited evidence supporting the superiority of CBT compared to other active psychotherapies or treatment modalities [15•, 16]. However, some evidence suggests that clinician-led CBT may produce more rapid change compared to IPT and self-help treatment, and was associated with lower drop-out compared to self-help treatment; additionally, CBT was more efficacious in reducing binge eating compared to behavioral weight loss interventions [15•, 17, 20].

Pharmacological and Medical Interventions

For adults with moderate-to-severe BED symptoms, evidence also supports the efficacy of pharmacotherapy alone or in conjunction with psychotherapy [15•, 21]. Although no drug has been developed specifically for binge eating or BED, several psychotropic medications targeting other conditions have shown to also have beneficial effects on binge eating and have been re-purposed for BED treatment. Currently, the stimulant medication lisdexamfetamine (LDX) is the only FDA-approved medication in the United States for BED treatment, though antidepressants (particularly selective serotonin reuptake inhibitors) and antiepileptics (particularly topiramate) have also shown modest efficacy in reducing binge eating in BED [21]. In addition, although bariatric surgery is not considered a primary intervention for BED, most studies have found that bariatric surgery is

related to initial improvements in binge eating and BED, which are usually shown through two years post-surgery [22]. However, while binge eating improves in the short-term after surgery, this behavior may rebound over time, and the course of postoperative binge eating becomes more variable over longer term follow-ups [22]. The initial declines in binge eating could be due in part to anatomical changes that limit the capacity to objectively overeat following surgery, and some research suggests other problematic eating behaviors may emerge or worsen during this time, such as loss of control eating and grazing (i.e., repetitive, unplanned consumption of small quantities of food over time) [22].

Limitations and Research Gaps

Despite the variety of available evidence-based interventions, there remain critical limitations and unanswered questions in the treatment of BED in adults. Access to and affordability of treatment remains a significant barrier across eating disorder diagnoses [23], and the majority of individuals with eating disorders, including those with BED, do not seek out and/or engage in treatment [24]. Treatment is often delayed due to under-detection and under-diagnosis of BED by healthcare professionals, particularly in primary care settings where individuals are most likely to first seek help [24, 25]. Additionally, those with BED may not identify their symptoms as problematic due to limited mental health literacy about eating disorders, and individuals are more likely to seek help for physical rather than mental health complaints [26, 27]. These barriers may be exacerbated in BED due to stereotypes that eating disorders are limited to those who are underweight or engage in restriction or purging behaviors, as well as experiences of stigma [28, 29]. For instance, compared to individuals with anorexia nervosa or bulimia nervosa, those with BED are more likely to be the target of stigmatizing attributions such as lack of self-control, personal responsibility, and blameworthiness [30]. These beliefs may also be present among family, friends, and healthcare professionals of those with BED, further hindering the recognition of BED symptoms and likelihood that affected individuals will seek out treatment.

Even when individuals with BED seek out help, most do not receive specialized eating disorder treatment [24]. And although more efficacious than inactive control groups, the outcomes of existing specialized treatments are modest, with about half of individuals with BED achieving abstinence from binge eating at end-of-treatment through 12-months post-treatment [19•]. Notably, despite the effects on binge eating, psychotherapies for BED do not lead to consistent changes in body weight [15•, 19•]. A recent comprehensive meta-analysis of BED treatment outcomes across 81 RCTs (including self-help, psychotherapy, and pharmacotherapy interventions) also demonstrated high heterogeneity across

studies, low methodological quality, and a lack of long-term follow-up data examining durability of effects [15•]. In addition, there were few head-to-head comparisons between BED treatments, limiting understanding of comparative efficacy of interventions [15•].

Finally, little is known regarding predictors, moderators, and mediators of BED treatment outcomes, which are crucial in helping to identify which treatments work best for whom, as well as pinpointing underlying mechanisms of action. Consideration of heterogeneity in treatment response is imperative given earlier research showing individuals with less severe BED symptoms responded to placebo [31], suggesting that a subset of individuals could benefit from lower intensity treatments and may experience more a transient course of illness. With respect to predictors of treatment outcomes, across treatment modalities, meta-analytic results identified some patient characteristics (i.e., lower age and BMI, female sex, and higher baseline binge-eating episodes) as non-specific predictors that related to greater reduction in binge-eating episodes [15•]. Thus far, early behavior change (characterized by a 65–70% reduction in binge eating by week 4 of treatment) has been the only consistent predictor of CBT outcome [32]. With respect to moderators, patient demographics and psychiatric comorbidities have not consistently predicted response to CBT compared to other treatments for BED, and inconsistent findings have emerged regarding overvaluation of shape and weight as a differential predictor of CBT outcomes compared to other psychotherapies [32]. With respect to mindfulness- and acceptance-based treatments interventions for BED, a recent review showed that few studies assessed moderators of outcomes [33]. Regarding mediators, research has been limited and inconsistent across treatment modalities. A recent study demonstrated support for mechanisms implicated by interpersonal theory, showing that the relationship between decreases in interpersonal problems and binge-eating reduction was mediated by improvement in negative affect in both CBT and IPT [34]. In addition, the review of mindfulness- and acceptance-based treatments suggested that pre- to post-treatment improvements in theoretically consistent mechanisms (e.g., psychological flexibility, emotion regulation) were associated with better outcomes, though rigorous statistical tests of mediation were lacking [33].

Child and Adolescent Treatment Approaches

Psychological Interventions

The prevalence of full-syndrome BED is relatively low in children and adolescents, and data suggest that subclinical symptoms (e.g., binge-eating episodes that fail to meet size or frequency criteria required for a diagnosis of BED) are

strongly associated with distress, impairment, and risk for full-syndrome eating disorders [35, 36]. Thus, interventions for youth have typically relied upon age-adapted criteria for BED, which include both objective and subjective binge episodes and allow for lower frequency and/or limited duration of binge eating relative to DSM-5 criteria [37]. Most youth interventions have focused on developmentally tailoring interventions with demonstrated efficacy in the treatment of adults [38]. These include CBT [39–41], IPT [42], and DBT [43, 44]. CBT has been shown to outperform delayed treatment in youth with recurrent binge eating [41] and age-adapted BED [39], although currently there are no studies comparing it to a credible control treatment. IPT has been found to produce greater improvements in loss of control eating (with or without consumption of an objectively large amount of food) among adolescents than a health education control condition in both a small pilot RCT [45] and a larger, fully powered RCT [42]. Finally, both a case report [44] and a small RCT [43] suggest that a DBT-informed approach may hold promise as an acceptable and efficacious treatment for adolescent binge eating.

Pharmacological and Medical Interventions

Research on pharmacological agents for the treatment of binge-type eating in youth has been extremely limited, although off-label usage of LDX has been reported in the literature with promising effects on binge-type eating in a single case [46]. Similar to adult studies [23], bariatric surgery in adolescents is associated with improvements in binge-type eating from pre-surgery through up to six years post-surgery [47–49], although prevalence rates tend to show gradual increases with longer passage of time since surgery.

Limitations and Research Gaps

While CBT, IPT, and DBT all show promise in the treatment of binge eating-related conditions in children and adolescents, few RCTs have been conducted, and even fewer have included a rigorous time- and attention-matched control condition. Therefore, it is premature to declare a “first line” evidence-based approach for this age group. There are several additional limitations and gaps that impede scientific progress in understanding best practices for the treatment of binge eating-related conditions in youth.

First, current treatments for binge-type eating disorders in youth are based on etiological models that were developed for adults, and either have not been adequately tested in youth or have received limited support in youth. For example, restraint theory (on which CBT is based) proposes that rigid and overly restrictive dieting attitudes, cognitions, and behaviors lead to over-reliance on non-physiological cues (e.g., pre-defined portion sizes) to guide eating behavior, increasing risk for

dysregulated eating when dietary restraint is interrupted [e.g., through breaking a dietary rule; 50]. Yet, there is mixed support for elevated dietary restraint and objective disturbances in eating behavior in adolescents with binge-type eating [51–61]. Similarly, affect regulation theories (on which IPT is based) suggest that binge-type eating represents an attempt to modulate negative mood [e.g., via affective trade-off or masking; 62]. Yet, while there is strong support for negative affect as a distal risk factor for later development of binge-type eating in adolescents [63, 64], momentary assessment methodologies have yielded no evidence that negative affect is a proximal precipitant of binge-type eating in youth [65–67]. Relatedly, current interventions for youth fail to account for other factors that may independently or interactively contribute to dysregulated eating in youth, such as the food environment and individual responses to palatable foods.

Second, existing interventions fail to adequately consider that self-regulation capacity [i.e., ability to override undesirable impulses in pursuit of longer-term goals; 68]. Self-regulation subsumes the three core executive functions of inhibition, working memory, and cognitive flexibility, and lower order derivative processes including planning, problem-solving, affect regulation, etc. [69]. Executive functions remain in flux throughout childhood, late adolescence, and into early adulthood [70, 71], underscored by developmental changes in neurobiology [72]. This may make it challenging for adolescents to implement skills learned during treatment that largely depend on self-control in the moments and contexts they are needed most (e.g., remembering to employ stimulus control to prevent binge-type in high-risk locations).

Third, as with adult interventions for BED, treatments targeting youth can be costly to administer and may not be adequately covered by insurance [especially those delivered in face-to-face formats; 73], often involve intensive participation (e.g., 16–20 sessions over 6–12 months), typically require extensive training of providers that may not be widely available, and are frequently inaccessible outside of metropolitan areas and specialty treatment settings. Digital interventions may present an alternative to offset some of these limitations, but few have been rigorously tested [40]. Those that have been evaluated utilized outdated technology (e.g., required use of a desktop or laptop computer and were not designed for use via Smartphone or other ambulatory devices) and were not informed by the needs, expectations, and preferences of intended users which may limit engagement and clinical impact.

Future Directions and Conclusion

Although significant progress has been made over the past several decades in developing and evaluating interventions for BED across the lifespan, there are several major research

gaps that should be a focus of research in the coming years. First, although there are several efficacious interventions for BED in adults, with many of these approaches demonstrating efficacy in youth as well, outcomes are heterogeneous and a significant proportion of individuals who receive these interventions fail to respond optimally (e.g., continue to meet criteria for BED or engage in subclinical binge-eating symptoms after a course of treatment). Alternative treatment options for initial non-responders to first-line treatments are not well-characterized, nor are characteristics of individuals who are likely to have poor response to particular treatment modalities. Understanding for whom, and under what conditions, individuals are likely to respond to particular treatment modalities could inform personalized medicine approaches and help with allocation of limited therapeutic resources (e.g., using a lower-touch intervention for individuals likely to respond to a range of approaches while preserving more time- and resource-intensive interventions for those likely to have a more recalcitrant course of illness). Adaptive stepped care interventions and use of sequential multiple assignment randomized trial (SMART) designs will be particularly useful to address these questions and have already shown promise in BED treatment [74•]. Relatedly, mechanisms of action for existing treatments are not well characterized. Identifying the most active ingredients of treatments using multiphase optimization strategies could improve both efficacy and efficiency (e.g., by focusing on only the most potent intervention strategies, thereby reducing the overall length of treatment [75•]). It will also be important to understand how to effectively target such mechanisms in real life. To this end, just-in-time adaptive interventions may assist in personalizing delivery of intervention components to capitalize on the times and places those components will be most impactful [76•].

Next, as mentioned earlier, access to evidence-based treatments is largely limited to those with adequate resources (e.g., insurance coverage, reliable transportation to travel to and from treatment) and/or geographic proximity to specialty centers (which are mostly centered in densely population urban regions). One consequence of the COVID-19 pandemic has been the pivot to telehealth/virtual healthcare for most mental health conditions; however, acceptability and efficacy of telehealth and other virtual approaches (e.g., eHealth/mHealth, digital interventions) are not well characterized.

Finally, identifying adjunctive treatment components that target mechanisms outside of the current theoretical frameworks may help to improve outcomes and address comorbidities that frequently co-occur with BED. For example, given evidence that individuals with BED display diminished neurocognitive performance relative to their peers [77], executive function training has been recommended as a way to improve adherence to psychological interventions [78, 79], particularly

those focused on momentary processes that may require high degrees of self-regulation to implement in real-world contexts. While current treatments for BED do not address the role of physical activity, there is also evidence that interventions promoting physical activity could exert effects on binge eating, potentially by improving self-regulation and hedonic hunger. For example, emerging physical activity interventions targeting binge eating in adults were related to reduced binge-eating episodes as well as improvements in depression, anxiety, and food craving [80]. Similarly, understanding ways to optimally integrate weight control strategies into interventions for BED could help streamline treatment for those with comorbid obesity, as weight loss following psychological treatment for BED is typically modest or non-existent [81]. Furthermore, existing evidence has suggested potential of weight management interventions to reduce binge eating in both adult [82] and pediatric populations [83–85], though effects are typically less robust for binge eating outcomes than targeted psychological approaches such as CBT [86–88].

Taken together, there are several efficacious treatment options for those with BED, including self-help, clinician-led psychotherapy, and medical interventions, though the development and evaluation of interventions for youth with BED are more limited compared to adults. However, there remain critical barriers to treatment and gaps in knowledge that underscore the need for continued research, treatment development, and potentially refinement of existing theoretical models. This will likely involve investigation of novel biological, social, and environmental factors that have not been adequately considered in prior work. Moreover, it is imperative that future research consider developmental processes occurring in youth and identify if and how the etiology and maintenance of BED in youth may diverge from adults.

Author Contributions KES and ABG wrote the main manuscript text and reviewed the manuscript.

Funding Open access funding provided by SCEL, Statewide California Electronic Library Consortium

Data Availability No datasets were generated or analysed during the current study.

Compliance with Ethical Standards

Conflict of Interest The authors declare no competing interests.

Human and Animal Rights and Informed Consent This article does not contain any studies with human or animal subjects performed by any of the authors.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

Papers of particular interest, published recently, have been highlighted as:

- Of importance

1. American Psychiatric Association. Diagnostic and statistical manual of mental disorders (5th ed., text rev.) 2022. <https://doi.org/10.1176/appi.books.9780890425787>
2. Lindvall Dahlgren C, Wisting L, Rø Ø. Feeding and eating disorders in the DSM-5 era: a systematic review of prevalence rates in non-clinical male and female samples. *J Eat Disord.* 2017;5:56.
3. Kjeldbjerg ML, Clausen L. Prevalence of binge-eating disorder among children and adolescents: a systematic review and meta-analysis. *Eur Child Adolesc Psychiatry.* 2023;32(4):549–74.
4. Kessler RC, Berglund PA, Chiu WT, Deitz AC, Hudson JI, Shahly V, et al. The prevalence and correlates of binge eating disorder in the World Health Organization World Mental Health Surveys. *Biol Psychiatry.* 2013;73(9):904–14.
5. Barnes RD, Blomquist KK, Grilo CM. Exploring pretreatment weight trajectories in obese patients with binge eating disorder. *Compr Psychiatry.* 2011;52(3):312–8.
6. Blomquist KK, Barnes RD, White MA, Masheb RM, Morgan PT, Grilo CM. Exploring weight gain in year before treatment for binge eating disorder: a different context for interpreting limited weight losses in treatment studies. *Int J Eat Disord.* 2011;44(5):435–9.
7. Fairburn CG, Cooper Z, Doll HA, Norman P, O'Connor M. The natural course of bulimia nervosa and binge eating disorder in young women. *Arch Gen Psychiatry.* 2000;57(7):659–65.
8. Mitchell JE. Medical comorbidity and medical complications associated with binge-eating disorder. *Int J Eat Disord.* 2016;49(3):319–23.

9. Olguin P, Fuentes M, Gabler G, Guerdjikova AI, Keck PE Jr, McElroy SL. Medical comorbidity of binge eating disorder. *Eat Weight Disord.* 2017;22(1):13–26.
10. Hudson JI, Hiripi E, Pope HG Jr, Kessler RC. The prevalence and correlates of eating disorders in the National Comorbidity Survey Replication. *Biol Psychiatry.* 2007;61(3):348–58.
11. Streatfeild J, Hickson J, Austin SB, Hutcheson R, Kandel JS, Lampert JG, et al. Social and economic cost of eating disorders in the United States: evidence to inform policy action. *Int J Eat Disord.* 2021;54(5):851–68.
12. Grilo CM, Juarascio A. Binge-eating disorder interventions: review, current status, and implications. *Curr Obes Rep.* 2023 Sep;12(3):406–16. <https://doi.org/10.1007/s13679-023-00517-0>.
13. Bohon C. Binge eating disorder in children and adolescents. *Child Adolesc Psychiatr Clin N Am.* 2019;28(4):549–55.
14. Fairburn CG, Cooper Z, Shafran R. Cognitive behaviour therapy for eating disorders: A “transdiagnostic” theory and treatment. *Behav Res Ther.* 2003;41(5):509–28.
15. Hilbert A, Petroff D, Herpertz S, Pietrowsky R, Tuschen-Caffier B, Vocks S, et al. Meta-analysis of the efficacy of psychological and medical treatments for binge-eating disorder. *J Consult Clin Psychol.* 2019;87(1):91–105. **This comprehensive meta-analysis surveyed 81 randomized control trials of BED treatment spanning psychotherapy, self-help treatment, and pharmacotherapy. Results supported the efficacy of psychotherapies and self-help treatment compared to inactive controls, and for pharmacotherapy compared to placebo. Notably, there was limited support for the superiority of one treatment modality compared to other active treatments. Few patient characteristics were identified as non-specific predictors of outcomes.**
16. Linardon J, Fairburn CG, Fitzsimmons-Craft EE, Wilfley DE, Brennan L. The empirical status of the third-wave behaviour therapies for the treatment of eating disorders: A systematic review. *Clin Psychol Rev.* 2017;58:125–40.
17. Linardon J, Wade TD, de la Piedad GX, Brennan L. The efficacy of cognitive-behavioral therapy for eating disorders: a systematic review and meta-analysis. *J Consult Clin Psychol.* 2017;85(11):1080–94.
18. Peterson CB, Engel SG, Crosby RD, Strauman T, Smith TL, Klein M, et al. Comparing integrative cognitive-affective therapy and guided self-help cognitive-behavioral therapy to treat binge-eating disorder using standard and naturalistic momentary outcome measures: a randomized controlled trial. *Int J Eat Disord.* 2020;53(9):1418–27.
19. Hilbert A, Petroff D, Herpertz S, Pietrowsky R, Tuschen-Caffier B, Vocks S, et al. Meta-analysis on the long-term effectiveness of psychological and medical treatments for binge-eating disorder. *Int J Eat Disord.* 2020;53(9):1353–76. **This is the first meta-analysis synthesizing research on long-term effectiveness of BED treatments. Results of 114 studies supported the efficacy of psychotherapy and self-help treatment up to 12 months post-treatment, though longer follow-up data were limited.**
20. de Zwaan M, Herpertz S, Zipfel S, Svaldi J, Friederich HC, Schmidt F, et al. Effect of internet-based guided self-help vs individual face-to-face treatment on full or subsyndromal binge eating disorder in overweight or obese patients: the INTERBED randomized clinical trial. *JAMA Psychiatr.* 2017;74(10):987–95.
21. McElroy SL, Guerdjikova AI, Mori N, Romo-Nava F. Medication for binge eating. In: Frank GKW, Berner LA, editors. *Binge Eating: A Transdiagnostic Psychopathology.* Cham, Switzerland: Springer; 2020. p. 227–41.
22. Spirou D, Raman J, Smith E. Psychological outcomes following surgical and endoscopic bariatric procedures: a systematic review. *Obesity reviews : an official journal of the International Association for the Study of Obesity.* 2020;21(6):e12998.
23. Innes NT, Clough BA, Casey LM. Assessing treatment barriers in eating disorders: a systematic review. *Eat Disord.* 2017;25(1):1–21.
24. Bilić S, Sander J, Bauer S. Overcoming barriers to the treatment of binge eating. In: Frank GKW, Berner LA, editors. *Binge Eating: A Transdiagnostic Psychopathology.* Chem, Switzerland: Springer; 2020. p. 311–21.
25. Chao AM, Rajagopalan AV, Tronieri JS, Walsh O, Wadden TA. Identification of binge eating disorder criteria: results of a national survey of healthcare providers. *J Nurs Scholarsh.* 2019;51(4):399–407.
26. Striegel-Moore RH, Dohm FA, Kraemer HC, Schreiber GB, Crawford PB, Daniels SR. Health services use in women with a history of bulimia nervosa or binge eating disorder. *Int J Eat Disord.* 2005;37(1):11–8.
27. Grillot CL, Keel PK. Barriers to seeking treatment for eating disorders: the role of self-recognition in understanding gender disparities in who seeks help. *Int J Eat Disord.* 2018;51(11):1285–9.
28. Barnes RD, Ivezaj V, Grilo CM. An examination of weight bias among treatment-seeking obese patients with and without binge eating disorder. *Gen Hosp Psychiatry.* 2014;36(2):177–80.
29. Beat. The cost of eating disorders; social, health and economic impacts. 2015. Available at: <https://www.beateatingdisorders.org/uploads/documents/2017/10/the-costs-of-eating-disorders-final-original.pdf>.
30. Puhl R, Suh Y. Stigma and eating and weight disorders. *Curr Psychiatry Rep.* 2015;17:1–10.
31. Jacobs-Pilipski MJ, Wilfley DE, Crow SJ, Walsh BT, Lilienfeld LR, West DS, et al. Placebo response in binge eating disorder. *Int J Eat Disord.* 2007;40(3):204–11.
32. Linardon J, de la Piedad GX, Brennan L. Predictors, moderators, and mediators of treatment outcome following manualised cognitive-behavioural therapy for eating disorders: a systematic review. *Eur Eat Disord Rev.* 2017;25(1):3–12.
33. Barney JL, Murray HB, Manasse SM, Dochat C, Juarascio AS. Mechanisms and moderators in mindfulness- and acceptance-based treatments for binge eating spectrum disorders: a systematic review. *Eur Eat Disord Rev.* 2019;27(4):352–80.
34. Karam AM, Eichen DM, Fitzsimmons-Craft EE, Wilfley DE. An examination of the interpersonal model of binge eating over the course of treatment. *Eur Eat Disord Rev.* 2020;28(1):66–78.
35. Hilbert A, Brauhardt A. Childhood loss of control eating over five-year follow-up. *Int J Eat Disord.* 2014;47(7):758–61.
36. Hilbert A, Hartmann AS, Czaja J, Schoebi D. Natural course of preadolescent loss of control eating. *J Abnorm Psychol.* 2013;122(3):684–93.
37. Schmidt R, Hilbert A. Predictors of symptom trajectories after cognitive-behavioral therapy in adolescents with an age-adapted diagnosis of binge-eating disorder. *Behav Ther.* 2022;53(1):137–49.
38. Byrne ME, LeMay-Russell S, Tanofsky-Kraff M. Loss-of-control eating and obesity among children and adolescents. *Curr Obes Rep.* 2019 Mar;8(1):33–42. <https://doi.org/10.1007/s13679-019-0327-1>.
39. Hilbert A, Petroff D, Neuhaus P, Schmidt R. Cognitive-behavioral therapy for adolescents with an age-adapted diagnosis of binge-eating disorder: a randomized clinical trial. *Psychother Psychosom.* 2020;89(1):51–3.
40. Jones M, Luce KH, Osborne MI, Taylor K, Cunning D, Doyle AC, et al. Randomized, controlled trial of an internet-facilitated intervention for reducing binge eating and overweight in adolescents. *Pediatrics.* 2008;121(3):453–62.
41. DeBar LL, Wilson GT, Yarborough BJ, Burns B, Oyler B, Hildebrandt T, et al. Cognitive behavioral treatment for recurrent binge eating in adolescent girls: a pilot trial. *Cogn Behav Pract.* 2013;20(2):147–61.

42. Tanofsky-Kraff M, Shomaker LB, Wilfley DE, Young JF, Sbrocco T, Stephens M, et al. Targeted prevention of excess weight gain and eating disorders in high-risk adolescent girls: a randomized controlled trial. *Am J Clin Nutr*. 2014;100(4):1010–8.
43. Mazzeo SE, Lydecker J, Harney M, Palmberg AA, Kelly NR, Gow RW, et al. Development and preliminary effectiveness of an innovative treatment for binge eating in racially diverse adolescent girls. *Eat Behav*. 2016;22:199–205.
44. Safer DL, Lock J, Couturier JL. Dialectical behavior therapy modified for adolescent binge eating disorder: A case report. *Cogn Behav Pract*. 2007;14(2):157–67.
45. Tanofsky-Kraff M, Wilfley DE, Young JF, Mufson L, Yanovski SZ, Glasofer DR, et al. A pilot study of interpersonal psychotherapy for preventing excess weight gain in adolescent girls at-risk for obesity. *Int J Eat Disord*. 2010;43(8):701–6.
46. Srivastava G, O'Hara V, Browne N. Use of lisdexamfetamine to treat obesity in an adolescent with severe obesity and binge eating. *Children (Basel)*. 2019 Feb 4;6(2):22. <https://doi.org/10.3390/children6020022>.
47. Järholm K, Bruze G, Peltonen M, Marcus C, Flodmark CE, Henfridsson P, et al. 5-year mental health and eating pattern outcomes following bariatric surgery in adolescents: a prospective cohort study. *Lancet Child Adolesc Health*. 2020;4(3):210–9.
48. Reddy SD, Sysko R, Mackey ER. Mental health, loss of control eating, and cognitive functioning in adolescents undergoing bariatric surgery. *Semin Pediatr Surg*. 2020;29(1):150892.
49. Goldschmidt AB, Khoury J, Jenkins TM, Bond DS, Thomas JG, Utzinger LM, et al. Adolescent loss-of-control eating and weight loss maintenance after bariatric surgery. *Pediatrics*. 2018 Jan;141(1):e20171659. <https://doi.org/10.1542/peds.2017-1659>.
50. Schaumberg K, Anderson DA, Anderson LM, Reilly EE, Gorrell S. Dietary restraint: what's the harm? A review of the relationship between dietary restraint, weight trajectory and the development of eating pathology. *Clin Obes*. 2016;6(2):89–100.
51. Goossens L, Braet C, Decaluwe V. Loss of control over eating in obese youngsters. *Behav Res Ther*. 2007;45(1):1–9.
52. Goossens L, Soenens B, Braet C. Prevalence and characteristics of binge eating in an adolescent community sample. *J Clin Child Adolesc Psychol*. 2009;38(3):342–53.
53. Claus L, Braet C, Decaluwe V. Dieting history in obese youngsters with and without disordered eating. *Int J Eat Disord*. 2006;39(8):721–8.
54. Decaluwe V, Braet C. Prevalence of binge-eating disorder in obese children and adolescents seeking weight-loss treatment. *Int J Obes Relat Metab Disord*. 2003;27(3):404–9.
55. Decaluwe V, Braet C. The cognitive behavioural model for eating disorders: a direct evaluation in children and adolescents with obesity. *Eat Behav*. 2005;6(3):211–20.
56. Decaluwe V, Braet C, Fairburn CG. Binge eating in obese children and adolescents. *Int J Eat Disord*. 2003;33(1):78–84.
57. Glasofer DR, Tanofsky-Kraff M, Eddy KT, Yanovski SZ, Theim KR, Mirch MC, et al. Binge eating in overweight treatment-seeking adolescents. *J Pediatr Psychol*. 2007;32(1):95–105.
58. Tanofsky-Kraff M, Faden D, Yanovski SZ, Wilfley DE, Yanovski JA. The perceived onset of dieting and loss of control eating behaviors in overweight children. *Int J Eat Disord*. 2005;38(2):112–22.
59. Tanofsky-Kraff M, Goossens L, Eddy KT, Ringham R, Goldschmidt A, Yanovski SZ, et al. A multisite investigation of binge eating behaviors in children and adolescents. *J Consult Clin Psychol*. 2007;75(6):901–13.
60. Tanofsky-Kraff M, McDuffie JR, Yanovski SZ, Kozlosky M, Schvey NA, Shomaker LB, et al. Laboratory assessment of the food intake of children and adolescents with loss of control eating. *Am J Clin Nutr*. 2009;89(3):738–45.
61. Tanofsky-Kraff M, Yanovski SZ, Wilfley DE, Marmarosh C, Morgan CM, Yanovski JA. Eating-disordered behaviors, body fat, and psychopathology in overweight and normal-weight children. *J Consult Clin Psychol*. 2004;72(1):53–61.
62. Haedt-Matt AA, Keel PK. Revisiting the affect regulation model of binge eating: a meta-analysis of studies using ecological momentary assessment. *Psychol Bull*. 2011;137(4):660–81.
63. Goldschmidt AB, Wall MM, Zhang J, Loth KA, Neumark-Sztainer D. Overeating and binge eating in emerging adulthood: 10-year stability and risk factors. *Dev Psychol*. 2016;52(3):475–83.
64. Stice E. Risk and maintenance factors for eating pathology: a meta-analytic review. *Psychol Bull*. 2002;128(5):825–48.
65. Ranzenhofer LM, Engel SG, Crosby RD, Anderson M, Vannucci A, Cohen LA, et al. Using ecological momentary assessment to examine interpersonal and affective predictors of loss of control eating in adolescent girls. *Int J Eat Disord*. 2014;47(7):748–57.
66. Hilbert A, Rief W, Tuschen-Caffier B, de Zwaan M, Czaja J. Loss of control eating and psychological maintenance in children: an ecological momentary assessment study. *Behav Res Ther*. 2009;47(1):26–33.
67. Goldschmidt AB, Smith KE, Crosby RD, Boyd HK, Dougherty E, Engel SG, et al. Ecological momentary assessment of maladaptive eating in children and adolescents with overweight or obesity. *Int J Eat Disord*. 2018;51(6):549–57.
68. Johnson F, Pratt M, Wardle J. Dietary restraint and self-regulation in eating behavior. *Int J Obes*. 2012;36(5):665–74.
69. Hofmann W, Schmeichel BJ, Baddeley AD. Executive functions and self-regulation. *Trends Cogn Sci*. 2012;16(3):174–80.
70. Best JR, Miller PH. A developmental perspective on executive function. *Child Dev*. 2010;81(6):1641–60.
71. Ferguson HJ, Brunson VEA, Bradford EEF. The developmental trajectories of executive function from adolescence to old age. *Sci Rep*. 2021;11(1):1382.
72. Casey BJ. Beyond simple models of self-control to circuit-based accounts of adolescent behavior. *Annu Rev Psychol*. 2015;66:295–319.
73. Weissman RS, Rosselli F. Reducing the burden of suffering from eating disorders: unmet treatment needs, cost of illness, and the quest for cost-effectiveness. *Behav Res Ther*. 2017;88:49–64.
74. Grilo CM, White MA, Masheb RM, Ivezaj V, Morgan PT, Georgieva R. Randomized controlled trial testing the effectiveness of adaptive “SMART” stepped-care treatment for adults with binge-eating disorder comorbid with obesity. *Am Psychol*. 2020;75(2):204–18. **This study describes the first use of a sequential multiple assignment randomized trial (SMART) design in the treatment of BED.**
75. Manasse SM, Clark KE, Juarascio AS, Forman EM. Developing more efficient, effective, and disseminable treatments for eating disorders: an overview of the multiphase optimization strategy. *Eat Weight Disord*. 2019;24(6):983–95. **This article describes the multiphase optimization strategy and potential applications to eating disorder interventions.**
76. Juarascio AS, Parker MN, Lagacey MA, Godfrey KM. Just-in-time adaptive interventions: a novel approach for enhancing skill utilization and acquisition in cognitive behavioral therapy for eating disorders. *Int J Eat Disord*. 2018;51(8):826–30. **This article reviews the methodology of just-in-time adaptive interventions and relevance to eating disorder treatment.**
77. Iceta S, Rodrigue C, Legendre M, Daoust J, Flaudias V, Michaud A, et al. Cognitive function in binge eating disorder and food addiction: a systematic review and three-level meta-analysis. *Prog Neuropsychopharmacol Biol Psychiatry*. 2021;111:110400.
78. Eichen DM, Strong DR, Twamley EW, Boutelle KN. Adding executive function training to cognitive behavioral therapy for binge eating disorder: A pilot randomized controlled trial. *Eat Behav*. 2023;51:101806.

79. Eichen DM, Matheson BE, Appleton-Knapp SL, Boutelle KN. Neurocognitive treatments for eating disorders and obesity. *Curr Psychiatry Rep.* 2017;19(9):62.
80. Blanchet C, Mathieu M, St-Laurent A, Fecteau S, St-Amour N, Drapeau V. A systematic review of physical activity interventions in individuals with binge eating disorders. *Curr Obes Rep.* 2018;7(1):76–88.
81. Palavras MA, Hay P, Filho CA, Claudino A. The efficacy of psychological therapies in reducing weight and binge eating in people with bulimia nervosa and binge eating disorder who are overweight or obese—a critical synthesis and meta-analysis. *Nutrients.* 2017 Mar 17;9(3):299. <https://doi.org/10.3390/nu9030299>.
82. Jebeile H, Libesman S, Melville H, Low-Wah T, Dammery G, Seidler AL, et al. Eating disorder risk during behavioral weight management in adults with overweight or obesity: a systematic review with meta-analysis. *Obesity reviews : an official journal of the International Association for the Study of Obesity.* 2023;24(6):e13561.
83. Jebeile H, Gow ML, Baur LA, Garnett SP, Paxton SJ, Lister NB. Treatment of obesity, with a dietary component, and eating disorder risk in children and adolescents: a systematic review with meta-analysis. *Obesity reviews : an official journal of the International Association for the Study of Obesity.* 2019;20(9):1287–98.
84. Moustafa AF, Quigley KM, Wadden TA, Berkowitz RI, Chao AM. A systematic review of binge eating, loss of control eating, and weight loss in children and adolescents. *Obesity.* 2021;29(8):1259–71.
85. Peckmezian T, Hay P. A systematic review and narrative synthesis of interventions for uncomplicated obesity: weight loss, well-being and impact on eating disorders. *J Eat Disord.* 2017;5:15.
86. Grilo CM, Masheb RM. A randomized controlled comparison of guided self-help cognitive behavioral therapy and behavioral weight loss for binge eating disorder. *Behav Res Ther.* 2005;43(11):1509–25.
87. Grilo CM, Masheb RM, Wilson GT, Gueorguieva R, White MA. Cognitive-behavioral therapy, behavioral weight loss, and sequential treatment for obese patients with binge-eating disorder: a randomized controlled trial. *J Consult Clin Psychol.* 2011;79(5):675–85. <https://doi.org/10.1037/a0025049>.
88. Wilson GT, Wilfley DE, Agras WS, Bryson SW. Psychological treatments of binge eating disorder. *Arch Gen Psychiatry.* 2010;67(1):94–101.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.