



Correction to: Multidisciplinary treatment for locally advanced breast cancer with internal mammary lymph node metastasis in an elderly patient

Haruko Takuwa¹ · Wakako Tsuji¹ · Yoshihiro Yamamoto² · Chikako Yamauchi³ · Fumiaki Yotsumoto¹

Published online: 3 September 2019
© The Japan Society of Clinical Oncology 2019

Correction to:
International Cancer Conference Journal (2019) 8:1–6
<https://doi.org/10.1007/s13691-018-0344-z>

In the original publication, in Table 1, the patient 3, 58 years of age, was scheduled to be administered eribulin after

gemcitabine as adjuvant therapy. However, the patient was deceased before administration due to sepsis. Eribulin was not administered actually and it needs to be deleted. The corrected Table 1 is given in this Correction.

The original article can be found online at <https://doi.org/10.1007/s13691-018-0344-z>.

✉ Haruko Takuwa
st24057@yahoo.co.jp

¹ Department of Breast Surgery, Shiga General Hospital, 5-4-30, Moriyama-shi, Moriyama, Shiga 524-8524, Japan

² Department of Pathology, Shiga General Hospital, Moriyama, Shiga, Japan

³ Department of Radiation Oncology, Shiga General Hospital, Moriyama, Shiga, Japan

Table 1 Patient characteristics with internal mammary lymph node metastasis at diagnosis

Age	Clinical stage	Location	Subtype	Neoadjuvant therapy	Operation	Radiation	Adjuvant therapy	Outcome
42	T3N3bM0	Upper-lateral	Luminal-HER2	EC-wPTX + HCPT	Bp + SLNB	60 Gy/30fr	HCPT + TAM	SD
44	T4bN3bM1 (synchronous bilateral)	Upper-mediolateral	Luminal-HER2	TC	Bt	–	TAM + zometa-DTX + Per + HCPT + zometa	SD
58	T4bN2bM0 (SCC)	Upper-medial	Luminal A-like	EC-wPTX-VNR	Bp + Ax	50 Gy/25fr	DTX-TS-1 + ANA-GEM	Death due to sepsis
68	T2N3bM0	Lower-medial	Luminal B-like	wPTX	Bp + Ax	50 Gy/25fr	LET	SD
82	T2N3bM0	Upper-medial	Luminal B-like	–	Bp + Ax	60 Gy/30fr	UFT + LET	SD

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.