

## Erratum to: Diuretics-assisted treatment of chronic laxative abuse

Yuichi Shirasawa · Michio Fukuda ·  
Genjiro Kimura

Published online: 25 May 2014  
© Japanese Society of Nephrology 2014

**Erratum to: CEN Case Rep**  
**DOI 10.1007/s13730-014-0120-7**

As the result of a change that occurred during editorial processing, Fig. 1b appeared incorrectly in the article cited above. The correct figure is shown here. The publisher sincerely apologizes for the error.


---

The online version of the original article can be found under  
doi:[10.1007/s13730-014-0120-7](https://doi.org/10.1007/s13730-014-0120-7).

---

Y. Shirasawa (✉) · M. Fukuda  
Department of Cardio-Renal Medicine and Hypertension,  
Nagoya City University School of Medicine, 1 Kawasumi,  
Mizuho-Cho, Mizuho Ward, Nagoya City,  
Aichi Prefecture 467-8601, Japan  
e-mail: [yuiville@med.nagoya-cu.ac.jp](mailto:yuiville@med.nagoya-cu.ac.jp)

G. Kimura  
Asahi Rosai Hospital, Japan Labour Health and Welfare  
Organization, 61 Hirakocho-Kita, Owariasahi City,  
Aichi Prefecture 488-8585, Japan



**Fig. 1** Temporal relationships of body weight and urine volume (a), blood pressure (b), and hematocrit and total protein and albumin levels (c) during the hospitalization