Correction to: Combined light chain crystalline tubulopathy, podocytopathy, and histiocytosis associated with Bence–Jones κ protein diagnosed via immuno-electron microscopy

Yoshihiro Nakamura¹ · Hiroshi Kitamura² · Hiroki Ikai¹ · Mari Yamamoto¹ · Yukari Murai³ · Tsuyoshi Watanabe¹ · Naoho Takizawa¹ · Waka Yokoyama-Kokuryo¹ · Takashi Ehara⁴ · Masaya Watarai⁵ · Hideaki Shimizu⁶ · Yoshiro Fujita^{1,3}

Published online: 19 March 2021
© Japanese Society of Nephrology 2021

Correction to: CEN Case Reports
<https://doi.org/10.1007/s13730-021-00588-9>

In the original publication of the article, the email address of the corresponding author has been published incorrectly as nakamurashift7@yahoo.co.jp. The correct email address is nakamurashift@yahoo.co.jp.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s13730-021-00588-9>.

✉ Yoshihiro Nakamura
nakamurashift@yahoo.co.jp

- ¹ Department of Rheumatology, Chubu Rosai Hospital, 2-10-15, Komei-cho, Minato-ku, Nagoya, Aichi 455-8530, Japan
- ² Department of Pathology, National Hospital Organization Chiba-East Hospital, Chiba, Japan
- ³ Department of Nephrology, Chubu Rosai Hospital, 2-10-15, Komei-cho, Minato-ku, Nagoya, Aichi 455-8530, Japan
- ⁴ Department of Histopathology, Shinshu University School of Medicine, Matsumoto, Japan
- ⁵ Department of Hematology, Daido Hospital, Nagoya, Japan
- ⁶ Department of Nephrology and Renal Replacement, Daido Hospital, Nagoya, Japan