

## Erratum to: Mechanical endovascular thrombectomy for acute ischemic stroke: a retrospective multicenter study in Belgium

Niels Fockaert<sup>1</sup> · Marieke Coninckx<sup>2</sup> · Sam Heye<sup>3</sup> · Luc Defreyne<sup>4</sup> ·  
Denis Brisbois<sup>5</sup> · Pierre Goffette<sup>6</sup> · Jan Gralla<sup>7</sup> · Pasquale Mordasini<sup>7</sup> ·  
Andre Peeters<sup>8</sup> · Philippe Desfontaines<sup>9</sup> · Dimitri Hemelsoet<sup>2</sup> · Vincent Thijs<sup>1,10,11</sup> ·  
Robin Lemmens<sup>1,10,11</sup>

Published online: 4 November 2015  
© Belgian Neurological Society 2015

**Erratum to: Acta Neurol Belg**  
**DOI 10.1007/s13760-015-0552-7**

In the online version of the original article, Figs. 1 and 2 captions have been erroneously switched. The correct version of the figures and their captions are given in the following page.

---

The online version of the original article can be found under  
doi:[10.1007/s13760-015-0552-7](https://doi.org/10.1007/s13760-015-0552-7).

---

✉ Robin Lemmens  
robin.lemmens@uzleuven.be

<sup>1</sup> Department of Neurology, University Hospitals Leuven, 3000 Leuven, Belgium

<sup>2</sup> Department of Neurology, University Hospitals Gent, 9000 Ghent, Belgium

<sup>3</sup> Department of Radiology, University Hospitals Leuven, 3000 Leuven, Belgium

<sup>4</sup> Department of Vascular and Interventional Radiology, University Hospitals Gent, 9000 Ghent, Belgium

<sup>5</sup> Department of Interventional Neuroradiology, Centre Hospitalier Chrétien, 4000 Liège, Belgium

<sup>6</sup> Department of Neuroradiology, Cliniques Universitaires Saint-Luc, Université catholique de Louvain, Brussels, Belgium

<sup>7</sup> Department of Diagnostic and Interventional Neuroradiology, University Hospital Bern, Bern, Switzerland



<sup>8</sup> Department of Neurology, Cliniques Universitaires Saint-Luc, Université catholique de Louvain, 1200 Brussels, Belgium

<sup>9</sup> Department of Neurology, Centre Hospitalier Chrétien, 4000 Liège, Belgium

<sup>10</sup> Department of Neurosciences, Experimental Neurology and Leuven Research Institute for Neuroscience and Disease (LIND), KU Leuven, University of Leuven, 3000 Leuven, Belgium

<sup>11</sup> VIB, Vesalius Research Center, Laboratory of Neurobiology, 3000 Leuven, Belgium

**Fig. 1** Distribution of post-procedure mTICI scores, stratified by site of AOL. Substantial reperfusion, defined as an mTICI 2B-3, was achieved in 78 % of patients in the overall patient series; in 73 % with an occlusion in the anterior circulation versus 100 % with an posterior circulation occlusion ( $p = 0.02$ ). *ICA* internal carotid artery, *MCA* medial cerebral artery, *BA* basilar artery, *AOL* arterial occlusive lesion, *mTICI* modified thrombolysis in cerebral infarction



**Fig. 2** Functional outcome in all patients, stratified by AOL. In the overall population, GFO was achieved in 42 % of patients and EFO in 26 % EFO. When stratified by anterior versus posterior circulation stroke, no differences were observed: The rate of GFO was 44 % in patients with anterior circulation stroke versus 34 % in patients with

basilar artery stroke. *ICA* internal carotid artery, *MCA* medial cerebral artery, *BA* basilar artery, *mRS* modified Rankin Scale, *GFO* good functional outcome (mRS 0–2), *EFO* excellent functional outcome (mRS 0–1)