

## Erratum to: Subacute onset Hirayama disease with extensive myelopathy on MRI in flexed posture

Merel Brems<sup>1</sup> · Johan Van Goethem<sup>1</sup> · Mania De Praeter<sup>1</sup> · Barbara Willekens<sup>1,2</sup>

Published online: 20 May 2017  
© Belgian Neurological Society 2017

**Erratum to: Acta Neurol Belg**  
**DOI 10.1007/s13760-017-0782-y**

Unfortunately, Figs. 1 and 2 were deleted in the original publication. Figures 1 and 2 are given below.

The original version is also updated.

---

The online version of the original article can be found under doi:[10.1007/s13760-017-0782-y](https://doi.org/10.1007/s13760-017-0782-y).

---

✉ Merel Brems  
merel.brems@gmail.com

<sup>1</sup> Universitair Ziekenhuis Antwerpen, Antwerp, Belgium

<sup>2</sup> Laboratory of Experimental Hematology, University of Antwerp, Antwerp, Belgium



**Fig. 1** A midsagittal T2-weighted MR-image shows focal medullary atrophy over the lower cervical segment from C4 to C7 with accompanying intramedullary T2 hyperintense signal

**Fig. 2** **a** Midsagittal T2 and **b** STIR MR-images in flexion. These images show severe spinal narrowing due to compression by an anterior displaced posterior dural lining with a secondary broadened posterior epidural space

