Correction to: HACE1, GLRX5, and ELP2 gene variant cause spastic paraplegies

Gunes Sager¹ · Ayberk Turkeyilmaz² · Esra Arslan Ates³ · Busra Kutlubay⁴

© Belgian Neurological Society 2021

Correction to: Acta Neurologica Belgica
<https://doi.org/10.1007/s13760-021-01649-7>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgement Patients with FA2H gene and HACE1 gene variants were presented as a case report at the 20th Turkish National Pediatric Neurology Symposium on May 2018 (Hereditär Spastik Paraplegie hastalık Ailesine Olgularla Yaklaşım, Gunes Sager, Gülten Öztürk, Ayberk Turkeyilmaz, Pınar Ata, Olcay Ünver, Dilsad Türkdogan P-050, ref no:313, 20th Turkish National Pediatric Neurology Symposium 2018). We gratefully thank Prof. Dr. Dilsad Türkdogan, Ass. Prof. Dr. Olcay Ünver and Dr. Gülten Öztürk for their contributions.

The original article can be found online at <https://doi.org/10.1007/s13760-021-01649-7>.

✉ Gunes Sager
sgunessenturk@gmail.com

¹ Department of Pediatric Neurology, Kartal Dr. Lutfi Kirdar City Hospital, Semsî Denizer Avenue, Cevizli, 34890 Kartal, Istanbul, Turkey

² Department of Medical Genetics, Erzurum City Hospital, Erzurum, Turkey

³ Department of Medical Genetics, Marmara University Pendik Training and Research Hospital, Istanbul, Turkey

⁴ Department of Pediatric Neurology, Umraniye Training and Research Hospital, Istanbul, Turkey