



Correction to: Anaerobic production of valeric acid from crude glycerol via chain elongation

S. T. S. Veras¹ · W. A. Cavalcante² · T. A. Gehring³ · A. R. Ribeiro⁴ · T. J. T. Ferreira⁵ · M. T. Kato¹ · P. Rojas-Ojeda⁶ · J. L. Sanz-Martin⁶ · R. C. Leitão⁷

Published online: 11 November 2019
© Islamic Azad University (IAU) 2019

Correction to:
International Journal of Environmental Science and Technology
<https://doi.org/10.1007/s13762-019-02562-6>

The original version of this article unfortunately contained a mistake in Fig. 2. The corrected figure is given below.

The original article can be found online at <https://doi.org/10.1007/s13762-019-02562-6>.

✉ R. C. Leitão
renato.leitao@embrapa.br

¹ Laboratory of Environmental Sanitation, Department of Civil and Environmental Engineering, Federal University of Pernambuco, Av. Acadêmico Hélio Ramos s/n, Cidade Universitária, Recife, PE 50740-530, Brazil

² Federal Institute of Science and Technology of Ceará State, Campus Tauá, Rua Antônio Teixeira Benevides 01, Tauá, CE 63660-000, Brazil

³ Institute of Urban Water Management and Environmental Engineering, Ruhr-University Bochum, Universitätsstrasse 150, 44801 Bochum, Germany

⁴ São Carlos School of Engineering, University of São Paulo, Av. Trabalhador São Carlense, 400, São Carlos, SP 13566-590, Brazil

⁵ Institute of Marine Science, Federal University of Ceará, Avenida da Abolição, 3207, Fortaleza, CE 60165-081, Brazil

⁶ Department of Molecular Biology, Autonomous University of Madrid, 28049 Madrid, Spain

⁷ Embrapa Agroindústria Tropical, Rua Dra. Sara Mesquita, 2270, Fortaleza, CE 60511-110, Brazil






Fig. 2 General distribution of carboxylates (a) and (b), ethanol and 1,3-propanediol (c) concentrations throughout the three operational phases (P1), (P2), and (P3)

