

Concurrent dengue virus and Japanese encephalitis virus infection

B. Joob · V. Wiwanitkit

Received: 18 June 2012 / Accepted: 21 July 2012 / Published online: 11 August 2012
© Springer-Verlag 2012

Sir,

The recent article on concurrent dengue virus and Japanese encephalitis virus infection is very interesting [1]. Indeed, the condition can be possible and this can be either co-infection or co-detection. The condition can be possible only if there are both pathogens carried by the required mosquito vectors. Also, the patients have to be bitten by the mosquito vectors of both diseases. The possibility might be expected. However, to fulfill the whole condition, it might be more difficult than a single infection. An interesting topic to be discussed is the accuracy of the diagnosis of the infection. The cross-reactivity can be expected in classical immunology diagnosis [2].

Conflict of interest None.

References

1. Garg RK, Malhotra HS, Gupta A, Kumar N, Jain A. Concurrent dengue virus and Japanese encephalitis virus infection of the brain: is it co-infection or co-detection? *Infection*. 14 June 2012. (Epub ahead of print).
2. Yabe S, Nakayama M, Yamada K, Kitano T, Arai Y, Horimoto T, Masuda G, Mitou A, Tashiro M. Laboratory virological diagnosis of imported dengue cases. *Kansenshogaku Zasshi*. 1996;70: 1160–9.

B. Joob (✉)
Sanitation 1 Medical Academic Center, Bangkok, Thailand
e-mail: beuyjoob@hotmail.com

V. Wiwanitkit
Wiwanitkit House, Bang Khae, Bangkok, Thailand

V. Wiwanitkit
Hainan Medical University (HMU), Haikou, China

V. Wiwanitkit
Joseph Ayo Babalola University, Ikeji-Arakeji, Nigeria