

Geometric models of twisted differential K -theory I

Byungdo Park¹

Received: 10 February 2016 / Accepted: 10 April 2017 / Published online: 13 May 2017
© Tbilisi Centre for Mathematical Sciences 2017

Abstract This is the first in a series of papers constructing geometric models of twisted differential K -theory. In this paper we construct a model of even twisted differential K -theory when the underlying topological twist represents a torsion class. By differential twists we will mean smooth $U(1)$ -gerbes with connection, and we use twisted vector bundles with connection as cocycles. The model we construct satisfies the axioms of Kahle and Valentino, including functoriality, naturality of twists, and the hexagon diagram. This paper confirms a long-standing hypothetical idea that twisted vector bundles with connection define twisted differential K -theory.

Keywords Twisted K -theory · Differential K -theory · Twisted Chern character

Mathematics Subject Classification Primary 19L50; Secondary 19L10

1 Introduction

There has been a considerable interest in twisted and differential refinements of generalized cohomology theories. Several aspects of topology, geometry, analysis, and physics coalesce in this area, and it has provided interesting applications of

Communicated by Thomas Schick.

We gratefully acknowledge a partial support on this work from Mahmoud Zeinalian's NSF Grant DMS-1309099 as well as support and hospitality from the Hausdorff Research Institute for Mathematics and Max Planck Institute for Mathematics during our visit.

✉ Byungdo Park
byungdpark@gmail.com

¹ Department of Mathematics, CUNY Graduate Center, 365 Fifth Ave., New York, NY 10016, USA

∞ -categorical machinery nicely packaged by ∞ -sheaves of spectra on the site of smooth manifolds. (See [5, 6] for example.)

One important question in generalized cohomology theories is whether one can represent an element of a given generalized cohomology theory of a space using geometric cocycles. For instance, an element of the singular cohomology group of a space can be represented by a singular cocycle in the space, and elements in the complex K -theory of a space can be represented by complex vector bundles over that space. However, geometric models are still unknown for most other cohomology theories such as topological modular forms, and even less is known for their twisted and differential refinements. This paper is the first in a series of papers to answer this question for the case of differential refinements of twisted complex K -theory.

Twisted K -theory was first introduced by Donovan and Karoubi in [10], where twists represent torsion classes in degree 3 integral cohomology, and Rosenberg [23] for all classes. More recently twisted K -theory has received much attention because of its applications in classifying D-brane charges in string theory [27], Verlinde algebras [12], and topological insulators [14].

An archetype of differential K -theory first appeared in Karoubi [18] as the *multiplicative K -theory*. This is nowadays known as the flat subgroup of the differential K -theory. Lott [21] used Karoubi's construction to develop an index theorem, but it was mostly applications in string theory that have rekindled a considerable interest in differential K -theory (see Freed [11] for example). Perhaps one of the most remarkable steps forward in differential cohomologies is due to Hopkins and Singer [16] wherein they construct a differential extension of any exotic cohomology theory in a homotopy-theoretic way. Following this work, Bunke and Schick [7], Freed and Lott [13], Klonoff [20], Simons and Sullivan [24], and Tradler et al. [25, 26] all came up with more concrete and geometric models of differential K -theory. For a more detailed survey on recent developments of differential K -theory, we refer the reader to Bunke and Schick [8].

There have been some attempts to twist differential K -theory. Carey et al. [9] gave a construction that satisfies the square diagram and short exact sequences. Kahle and Valentino [17], in an attempt to precisely formulate the T -duality for Ramond–Ramond fields in the presence of a B -field, gave a list of axioms for twisted differential K -theory, which can be generalized as axioms for any twisted differential cohomology theory. They construct a canonical differential twist for the differential K -theory of the total space of any torus bundle [17, Section 2.2]. However, a construction of twisted differential K -theory that satisfies Kahle–Valentino axioms had not been found until very recently: in 2014, Bunke and Nikolaus [5] constructed a differential refinement of any twisted cohomology theory. Their construction of twisted differential K -theory satisfies several properties we would expect, including all Kahle–Valentino axioms, except the push-forward axiom which is not addressed in [5]. The construction of Bunke and Nikolaus provides a correct model for twisted differential cohomology theory in that their model combines twisted cohomology groups and twisted differential forms in a homotopy theoretic way, analogous to what Hopkins and Singer did in the untwisted case. However, the Bunke–Nikolaus model is not very geometric just as the Hopkins–Singer model is not. We might hope that there exists a more geometric model for a twisted differential cohomology theory, at least in the case of K -theory.

The goal of this paper is to construct such a geometric model of twisted differential even K -theory in the case that the underlying topological twists represent torsion classes in degree 3 integral cohomology. We use $U(1)$ -gerbes with connection as differential twists and twisted vector bundles with connection as cycles. We also have constructed a twisted differential K -theory for both torsion and non-torsion twistings using lifting bundle gerbes with connection and curving as differential twists and U_{tr} -bundle gerbe modules with connection (due to Bouwknegt et al. [3]) as cycles, which will be discussed in the second paper of this series. Both of our models satisfy all of the Kahle-Valentino axioms except the push-forward axiom which, together with a model of odd twisted differential K -theory, will be discussed in subsequent papers.

This paper is organized as follows. In Sect. 2, we review twisted vector bundles and set up some notation. Section 3 constructs the twisted Chern character form and the twisted Chern-Simons form. We also verify several properties which will be needed in later sections. Section 4 defines differential twists and constructs an even twisted differential K -group. We then show that our construction is functorial, natural with respect to change of differential twist, define maps into and out of the twisted differential K -groups, and verify that our model fits into a twisted analogue of the differential K -theory hexagon diagram à la Simons and Sullivan [24].

Having constructed geometric models of the even twisted differential K -theory, a natural question arises: “Is there a map between our geometric model and the Bunke–Nikolaus model?” Bunke et al. [6] answered this question for the case of untwisted differential K -theory. In this case, there is a way to obtain a sheaf of spectra on the site of smooth manifolds using the symmetric monoidal category of vector bundles with connection. In [6], they obtain a map between this sheaf of spectra into a Hopkins–Singer sheaf of spectra by the universal property of the pullback. The induced map between abelian groups is called a *cycle map*. Constructing a twisted analogue of the cycle map along this vein is work in progress, which we hope to complete in the near future.

2 Review of twisted vector bundles and twisted K -theory

In this section, we set up notations and briefly review λ -twisted vector bundles. A good reference on twisted vector bundles is Karoubi [19], which has a broader account.

Notation 2.1 Throughout this paper, all of our manifolds are connected compact smooth manifolds, and all our maps are smooth maps unless specified otherwise. In particular, X and Y always denote manifolds. We will use the notation $U_{i_1 \dots i_n}$ to denote an n -fold intersection $U_{i_1} \cap \dots \cap U_{i_n}$. If an open cover is locally finite and every n -fold intersection is contractible for all $n \in \mathbb{Z}^+$, we will call it a good cover.

Definition 2.2 Let $\mathcal{U} = \{U_i\}_{i \in I}$ be an open cover of X , and λ be a $U(1)$ -valued completely normalized Čech 2-cocycle as recalled below. A λ -**twisted vector bundle** E of rank n over X consists of a family of product bundles $\{U_i \times \mathbb{C}^n : U_i \in \mathcal{U}\}_{i \in \Lambda}$ together with transition maps

$$g_{ji} : U_{ij} \rightarrow U(n)$$

satisfying

$$g_{ii} = \mathbf{1}, \quad g_{ji} = g_{ij}^{-1}, \quad g_{kj}g_{ji} = g_{ki}\lambda_{kji}.$$

- Remark 2.3* (1) Recall that a Čech cocycle $\zeta = (\zeta_{i_1 \dots i_n})$ is called *completely normalized* if $\zeta_{i_1 \dots i_n} \equiv 1$ whenever there is a repeated index, and $\zeta_{\sigma(i_1) \dots \sigma(i_n)} = (\zeta_{i_1 \dots i_n})^{\text{sign}(\sigma)}$ for any $\sigma \in \mathfrak{S}_n$, where \mathfrak{S}_n is the symmetric group on n letters.
- (2) We write a λ -twisted vector bundle E of rank n as a triple $(\mathcal{U}, \{g_{ji}\}, \{\lambda_{kji}\})$, or a pair $(\{g_{ji}\}, \{\lambda_{kji}\})$ if the open cover \mathcal{U} is clear from the context. When the rank n is zero, there exists a λ -twisted vector bundle $\mathcal{O} = (\{g_{ji}^{\mathcal{O}}\}, \{\lambda_{kji}\})$ with $g_{ji}^{\mathcal{O}} = \mathbf{1}$ for all $i, j \in \Lambda$. We call it the *zero λ -twisted vector bundle*.

Definition 2.4 A **morphism** f from a λ -twisted vector bundle $E = (\{g_{ji}\}, \{\lambda_{kji}\})$ of rank n to a λ -twisted vector bundle $F = (\{h_{ji}\}, \{\lambda_{kji}\})$ of rank n , with respect to the same open cover $\{U_i\}_{i \in \Lambda}$ of the base X , is a family of maps $\{f_i : U_i \rightarrow U(n)\}_{i \in \Lambda}$ such that

$$f_j(x)g_{ji}(x) = h_{ji}(x)f_i(x) \quad \text{for all } x \in U_{ij} \quad \text{and all } i, j \in \Lambda.$$

Definition 2.5 Let $E = (\{g_{ji}\}, \{\lambda_{kji}\})$ and $F = (\{h_{ji}\}, \{\lambda_{kji}\})$ be λ -twisted vector bundles of rank n and m with respect to the same covering $\mathcal{U} = \{U_i\}$ of X . The **direct sum** $E \oplus F$ is defined by $(\{g_{ji} \oplus h_{ji}\}, \{\lambda_{kji}\})$, and is a λ -twisted vector bundle of rank $n + m$. The symbol \oplus between two transition maps denotes the block sum of matrices.

We denote the category of λ -twisted vector bundles over X defined on an open cover \mathcal{U} by $\mathbf{Bun}(\mathcal{U}, \lambda)$. The category $\mathbf{Bun}(\mathcal{U}, \lambda)$ is an additive category with respect to the direct sum \oplus .

Definition 2.6 The **twisted K -theory** of X defined on an open cover \mathcal{U} with a $U(1)$ -gerbe twisting λ , denoted by $K^0(\mathcal{U}, \lambda)$, is the Grothendieck group of the commutative monoid $\mathbf{Vect}(\mathcal{U}, \lambda)$ of isomorphism classes of λ -twisted vector bundles on \mathcal{U} .

Remark 2.7 The isomorphism class of the group $K^0(\mathcal{U}, \lambda)$ for a fixed \mathcal{U} depends only on the Čech cohomology class of λ . To see this, let \mathcal{C} and \mathcal{D} be additive categories. Recall that if an additive functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is an equivalence of additive categories, then it induces an isomorphism of groups $F_* : K(\mathcal{C}) \rightarrow K(\mathcal{D})$, where K denotes the K -theory functor from additive categories to abelian groups. Let σ and λ be cohomologous $U(1)$ -valued Čech 2-cocycles defined on \mathcal{U} , i.e., $\lambda_{kji} = \sigma_{kji}\chi_{ji}\chi_{ik}\chi_{kj}$ for some Čech 1-cochain χ . There is an additive functor $\Phi : \mathbf{Bun}(\mathcal{U}, \sigma) \rightarrow \mathbf{Bun}(\mathcal{U}, \lambda)$ that is an isomorphism of categories. The functor Φ takes a σ -twisted vector bundle $(\{g_{ji}\}, \{\sigma_{kji}\})$ to a λ -twisted vector bundle $(\{g_{ji}\chi_{ji}\}, \{\lambda_{kji}\})$ and takes a morphism between σ -twisted vector bundles to itself. The inverse of Φ is defined similarly by taking a λ -twisted vector bundle $(\{g_{ji}\}, \{\lambda_{kji}\})$ to the σ -twisted vector bundle $(\{g_{ji}\chi_{ji}^{-1}\}, \{\sigma_{kji}\})$. Therefore, the induced map of groups $\Phi_* : K^0(\mathcal{U}, \sigma) \rightarrow K^0(\mathcal{U}, \lambda)$ is an isomorphism.

Definition 2.8 Let $f : Y \rightarrow X$ be a map, and $E = (\{g_{ji}\}, \{\lambda_{kji}\})$ be a λ -twisted vector bundle defined on a covering $\mathcal{U} = \{U_i\}$ of X . Let $f^{-1}\mathcal{U}$ denote the open cover on Y consisting of open sets of the form $f^{-1}(U_i)$. The **pull-back** of the λ -twisted vector bundle E is a $(\lambda \circ f)$ -twisted vector bundle $(f^{-1}\mathcal{U}, \{g_{ji} \circ f\}, \{\lambda_{kji} \circ f\})$ on Y denoted by $f^*(E)$.

Proposition 2.9 *The map*

$$f^* : \text{Vect}(\mathcal{U}, \lambda) \rightarrow \text{Vect}(f^{-1}\mathcal{U}, \lambda \circ f)$$

$$[E] \mapsto [f^*E]$$

is a monoid homomorphism with respect to \oplus and therefore induces a group homomorphism

$$f^* : K^0(\mathcal{U}, \lambda) \rightarrow K^0(f^{-1}\mathcal{U}, \lambda \circ f)$$

$$[E] - [F] \mapsto [f^*E] - [f^*F].$$

Proof The map is well-defined on $\text{Vect}(\mathcal{U}, \lambda)$. Given another λ -twisted vector bundle F , we have $f^*(E) \oplus f^*(F) = f^*(E \oplus F)$. Hence f^* is a monoid homomorphism, which induces a group homomorphism f^* between K -groups. \square

3 Chern–Weil theory of twisted vector bundles

In this section, we review Chern–Weil theory of twisted vector bundles and define twisted Chern–Simons forms. We will also prove several lemmata which will be needed in subsequent sections. We begin with a summary of the language of $U(1)$ -gerbes with connection used in this paper. We refer the reader to Gawędzki and Reis [15] for more details.

Definition 3.1 Let X be a manifold and $\mathcal{U} := \{U_i\}_{i \in \Lambda}$ an open cover of X . A $U(1)$ -gerbe over X subordinate to \mathcal{U} is a $U(1)$ -valued completely normalized Čech 2-cocycle $\{\lambda_{kji}\} \in \check{Z}^2(\mathcal{U}, U(1))$. A **connection** on a $U(1)$ -gerbe $\{\lambda_{kji}\}$ on \mathcal{U} is a pair $(\{A_{ji}\}, \{B_i\})$ consisting of a family of differential 1-forms $\{A_{ji} \in \Omega^1(U_{ij}; \sqrt{-1}\mathbb{R})\}_{i,j \in \Lambda}$ and a family of differential 2-forms $\{B_i \in \Omega^2(U_i; \sqrt{-1}\mathbb{R})\}_{i \in \Lambda}$ satisfying the following relations:

- C1.** $\lambda_{kji}\lambda_{lji}^{-1}\lambda_{lki}\lambda_{lkj}^{-1} = 1$
- C2.** $d \log \lambda_{kji} = A_{ji} + A_{ik} + A_{kj}$
- C3.** $B_j - B_i = dA_{ji}$

- Remark 3.2* (1) A $U(1)$ -gerbe with connection on \mathcal{U} is therefore a Deligne cocycle of degree 2. Notice that our total differential is $D = d + (-1)^q \delta$ on $\check{C}^p(\mathcal{U}, \Omega^q)$.
- (2) From $dB_i = dB_j$ for all $i, j \in \Lambda$, the family of exact 3-forms $\{dB_i\}_{i \in \Lambda}$ defines a global closed differential 3-form H . The differential form H is called the *curvature* of the $U(1)$ -gerbe or the *Neveu–Schwarz 3-form*.
- (3) Let $\{\lambda_{kji}\} \in \check{Z}^2(\mathcal{U}, U(1))$ be a $U(1)$ -gerbe, and $\delta : \check{H}^2(\mathcal{U}, U(1)) \rightarrow H^3(X; 2\pi i\mathbb{Z})$ be the connecting map. The image in $H^3_{\text{dR}}(X; \sqrt{-1}\mathbb{R})$ of the cohomology class $\delta([\lambda]) \in H^3(X; 2\pi i\mathbb{Z})$ coincides with the cohomology class of $H \in H^3_{\text{dR}}(X; \sqrt{-1}\mathbb{R})$. (See Brylinski [4, p.175] Corollary 4.2.8.)

Throughout the rest of this paper $\widehat{\lambda} = (\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\})$ always denotes a $U(1)$ -gerbe with connection defined on an open cover $\mathcal{U} = \{U_i\}_{i \in \Lambda}$ of X and H denotes the 3-curvature form of $\widehat{\lambda}$. We assume that the Dixmier–Douady class of λ is a torsion class in $H^3(X; \mathbb{Z})$.

Definition 3.3 Let $\widehat{\lambda} = (\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\})$ be as above, $E = (\mathcal{U}, \{g_{ji}\}, \{\lambda_{kji}\})$ a smooth λ -twisted vector bundle of rank n . A **connection** on E compatible with $\widehat{\lambda}$ is a family $\Gamma = \{\Gamma_i \in \Omega^1(U_i; \mathfrak{u}(n))\}_{i \in \Lambda}$ satisfying

$$\Gamma_i - g_{ji}^{-1} \Gamma_j g_{ji} - g_{ji}^{-1} dg_{ji} = -A_{ji} \cdot \mathbf{1}, \tag{1}$$

where $A_{ji} \in \Omega^1(U_{ij}; i\mathbb{R})$. Here $\mathfrak{u}(n)$ denotes the Lie algebra of $U(n)$, and $\mathbf{1}$ the identity matrix.

Lemma 3.4 *In the notation of Definition 3.3, $A_{ji} \cdot \mathbf{1} - A_{ki} \cdot \mathbf{1} + A_{kj} \cdot \mathbf{1} = \lambda_{kji}^{-1} d\lambda_{kji} \cdot \mathbf{1}$.*

Proof

$$\begin{aligned} (A_{ji} - A_{ki} + A_{kj}) \cdot \mathbf{1} &= -(\Gamma_i - g_{ji}^{-1} \Gamma_j g_{ji} - g_{ji}^{-1} dg_{ji}) \\ &\quad + (\Gamma_i - g_{ki}^{-1} \Gamma_k g_{ki} - g_{ki}^{-1} dg_{ki}) + A_{kj} \cdot \mathbf{1} \\ &= g_{ji}^{-1} (g_{kj}^{-1} \Gamma_k g_{kj} + g_{kj}^{-1} dg_{kj} - A_{kj} \cdot \mathbf{1}) g_{ji} + g_{ji}^{-1} dg_{ji} \\ &\quad - \lambda_{kji} g_{ij} g_{jk} \Gamma_k g_{kj} g_{ji} \lambda_{kji}^{-1} - g_{ki}^{-1} dg_{ki} + A_{kj} \cdot \mathbf{1} \\ &= g_{ji}^{-1} g_{kj}^{-1} \Gamma_k g_{kj} g_{ji} + g_{ji}^{-1} g_{kj}^{-1} dg_{kj} g_{ji} - A_{kj} \cdot \mathbf{1} + g_{ji}^{-1} dg_{ji} \\ &\quad - g_{ij} g_{jk} \Gamma_k g_{kj} g_{ji} - g_{ki}^{-1} dg_{ki} + A_{kj} \cdot \mathbf{1} \\ &= g_{ji}^{-1} g_{kj}^{-1} (-g_{kj} dg_{ji} + dg_{ki} \lambda_{kji} + g_{ki} d\lambda_{kji}) + g_{ji}^{-1} dg_{ji} \\ &\quad - g_{ki}^{-1} dg_{ki} \\ &= g_{ik} g_{kj} g_{ji} g_{ji}^{-1} g_{kj}^{-1} dg_{ki} + g_{ji}^{-1} g_{kj}^{-1} g_{ki} d\lambda_{kji} - g_{ki}^{-1} dg_{ki} \\ &= \lambda_{kji}^{-1} d\lambda_{kji} \cdot \mathbf{1}. \end{aligned}$$

□

Remark 3.5 For any λ -twisted vector bundle E , there exists a connection on E associated with $\widehat{\lambda}$. See [19, p. 244].

Definition 3.6 Let $\widehat{\lambda} = (\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\})$ be as above, (E, Γ) a λ -twisted vector bundle $(\mathcal{U}, \{g_{ji}\}, \{\lambda_{kji}\})$ of rank n with a connection Γ compatible with $\widehat{\lambda}$. The **curvature form** of Γ is the family $R = \{R_i \in \Omega^2(U_i; \mathfrak{u}(n))\}_{i \in \Lambda}$, where $R_i := d\Gamma_i + \Gamma_i \wedge \Gamma_i$.

Lemma 3.7 *For each $m \in \mathbb{Z}^+$, the differential forms $\text{tr}[(R_i - B_i \cdot \mathbf{1})^m]$ over the open sets U_i glue together to define a global differential form on X . Here $B_i \in \Omega^2(U_i; i\mathbb{R})$ is given by $\widehat{\lambda}$.*

Proof From (1), it follows that $R_i = g_{ji}^{-1}R_jg_{ji} - dA_{ji} \cdot \mathbf{1}$. Then

$$\begin{aligned} \text{tr}[(R_i - B_i \cdot \mathbf{1})^m] &= \text{tr}[(g_{ji}^{-1}R_jg_{ji} - dA_{ji} \cdot \mathbf{1} - B_i \cdot \mathbf{1})^m] \\ &= \text{tr}[(g_{ji}^{-1}R_jg_{ji} - B_j \cdot \mathbf{1})^m] \\ &= \text{tr} \left[\sum_{r=0}^m \binom{m}{r} g_{ji}^{-1} R_j^{m-r} g_{ji} (-1)^r (B_j \cdot \mathbf{1})^r \right] \\ &\stackrel{*}{=} \text{tr} \left[\sum_{r=0}^m \binom{m}{r} R_j^{m-r} (-1)^r (B_j \cdot \mathbf{1})^r \right] = \text{tr}[(R_j - B_j \cdot \mathbf{1})^m], \end{aligned}$$

where $\binom{m}{r}$ is the binomial coefficient m choose r , and at $*$, we have used $\text{tr}(AB) = \text{tr}(BA)$ and the fact that $B_j \cdot \mathbf{1}$ commutes with other matrices. \square

Definition 3.8 Let $\widehat{\lambda} = (\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\})$ be as above, H the 3-curvature of $\widehat{\lambda}$, and (E, Γ) a λ -twisted vector bundle with connection compatible with $\widehat{\lambda}$. For $m \in \mathbb{Z}^+$, the m th **twisted Chern character form** is defined by

$$\text{ch}_{(m)}(\Gamma)(x) := \text{tr}(R_i(x) - B_i(x) \cdot \mathbf{1})^m \quad x \in U_i.$$

When $m = 0$, define $\text{ch}_{(0)}(\Gamma)$ to be the rank of E . The **total twisted Chern character form** is defined by

$$\text{ch}(\Gamma) := \text{rank}(E) + \sum_{m=1}^{\infty} \frac{1}{m!} \text{ch}_{(m)}(\Gamma),$$

which will be sometimes denoted by $\text{ch}(E, \Gamma)$.

Remark 3.9 Recall that the \mathbb{Z}_2 -graded sequence of differential forms $\dots \rightarrow \Omega^{\text{even}}(X) \xrightarrow{d+H} \Omega^{\text{odd}}(X) \xrightarrow{d+H} \dots$ is a complex if H is a closed 3-form on X . Then the *twisted de Rham cohomology* of X is the cohomology of this complex and denoted by $H_H^{\text{even/odd}}(X)$. If closed 3-forms H and H' are cohomologous, i.e. $H' = H + d\xi$, the multiplication by $\exp(\xi)$ induces an isomorphism $H_H^\bullet(X) \rightarrow H_{H'}^\bullet(X)$. We refer the reader to Atiyah and Segal [1] for more details on twisted cohomology.

The following fact is well-known. (See [3, p. 29] for example.)

Proposition 3.10 For each $m \in \mathbb{Z}^+$,

$$d\text{ch}_{(m)}(\Gamma) = m\text{ch}_{(m-1)}(\Gamma)H.$$

Hence the total twisted Chern character form $\text{ch}(\Gamma)$ is $(d + H)$ -closed.

Proposition 3.11 The m th twisted chern character form is additive for all $m \geq 0$, i.e.,

$$\text{ch}_{(m)}(\Gamma^E \oplus \Gamma^F) = \text{ch}_{(m)}(\Gamma^E) + \text{ch}_{(m)}(\Gamma^F).$$

Definition 3.12 Let $\widehat{\lambda} = (\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\})$ be as above, and (E, Γ) a λ -twisted vector bundle $(\mathcal{U}, \{g_{ji}\}, \{\lambda_{kji}\})$ of rank n with a connection Γ compatible with $\widehat{\lambda}$. Let $f : (Y, \mathcal{V}) \rightarrow (X, \mathcal{U})$ be any map provided that $\mathcal{V} = f^{-1}\mathcal{U}$. The **pullback** of (E, Γ) along f is $f^*(E)$ together with the family

$$f^*\Gamma := \{f^*\Gamma_i\}_{i \in \Lambda},$$

where $f^*\Gamma_i \in \Omega^1(f^{-1}(U_i); \mathfrak{u}(n))$ is defined by entrywise pullback.

Proposition 3.13 $f^*\Gamma$ is a connection on the $(\lambda \circ f)$ -twisted vector bundle $(\mathcal{V}, \{g_{ji} \circ f\}, \{\lambda_{kji} \circ f\})$ of rank n compatible with $f^*\widehat{\lambda} = (\{\lambda_{kji} \circ f\}, \{f^*A_{ji}\}, \{f^*B_i\})$ and $f^*ch(\Gamma) = ch(f^*\Gamma)$.

Proposition 3.14 Let $\widehat{\lambda} = (\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\})$ be as above and $\varphi : E \rightarrow F$ an isomorphism of λ -twisted vector bundles over X with respect to the same open cover \mathcal{U} . Let Γ^E be a connection on E associated with $\widehat{\lambda}$ and Γ^F a connection on F associated with $\widehat{\lambda}$. Then $ch(\Gamma^F) = ch(\varphi^*\Gamma^E)$.

Remark 3.15 We shall prove in Proposition 3.26 that the total twisted Chern character in twisted de Rham cohomology group is independent of the choice of connection.

Proposition 3.16 Let $\widehat{\lambda} = (\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\})$ and $\widehat{\lambda}' = (\{\lambda'_{kji}\}, \{A'_{ji}\}, \{B'_i\})$ be two $U(1)$ -gerbes with connection defined on an open cover $\mathcal{U} = \{U_i\}_{i \in \Lambda}$ over X . Suppose $\widehat{\lambda}$ and $\widehat{\lambda}'$ are cohomologous as Deligne 2-cocycles such that $\widehat{\lambda}' = \widehat{\lambda} + D\widehat{\alpha}$, where $\widehat{\alpha} = (\{\chi_{ji}\}, \{\Pi_i\}) \in \check{C}^1(\mathcal{U}, \Omega^1)$. (See Remark 3.2 for the definition of D .) Let $E = (\mathcal{U}, \{g_{ji}\}, \{\lambda_{kji}\})$ be a λ -twisted vector bundle of rank n and $\Gamma = \{\Gamma_i\}_{i \in \Lambda}$ a connection on E compatible with $\widehat{\lambda}$. Define a λ' -twisted vector bundle E' with connection Γ' compatible with $\widehat{\lambda}'$ by

$$\begin{aligned} E' &:= (\mathcal{U}, \chi_{ji}g_{ji}, \lambda'_{kji}) \\ \Gamma' &:= \{\Gamma'_i\}_{i \in \Lambda}, \text{ where } \Gamma'_i := \Gamma_i + \Pi_i \cdot \mathbf{1}. \end{aligned}$$

Then $ch(\Gamma) = ch(\Gamma')$.

Remark 3.17 Since $\widehat{\lambda}$ and $\widehat{\lambda}'$ are cohomologous, their 3-curvatures are the same.

Proof of Proposition 3.16 From

$$\begin{aligned} R'_i &= d\Gamma'_i + \Gamma'_i \wedge \Gamma'_i = d(\Gamma_i + \Pi_i \cdot \mathbf{1}) + (\Gamma_i + \Pi_i \cdot \mathbf{1}) \wedge (\Gamma_i + \Pi_i \cdot \mathbf{1}) \\ &= d\Gamma_i + d\Pi_i \cdot \mathbf{1} + \Gamma_i \wedge \Gamma_i + \Gamma_i \wedge \Pi_i \cdot \mathbf{1} + \Pi_i \cdot \mathbf{1} \wedge \Gamma_i \\ &\quad + \Pi_i \cdot \mathbf{1} \wedge \Pi_i \cdot \mathbf{1} \\ &= R_i + d\Pi_i \cdot \mathbf{1}, \end{aligned}$$

it follows that

$$ch_{(m)}(E, \Gamma) - ch_{(m)}(E', \Gamma') = \text{tr}(R_i - B_i \cdot \mathbf{1})^m - \text{tr}(R'_i - B'_i \cdot \mathbf{1})^m = 0$$

since $B'_i - B_i = d\Pi_i$. □

Notation 3.18 Given $\widehat{\lambda}$ and $\xi \in \Omega^2(X; i\mathbb{R})$, we denote by $\widehat{\lambda}_\xi$ the $U(1)$ -gerbe with connection $(\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i + \xi|_{U_i}\})$. Let E be a λ -twisted vector bundle and $\Gamma = \{\Gamma_i\}_{i \in \mathcal{A}}$ be a connection on E associated with $\widehat{\lambda}$. We denote the same family $\{\Gamma_i\}_{i \in \mathcal{A}}$ on E that is associated with $\widehat{\lambda}_\xi$ as a connection on E by Γ_ξ . We also denote $\xi|_{U_i}$ by ξ_i .

Proposition 3.19 Let $\widehat{\lambda} = (\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\})$ be as above, E a λ -twisted vector bundle, Γ a connection on E , and $\xi \in \Omega^2(X; i\mathbb{R})$. Then $ch(\Gamma_{-\xi}) = ch(\Gamma) \wedge \exp(\xi)$.

Proof

$$\begin{aligned} ch(\Gamma_{-\xi}) &= \sum_{m=0}^{\infty} \frac{1}{m!} ch_{(m)}(\Gamma_{-\xi}) = \sum_{m=0}^{\infty} \frac{1}{m!} \text{tr}[(R_i - B_i \cdot \mathbf{1} + \xi_i \cdot \mathbf{1})^m] \\ &= \sum_{m=0}^{\infty} \frac{1}{m!} \text{tr}\left(\sum_{r=0}^m \binom{m}{r} (R_i - B_i \cdot \mathbf{1})^{m-r} \xi_i^r \cdot \mathbf{1}\right) \\ &= \sum_{m=0}^{\infty} \frac{1}{m!} \sum_{r=0}^m \left(\binom{m}{r} \text{tr}(R_i - B_i \cdot \mathbf{1})^{m-r}\right) \wedge \xi_i^r \\ &= \sum_{m=0}^{\infty} \frac{1}{m!} \sum_{r=0}^m \left(\frac{m!}{(m-r)!r!} ch_{(m-r)}(\Gamma)\right) \wedge \xi_i^r = \sum_{m=0}^{\infty} \sum_{r=0}^m \frac{ch_{(m-r)}(\Gamma)}{(m-r)!} \wedge \frac{\xi_i^r}{r!} \\ &= \sum_{m=0}^{\infty} \sum_{r=0}^{\infty} \frac{ch_{(m)}(\Gamma)}{(m)!} \wedge \frac{\xi_i^r}{r!} \quad \text{since } \sum_{m=0}^{\infty} \sum_{r=0}^m a_{r,m-r} = \sum_{m=0}^{\infty} \sum_{r=0}^{\infty} a_{r,m} \\ &= ch(\Gamma) \wedge \exp(\xi). \end{aligned}$$

□

Now we discuss the Chern–Simons transgression form in the twisted case.

Lemma 3.20 Let $\widehat{\lambda} = (\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\})$ be as above and Γ_0 and Γ_1 be connections on a λ -twisted vector bundle $E = (\mathcal{U}, \{g_{ji}\}, \{\lambda_{kji}\})$ such that both are compatible with $\widehat{\lambda}$. Then for each $t \in \mathbb{R}$,

$$\Gamma_t := (1 - t)\Gamma_0 + t\Gamma_1$$

is a connection on E compatible with $\widehat{\lambda}$, i.e., the space of $\widehat{\lambda}$ -compatible connections on E is an affine space modeled over $\Omega^1(X; \text{End}(E))$.

Proof

$$\begin{aligned} \Gamma_{ti} - g_{ji}^{-1} \Gamma_{tj} g_{ji} - g_{ji}^{-1} dg_{ji} &= (1 - t)\Gamma_{0i} + t\Gamma_{1i} - g_{ji}^{-1}((1 - t)\Gamma_{0j} + t\Gamma_{1j})g_{ji} \\ &\quad - g_{ji}^{-1} dg_{ji} \\ &= (1 - t)\Gamma_{0i} - (1 - t)g_{ji}^{-1} \Gamma_{0j} g_{ji} - (1 - t)g_{ji}^{-1} dg_{ji} \\ &\quad + t\Gamma_{1i} - t g_{ji}^{-1} \Gamma_{1j} g_{ji} - t g_{ji}^{-1} dg_{ji} \\ &= -(1 - t)A_{ji} \cdot \mathbf{1} - tA_{ji} \cdot \mathbf{1} = -A_{ji} \cdot \mathbf{1}. \end{aligned}$$

If Γ and Γ' are two different $\widehat{\lambda}$ -compatible connections on E , we have $\Gamma_i - \Gamma'_i = g_{ji}^{-1}(\Gamma_j - \Gamma'_j)g_{ji}$, so the space of $\widehat{\lambda}$ -compatible connections on E is an affine space modelled over $\Omega^1(X; \text{End}(E))$. Notice that $\text{End}(E)$ is an ordinary vector bundle. \square

Corollary 3.21 *Let Γ_0 and Γ_1 be as in Lemma 3.20. Let α_t and γ_t with $t \in I$ be two paths of connections each starting at Γ_0 and ending at Γ_1 and both α_t and γ_t are compatible with $\widehat{\lambda}$ for all $t \in I$. Then there exists a bigon (a polygon with two sides) of connections with edges α_t and γ_t such that every point on the bigon is a $\widehat{\lambda}$ -compatible connection on E .*

Proof By Lemma 3.20, for each fixed $t \in I$ and $s \in I$, the connection $(1 - s)\alpha_t + s\gamma_t$ is $\widehat{\lambda}$ -compatible. \square

Notation 3.22 We shall denote the projection map $X \times I \rightarrow X$ onto the first factor by p .

Lemma 3.23 *Let $\widehat{\lambda} = (\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\})$ be as above, E a λ -twisted vector bundle $(\mathcal{U}, \{g_{ji}\}, \{\lambda_{kji}\})$, and Γ_t a connection on E compatible with $\widehat{\lambda}$ for each $t \in I$. Then the family $\{\widetilde{\Gamma}_i\}_{i \in \Lambda}$ defined by $\widetilde{\Gamma}_i(x, t) := (p^*\Gamma_t)(x, t)$ is a connection on the $(\lambda \circ p)$ -twisted vector bundle $p^*E = (\mathcal{U} \times I, \{g_{ji} \circ p\}, \{\lambda_{kji} \circ p\})$ compatible with the pull-back $U(1)$ -gerbe with connection $p^*\widehat{\lambda} = (\{\lambda_{kji} \circ p\}, \{p^*A_{ji}\}, \{p^*B_i\})$.*

We refer the reader to Bott and Tu [2] for an account of the integration along the fiber.

Definition 3.24 Let $\widehat{\lambda} = (\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\})$ be as above, E a λ -twisted vector bundle over X , and $\gamma : t \mapsto \Gamma_t$ a path of connections on E such that each Γ_t is compatible with $\widehat{\lambda}$. The **twisted Chern–Simons form** of γ is the integration along the fiber:

$$cs(\gamma) := \int_I ch(\widetilde{\Gamma}) \in \Omega^{\text{odd}}(X; \mathbb{C}),$$

where $\widetilde{\Gamma}$ is the connection on p^*E defined by $\widetilde{\Gamma}(x, t) = (p^*\Gamma_t)(x, t)$.

The following lemma is certainly well-known, but we did not find a reference.

Lemma 3.25 *Let E be a smooth fiber bundle over X with fiber F a compact oriented smooth k -manifold with corners. Let $\int_F : \Omega^\bullet(E; \mathbb{C}) \rightarrow \Omega^{\bullet-k}(X; \mathbb{C})$ be the integration along the fiber map and $\omega \in \Omega^n(E; \mathbb{C})$ for $n \geq k$. Then*

$$d \int_F \omega = \int_F d\omega + (-1)^{n-k} \int_{\partial F} \omega. \tag{2}$$

Proposition 3.26 *Let $\widehat{\lambda} = (\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\})$ be as above, $E = (\mathcal{U}, \{g_{ji}\}, \{\lambda_{kji}\})$ a λ -twisted vector bundle of rank n , and $\gamma : t \mapsto \Gamma_t$ a path of connections on E joining Γ_0 and Γ_1 such that each Γ_t is compatible with $\widehat{\lambda}$. Then*

$$ch(\Gamma_0) - ch(\Gamma_1) = (d + H)cs(\gamma).$$

Proof Let $\tilde{\Gamma}$ be a connection on p^*E defined by $\tilde{\Gamma}(x, t) = (p^*\Gamma_t)(x, t)$. By Lemma 3.25, $d \int_I \text{ch}(\tilde{\Gamma}) = \int_I d\text{ch}(\tilde{\Gamma}) - \int_{\partial I} \text{ch}(\tilde{\Gamma}) = - \int_I \text{ch}(\tilde{\Gamma}) \wedge p^*H - \int_{\partial I} \text{ch}(\tilde{\Gamma}) = -H \wedge (\int_I \text{ch}(\tilde{\Gamma})) + \text{ch}(\Gamma_0) - \text{ch}(\Gamma_1)$. Hence the result. \square

Definition 3.27 The **twisted total Chern character** of E , denoted by $\text{ch}(E)$, is the twisted cohomology class of $\text{ch}(\Gamma)$ for any connection Γ on E .

Proposition 3.28 *The assignment*

$$\begin{aligned} \text{ch} : K^0(\mathcal{U}, \lambda) &\rightarrow H_H^{\text{even}}(X; \mathbb{C}) \\ [E] - [F] &\mapsto [\text{ch}(\Gamma^E)] - [\text{ch}(\Gamma^F)], \end{aligned}$$

with $(\{A_{ji}\}, \{B_i\})$ a connection on λ and Γ^E and Γ^F connections on λ -twisted vector bundles E and F , respectively, both compatible with $\hat{\lambda}$, is a well-defined group homomorphism called the **twisted Chern character**.

Before proving Proposition 3.28, we recall the following lemma and its generalizations, which are certainly well-known. We include a proof here for sake of completeness. (See also Bunke and Nikolaus [5], Section 7).

Lemma 3.29 *Suppose $U(1)$ -gerbes λ and λ' defined on a good open cover \mathcal{U} of X are isomorphic: $\lambda'_{kji} = \lambda_{kji} + (\delta\chi)_{kji}$. Let $(\{A_{ji}\}, \{B_i\})$ on λ and $(\{A'_{ji}\}, \{B'_i\})$ on λ' be arbitrarily chosen connections. Then there exists a Deligne 1-cochain $\hat{\alpha} = (\{\chi_{ji}\}, \{\Pi_i\})$ and $\hat{\xi} \in \Omega^2(X; i\mathbb{R})$ such that $\hat{\lambda}' = \hat{\lambda}_{\hat{\xi}} + D\hat{\alpha}$, where $\hat{\lambda} = (\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\})$ and $\hat{\lambda}' = (\{\lambda'_{kji}\}, \{A'_{ji}\}, \{B'_i\})$.*

We need the following lemma, which is well-known, see e.g. Bott and Tu [2, p. 94], Proposition 8.5.

Lemma 3.30 $\check{H}^p(\mathcal{U}, \Omega^q) = 0$, for all $p \geq 1$.

Proof of Lemma 3.29 We denote the 3-curvature of $\hat{\lambda}$ and $\hat{\lambda}'$ by H and H' , respectively. The curvature 3-forms of $\hat{\lambda}$ and $\hat{\lambda}'$ are cohomologous, i.e. $H' - H = d\zeta$ for some $\zeta \in \Omega^2(X; \sqrt{-1}\mathbb{R})$. Over each open set U_i , we have $d(B'_i - B_i) = d\zeta_i$, where $\zeta_i := \zeta|_{U_i}$, and by Poincaré’s Lemma, there exists a 1-form ω_i on each U_i such that $B'_i = B_i + \zeta_i + d\omega_i$. Now by the cocycle condition, $dA'_{ji} = B'_j - B'_i = B_j - B_i + d(\omega_j - \omega_i) = dA_{ji} + d(\omega_j - \omega_i)$ and so there exists a $U(1)$ -valued function μ_{ji} over each U_{ij} such that $A'_{ji} = A_{ji} + \omega_j - \omega_i + d \log \mu_{ji}$. Take the Čech differential of both sides and get $\delta(d \log(\chi_{ji} \mu_{ji}^{-1}))_{kji} = 0$. By Lemma 3.30, $d \log(\chi_{ji} \mu_{ji}^{-1}) = \gamma_j - \gamma_i$ for some $\gamma \in \check{C}^1(\mathcal{U}, \Omega^1)$, hence $d \log(\chi_{ji}) = d \log(\mu_{ji}) + (\gamma_j - \gamma_i)$. Notice that the family $\{d\gamma_i\}_{i \in \Delta}$ defines a global 2-form on X . Thus setting $\hat{\alpha} = (\{\chi_{ji}\}, \{\omega_i - \gamma_i\})$ and $\hat{\xi}|_{U_i} = \zeta_i + d\gamma_i$ proves the claim. \square

Remark 3.31 When the underlying gerbes λ and λ' are identical, a special case of Lemma 3.29 indicates that, under different choices of connection on λ , the corresponding twisted Chern characters are related by exp of a global 2-form.

Proof of Proposition 3.28 Well-definedness: Let $\widehat{\lambda}$ be fixed. The image of ch is independent of the choice of connections on the twisted vector bundles E and F by Proposition 3.26.

Suppose $\widehat{\lambda}$ and $\widehat{\lambda}'$ are the same $U(1)$ -gerbes λ endowed with different connections and $\widehat{\lambda}' = \widehat{\lambda} + D\widehat{\alpha}$. Then by Proposition 3.16, the image of ch is invariant under cohomologous change of $U(1)$ -gerbe connection. (Notice that the image of ch is not invariant under the arbitrary change of $U(1)$ -gerbe connection. See Remark 3.31.)

Suppose there exists a λ -twisted vector bundle G with an isomorphism $\varphi : E \oplus G \rightarrow \overline{E} \oplus G$. Let Γ^G be an arbitrary connection on G . By Lemma 3.11 and Proposition 3.14,

$$\begin{aligned} [\text{ch}(\Gamma^E)] + [\text{ch}(\Gamma^G)] &= [\text{ch}(\Gamma^E \oplus \Gamma^G)] = [\text{ch}(\varphi^*(\Gamma^{\overline{E}} \oplus \Gamma^G))] = [\text{ch}(\Gamma^{\overline{E}} \oplus \Gamma^G)] \\ &= [\text{ch}(\Gamma^{\overline{E}})] + [\text{ch}(\Gamma^G)]. \end{aligned}$$

From this, well-definedness of ch on $K^0(\mathcal{U}, \lambda)$ follows.

Group homomorphism: This follows from Lemma 3.11:

$$\begin{aligned} \text{ch}([E] - [F] + [\overline{E}] - [\overline{F}]) &= \text{ch}([E \oplus \overline{E}] - [F \oplus \overline{F}]) = \text{ch}(\Gamma^E \oplus \Gamma^{\overline{E}}) \\ &\quad - \text{ch}(\Gamma^F \oplus \Gamma^{\overline{F}}) \\ &= \text{ch}(\Gamma^E) - \text{ch}(\Gamma^F) + \text{ch}(\Gamma^{\overline{E}}) - \text{ch}(\Gamma^{\overline{F}}) \\ &= \text{ch}([E] - [F]) + \text{ch}([\overline{E}] - [\overline{F}]). \end{aligned}$$

□

Proposition 3.32 *Let $\widehat{\lambda} = (\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\})$ be as above, E a λ -twisted vector bundle $(\mathcal{U}, \{g_{ji}\}, \{\lambda_{kji}\})$, and Γ_0 and Γ_1 two connections on E joined by two different paths of connections α_t and γ_t on E , such that each of α_t and γ_t is compatible with $\widehat{\lambda}$ for all $t \in I$. Then*

$$cs(\gamma) - cs(\alpha) \in \text{Im}(d + H).$$

Proof The paths α and γ define connections on p^*E over $X \times I$, which we denote by $\widetilde{\alpha}$ and $\widetilde{\gamma}$, respectively. Then there exists a path of connections on p^*E interpolating between $\widetilde{\alpha}$ and $\widetilde{\gamma}$ (by Corollary 3.21). Accordingly this path of connection defines a connection $\widetilde{\beta}$ on q^*p^*E over $X \times I \times I$, where $q : X \times I \times I \rightarrow X \times I$ is the projection map forgetting the third factor. By applying Lemma 3.25 to the twisted Chern character form $\text{ch}(q^*p^*E, \widetilde{\beta})$, we get

$$\begin{aligned} d \int_{I \times I} \text{ch}(q^*p^*E, \widetilde{\beta}) &= \int_{I \times I} d\text{ch}(q^*p^*E, \widetilde{\beta}) + \int_{\partial(I \times I)} \text{ch}(q^*p^*E, \widetilde{\beta}) \\ &= - \left(\int_{I \times I} \text{ch}(q^*p^*E, \widetilde{\beta}) \right) \wedge H + \int_I \text{ch}(p^*E, \widetilde{\gamma}) \\ &\quad - \int_I \text{ch}(p^*E, \widetilde{\alpha}) \end{aligned}$$

Hence,

$$\text{cs}(\gamma) - \text{cs}(\alpha) = (d + H) \int_{I \times I} \text{ch}(q^* p^* E, \tilde{\beta})$$

□

Proposition 3.33 *Let $\varphi : E \rightarrow F$ be an isomorphism of λ -twisted vector bundles over X . Let $\gamma : t \mapsto \Gamma^t$ be a path of connections on F . Then $\text{cs}(\varphi^* \gamma) = \text{cs}(\gamma)$.*

4 Twisted differential K -theory

This section constitutes the main part of this paper. We define differential twists and construct a twisted differential K -group (Sects. 4.1, 4.2) using triples consisting of a twisted vector bundle, a connection, and an odd differential form modulo exact forms in a twisted de Rham complex. We verify that our construction is functorial (Sect. 4.3) and natural with respect to change of twists (Sect. 4.4). In Sects. 4.5 and 4.7, we define the I , R , and a maps and verify the exact sequence involving the a and I maps. Finally, we show commutativity of diagrams and exactness of sequences consisting the hexagon diagram à la Simons and Sullivan [24] (Sect. 4.8), and verify that the maps I , R , and a are compatible with change of twists (Sect. 4.9).

4.1 Differential twists

Definition 4.1 The **torsion differential K -twists** for an open cover \mathcal{U} of X , denoted by $\text{Twist}_K^{\text{tor}}(\mathcal{U})$, is the groupoid whose objects are $U(1)$ -gerbes with connection $\hat{\lambda} = (\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\})$ each of which has an underlying $U(1)$ -gerbe representing a torsion class in $H^3(X; \mathbb{Z})$. For any two objects $\hat{\lambda}_1$ and $\hat{\lambda}_2$ in this groupoid, the Hom set is defined by $\text{Hom}(\hat{\lambda}_1, \hat{\lambda}_2) = \{(\hat{\alpha}, \xi) \in \check{C}^1(\mathcal{U}; \Omega^0) \oplus \check{C}^0(\mathcal{U}; \Omega^1) \oplus \Omega^2(X; i\mathbb{R})\} : \hat{\lambda}_2 = \hat{\lambda}_1 + D\hat{\alpha} + \xi$.

Proposition 4.2 (Existence) *Given any manifold X with an open cover \mathcal{U} , the torsion differential twist $\text{Twist}_K^{\text{tor}}(\mathcal{U})$ consists of at least one object.*

Proof The statement amounts to the existence of a connection on a local $U(1)$ -bundle gerbe, which follows from the existence of partitions of unity as shown in Murray [22]. □

Notation 4.3 (1) The **torsion topological K -twists** for an open cover \mathcal{U} of a manifold X is the groupoid, denoted by $\text{Twist}_K^{\text{top}}(\mathcal{U})$, whose objects are $U(1)$ -gerbes $\lambda = \{\lambda_{kji}\}$ each representing a torsion class in $H^3(X; \mathbb{Z})$. A morphism from λ_1 to λ_2 is a Čech 1-cochain $\alpha = (\chi_{ji}) \in \check{C}^1(\mathcal{U}, U(1))$ such that $\lambda_1 = \lambda_2 + \delta\alpha$.
 (2) Define the groupoid $\Omega_{\text{cl}}^3(X; i\mathbb{R})$ of $i\mathbb{R}$ -valued closed differential 3-forms on X as follows. Objects are $i\mathbb{R}$ -valued closed differential 3-forms on X . A morphism from ω to ω' is a differential 2-form α on X modulo exact forms satisfying $\omega = \omega' + d\alpha$.

Definition 4.4 The **forgetful** and **curvature** functors are given by the assignments

$$\begin{aligned} \mathcal{F} : \text{Twist}_{\widehat{K}}^{\text{tor}}(\mathcal{U}) &\rightarrow \text{Twist}_K^{\text{tor}}(\mathcal{U}) & \text{Curv} : \text{Twist}_{\widehat{K}}^{\text{tor}}(\mathcal{U}) &\rightarrow \Omega_{\text{cl}}^3(X; i\mathbb{R}) \\ \widehat{\lambda} = (\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\}) &\mapsto \{\lambda_{kji}\} & \widehat{\lambda} = (\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\}) &\mapsto \text{Curv}(\widehat{\lambda}) = H, \\ (\widehat{\alpha}, \xi) &\mapsto \alpha & (\widehat{\alpha}, \xi) &\mapsto d\xi \end{aligned}$$

where $\widehat{\alpha} = (\{\chi_{ji}\}, \{\Pi_i\})$, $\alpha = (\{\chi_{ji}\})$, and $H|_{U_i} = dB_i$ for all $i \in \Lambda$.

Remark 4.5 Let $f : (Y, \mathcal{V}) \rightarrow (X, \mathcal{U})$ be a map with $\mathcal{V} = f^{-1}\mathcal{U}$. The following diagrams commute:

$$\begin{array}{ccc} \text{Twist}_{\widehat{K}}^{\text{tor}}(\mathcal{V}) & \xrightarrow{\mathcal{F}} & \text{Twist}_K^{\text{tor}}(\mathcal{V}) & \text{Twist}_{\widehat{K}}^{\text{tor}}(\mathcal{V}) & \xrightarrow{\text{Curv}} & \Omega_{\text{cl}}^3(Y; i\mathbb{R}) \\ f^* \uparrow & & f^* \uparrow & f^* \uparrow & & f^* \uparrow \\ \text{Twist}_{\widehat{K}}^{\text{tor}}(\mathcal{U}) & \xrightarrow{\mathcal{F}} & \text{Twist}_K^{\text{tor}}(\mathcal{U}) & \text{Twist}_{\widehat{K}}^{\text{tor}}(\mathcal{U}) & \xrightarrow{\text{Curv}} & \Omega_{\text{cl}}^3(X; i\mathbb{R}) \end{array}$$

Here f^* on torsion differential twists takes each torsion $U(1)$ -gerbe with connection to its pullback $U(1)$ -gerbe with pullback connection, and f^* on topological twists does the same on torsion $U(1)$ -gerbes.

Notation 4.6 Throughout this section, we shall use the notation $\widehat{\lambda}$ to denote a differential twist $(\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\}) \in \text{Twist}_{\widehat{K}}^{\text{tor}}(\mathcal{U})$, H for $\text{Curv}(\widehat{\lambda})$, and λ for $\mathcal{F}(\widehat{\lambda})$.

4.2 Twisted differential K -group

Definition 4.7 A $\widehat{K}^0(\mathcal{U}; \widehat{\lambda})$ -**generator** is a triple (E, Γ, ω) consisting of a λ -twisted vector bundle E defined on the open cover $\mathcal{U} = \{U_i\}_{i \in \Lambda}$ on X , a connection Γ on E compatible with $\widehat{\lambda}$, and $\omega \in \Omega^{\text{odd}}(X; \mathbb{C})/\text{Im}(d + H)$.

Definition 4.8 Let E be any λ -twisted vector bundle with a path of connections γ joining Γ_0 and Γ_1 and each connection on the path being compatible with $\widehat{\lambda}$. Define $\text{CS}(\Gamma_0, \Gamma_1) := \text{cs}(\gamma) \pmod{\text{Im}(d + H)}$.

Remark 4.9 By Proposition 3.32, Definition 4.8 is independent of the choice of path of connections. Furthermore, we have $\text{CS}(\Gamma_0, \Gamma_1) + \text{CS}(\Gamma_1, \Gamma_2) = \text{CS}(\Gamma_0, \Gamma_2)$.

Definition 4.10 Two $\widehat{K}^0(\mathcal{U}; \widehat{\lambda})$ -generators (E, Γ, ω) and (E', Γ', ω') are **equivalent** if there exists a λ -twisted vector bundle with connection (F, Γ^F) and a λ -twisted vector bundle isomorphism $\varphi = \{\varphi_i\}_{i \in \Lambda} : E \oplus F \rightarrow E' \oplus F$ such that $\text{CS}(\Gamma \oplus \Gamma^F, \varphi^*(\Gamma' \oplus \Gamma^F)) = \omega - \omega'$.

Lemma 4.11 *The relation between triples in Definition 4.10 is an equivalence relation.*

Proof The relation is reflexive since cs of a loop is $(d + H)$ -exact. For symmetry, suppose (E, Γ, ω) and (E', Γ', ω') are equivalent, i.e., there exists a λ -twisted vector bundle with connection (F, Γ^F) whose connection is compatible with $\widehat{\lambda}$ such that there is an isomorphism of λ -twisted vector bundles $\varphi : E \oplus F \xrightarrow{\cong} E' \oplus F$, and $CS(\Gamma \oplus \Gamma^F, \varphi^*(\Gamma' \oplus \Gamma^F)) = \omega - \omega'$. By Proposition 3.33,

$$CS(\Gamma \oplus \Gamma^F, \varphi^*(\Gamma' \oplus \Gamma^F)) = CS((\varphi^{-1})^*(\Gamma \oplus \Gamma^F), \Gamma' \oplus \Gamma^F).$$

This proves the symmetry. For transitivity, suppose (E, Γ, ω) is equivalent to (E', Γ', ω') and (E', Γ', ω') is equivalent to $(E'', \Gamma'', \omega'')$, i.e., there exists a λ -twisted vector bundle with connection (F, Γ^F) whose connection is compatible with $\widehat{\lambda}$ such that there is an isomorphism of λ -twisted vector bundles $\varphi : E \oplus F \xrightarrow{\cong} E' \oplus F$, and $CS(\Gamma \oplus \Gamma^F, \varphi^*(\Gamma' \oplus \Gamma^F)) = \omega - \omega'$, and there exists a λ -twisted vector bundle with connection $(F', \Gamma^{F'})$ whose connection is compatible with $\widehat{\lambda}$ such that there is an isomorphism of λ -twisted vector bundles $\varphi' : E' \oplus F' \xrightarrow{\cong} E'' \oplus F'$, and $CS(\Gamma' \oplus \Gamma^{F'}, \varphi'^*(\Gamma'' \oplus \Gamma^{F'})) = \omega' - \omega''$. Then by taking the λ -twisted vector bundle with connection $(F \oplus F', \Gamma^F \oplus \Gamma^{F'})$, the isomorphism of λ -twisted vector bundles $\psi : E \oplus F \oplus F' \rightarrow E'' \oplus F \oplus F'$ is defined by the composition

$$E \oplus F \oplus F' \xrightarrow{\varphi \oplus \mathbf{1}} E' \oplus F \oplus F' \xrightarrow{\mathbf{1} \oplus \sigma} E' \oplus F' \oplus F \xrightarrow{\varphi' \oplus \mathbf{1}} E'' \oplus F' \oplus F \xrightarrow{\mathbf{1} \oplus \sigma^{-1}} E'' \oplus F \oplus F',$$

each of which is an isomorphism, and σ is the canonical λ -twisted vector bundle isomorphism $F \oplus F' \rightarrow F' \oplus F$. Furthermore,

$$\begin{aligned} &CS(\Gamma \oplus \Gamma^F \oplus \Gamma^{F'}, \psi^*(\Gamma'' \oplus \Gamma^F \oplus \Gamma^{F'})) \\ &= CS(\Gamma \oplus \Gamma^F \oplus \Gamma^{F'}, (\varphi \oplus \mathbf{1})^*(\Gamma' \oplus \Gamma^F \oplus \Gamma^{F'})) \\ &\quad + CS((\varphi \oplus \mathbf{1})^*(\Gamma' \oplus \Gamma^F \oplus \Gamma^{F'}), \psi^*(\Gamma'' \oplus \Gamma^F \oplus \Gamma^{F'})) \text{ by Remark 4.9} \\ &\stackrel{*}{=} \omega - \omega' + CS(\Gamma' \oplus \Gamma^F \oplus \Gamma^{F'}, ((\mathbf{1} \oplus \sigma^{-1}) \circ (\varphi' \oplus \mathbf{1}) \circ (\mathbf{1} \oplus \sigma))^*(\Gamma'' \oplus \Gamma^F \oplus \Gamma^{F'})) \\ &\stackrel{**}{=} \omega - \omega' + CS(\Gamma' \oplus \Gamma^{F'} \oplus \Gamma^F, ((\mathbf{1} \oplus \sigma^{-1}) \circ (\varphi' \oplus \mathbf{1}))^*(\Gamma'' \oplus \Gamma^F \oplus \Gamma^{F'})) \\ &= \omega - \omega' + CS(\Gamma' \oplus \Gamma^{F'} \oplus \Gamma^F, (\varphi' \oplus \mathbf{1})^*(\Gamma'' \oplus \Gamma^{F'} \oplus \Gamma^F)) \\ &= \omega - \omega' + \omega' - \omega'' = \omega - \omega'', \end{aligned}$$

At $*$ and $**$, 3.33 for the twisted bundle isomorphism $(\varphi \oplus \mathbf{1})^{-1}$ and $(\mathbf{1} \oplus \sigma^{-1})^{-1}$, respectively. Hence (E, Γ, ω) is equivalent to $(E'', \Gamma'', \omega'')$. \square

Lemma 4.12 *Let $[(E, \Gamma^E, \omega)]$ and $[(F, \Gamma^F, \eta)]$ be equivalence classes of $K^0(\mathcal{U}; \widehat{\lambda})$ -generators. The equivalence class of the $K^0(\mathcal{U}; \widehat{\lambda})$ -generator $(E \oplus F, \Gamma^E \oplus \Gamma^F, \omega + \eta)$ is independent of the choice of representatives of $[(E, \Gamma^E, \omega)]$ and $[(F, \Gamma^F, \eta)]$.*

Definition 4.13 The **addition** $+$ between two equivalence classes of $\widehat{K}^0(\mathcal{U}; \widehat{\lambda})$ -generators is defined by $[(E, \Gamma^E, \omega)] + [(F, \Gamma^F, \eta)] := [(E \oplus F, \Gamma^E \oplus \Gamma^F, \omega + \eta)]$.

Hence the set of all equivalence classes of $\widehat{K}^0(\mathcal{U}; \widehat{\lambda})$ -generators forms a commutative monoid $(\mathfrak{G}, +)$.

Definition 4.14 Let $\widehat{\lambda} \in \text{Twist}_{\widehat{K}}^{\text{tor}}(\mathcal{U})$. The **twisted differential K -group** is

$$\widehat{K}^0(\mathcal{U}, \widehat{\lambda}) := K(\mathfrak{G}),$$

where K denotes the group completion functor from commutative monoid to abelian groups.

4.3 Functoriality

Lemma 4.15 Let $\widehat{\lambda}$ be a differential twist, $E = (\mathcal{U}, \{g_{ji}\}, \{\lambda_{kji}\})$ a λ -twisted vector bundle of rank n , and $\Gamma = \{\Gamma_i\}_{i \in \Lambda}$ connection on E compatible with $\widehat{\lambda}$. Also let $f : (Y, \mathcal{V}) \rightarrow (X, \mathcal{U})$ be a map with $\mathcal{V} = f^{-1}\mathcal{U}$. If two triples (E, Γ, ω) and (E', Γ', ω') are equivalent, then $(f^*E, f^*\Gamma, f^*\omega)$ and $(f^*E', f^*\Gamma', f^*\omega')$ are equivalent.

Proposition 4.16 Given a map $f : (Y, \mathcal{V}) \rightarrow (X, \mathcal{U})$ with $\mathcal{V} = f^{-1}\mathcal{U}$, the assignment

$$f^* : \widehat{K}^0(\mathcal{U}, \widehat{\lambda}) \rightarrow \widehat{K}^0(\mathcal{V}, f^*\widehat{\lambda})$$

$$[(E, \Gamma^E, \omega)] - [(F, \Gamma^F, \eta)] \mapsto [(f^*E, f^*\Gamma^E, f^*\omega)] - [(f^*F, f^*\Gamma^F, f^*\eta)]$$

is a well-defined group homomorphism.

Let **Man** be the category whose objects are connected compact smooth manifolds equipped with an open cover. A morphism from (Y, \mathcal{V}) to (X, \mathcal{U}) is a smooth map satisfying $\mathcal{V} = f^{-1}(\mathcal{U})$. Also let **Ab** be the category of abelian groups.

Corollary 4.17 $\widehat{K}^0(-, \widehat{\lambda}) : \mathbf{Man}^{op} \rightarrow \mathbf{Ab}$ is a functor.

4.4 Naturality of twists

Proposition 4.18 Let $\widehat{\lambda} = (\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\})$ and $\widehat{\lambda}' = (\{\lambda'_{kji}\}, \{A'_{ji}\}, \{B'_i\})$ be any two objects of $\text{Twist}_{\widehat{K}}^{\text{tor}}(\mathcal{U})$ satisfying that $\widehat{\lambda}' = \widehat{\lambda} + D\widehat{\alpha}$ for some $\widehat{\alpha} = (\{\chi_{ji}\}, \{\Pi_i\})$. Let $E = (\mathcal{U}, \{g_{ji}\}, \{\lambda_{kji}\})$ be a λ -twisted vector bundle of rank n and $\Gamma = \{\Gamma_i\}_{i \in \Lambda}$ a connection on E compatible with $\widehat{\lambda}$. Define:

$$E' := (\mathcal{U}, \chi_{ji}g_{ji}, \lambda'_{kji})$$

$$\Gamma' := \{\Gamma'_i\}_{i \in \Lambda} \text{ where } \Gamma'_i := \Gamma_i + \Pi_i \cdot \mathbf{1} \tag{3}$$

$$\omega' := \omega$$

(1) The assignment

$$\phi_{\widehat{\alpha}} : \widehat{K}^0(\mathcal{U}; \widehat{\lambda}) \xrightarrow{\cong} \widehat{K}^0(\mathcal{U}; \widehat{\lambda}')$$

$$[(E, \Gamma, \omega)] - [(F, \nabla, \eta)] \mapsto [(E', \Gamma', \omega')] - [(F', \nabla', \eta')]$$

is an induced group isomorphism that is natural in \mathcal{U} .

(2) Let $\xi \in \Omega^2(X; i\mathbb{R})$. The assignment

$$\begin{aligned} \mathcal{E} : \widehat{K}^0(\mathcal{U}; \widehat{\lambda}) &\xrightarrow{\cong} \widehat{K}^0(\mathcal{U}; \widehat{\lambda}_\xi) \\ [(E, \Gamma, \omega)] - [(F, \nabla, \eta)] &\mapsto [(E, \Gamma_\xi, \omega \wedge \exp(-\xi))] - [(F, \nabla_\xi, \eta \wedge \exp(-\xi))] \end{aligned}$$

is a group isomorphism that is natural in \mathcal{U} .

Remark 4.19 The family Γ' above is a connection on the λ' -twisted vector bundle E' compatible with $\widehat{\lambda}'$:

$$\begin{aligned} &g_{ji}^{-1} \chi_{ji}^{-1} \Gamma'_j \chi_{ji} g_{ji} + g_{ji}^{-1} \chi_{ji}^{-1} d(\chi_{ji} g_{ji}) - A'_{ji} \cdot \mathbf{1} \\ &= g_{ji}^{-1} \Gamma_j g_{ji} + \Pi_j \cdot \mathbf{1} + \chi_{ji}^{-1} d\chi_{ji} \cdot \mathbf{1} + g_{ji}^{-1} dg_{ji} - (A_{ji} + \Pi_j - \Pi_i + d \log \chi_{ji}) \cdot \mathbf{1} \\ &= \Gamma_i + \Pi_i \cdot \mathbf{1} = \Gamma'_i. \end{aligned}$$

Proof of Proposition 4.18 (1) Suppose $(E, \Gamma, \omega) \sim (\overline{E}, \overline{\Gamma}, \overline{\omega})$, i.e., there exists a twisted vector bundle F and a connection Γ^F compatible with $\widehat{\lambda}$ and a λ -twisted vector bundle isomorphism $\varphi = \{\varphi_i\} : E \oplus F \rightarrow \overline{E} \oplus F$ such that

$$\text{CS}(\Gamma \oplus \Gamma^F, \varphi^*(\overline{\Gamma} \oplus \Gamma^F)) = \omega - \overline{\omega}.$$

We verify that (E', Γ', ω') and $(\overline{E}', \overline{\Gamma}', \overline{\omega}')$ are equivalent so that well-definedness of the map follows. We take a λ' -twisted vector bundle F' and a connection Γ'^F compatible with $\widehat{\lambda}'$ by applying the same rule in (3) to (F, Γ^F) . There exists a λ' -twisted vector bundle isomorphism $\varphi = \{\varphi_i\} : E' \oplus F' \rightarrow \overline{E}' \oplus F'$ defined exactly the same as the above φ .¹ We have to show that

$$\text{CS}(\Gamma' \oplus \Gamma'^F, \varphi^*(\overline{\Gamma}' \oplus \Gamma'^F)) = \omega' - \overline{\omega}'.$$

Suppose $\widetilde{\Gamma}$ is a connection on $p^*(E \oplus F)$ over $X \times I$ defined by a path of connections joining $\Gamma \oplus \Gamma^F$ and $\varphi^*(\overline{\Gamma} \oplus \Gamma^F)$ on $E \oplus F$ over X . By definition,

$$\text{cs}(\Gamma^t) := \int_I \text{ch}(\widetilde{\Gamma}) = \int_I \text{rank}(E \oplus F) + \sum_{m=1}^\infty \frac{1}{m!} \text{tr}(\widetilde{R}_i - p^* B_i \cdot \mathbf{1})^m.$$

¹ Let $\overline{g}_{ji}, h_{ji}$ be transition maps of \overline{E} and F , respectively. Since φ is an isomorphism, we have

$$\varphi_j(x)(\overline{g}_{ji}(x) \oplus h_{ji}(x)) = (\overline{g}_{ji}(x) \oplus h_{ji}(x))\varphi_i(x)$$

for all $x \in U_{ij}$. From this we have

$$\varphi_j(x)(\overline{g}_{ji}(x)\chi_{ji}(x) \oplus h_{ji}(x)\chi_{ji}(x)) = (\overline{g}_{ji}(x)\chi_{ji}(x) \oplus h_{ji}(x)\chi_{ji}(x))\varphi_i(x)$$

for all $x \in U_{ij}$.

We see that:

$$\begin{aligned}
 \text{CS}(\Gamma' \oplus \Gamma'^F, \varphi^*(\overline{\Gamma}' \oplus \Gamma'^F)) &= \text{CS}(\Gamma \oplus \Gamma^F + \Pi \cdot \mathbf{1}, \varphi^*(\overline{\Gamma} \oplus \Gamma^F + \Pi \cdot \mathbf{1})) \\
 &= \text{CS}(\Gamma \oplus \Gamma^F + \Pi \cdot \mathbf{1}, \varphi^*(\overline{\Gamma} \oplus \Gamma^F) + \Pi \cdot \mathbf{1}) \\
 &= \text{cs}(\Gamma^t + \Pi \cdot \mathbf{1}) \pmod{\text{Im}(d + H)} \\
 &= \int_I \text{ch}(\tilde{\Gamma} + p^* \Pi \cdot \mathbf{1}) \pmod{\text{Im}(d + H)}. \tag{4}
 \end{aligned}$$

Since

$$d(\tilde{\Gamma}_i + p^* \Pi_i \cdot \mathbf{1}) + (\tilde{\Gamma}_i + p^* \Pi_i \cdot \mathbf{1}) \wedge (\tilde{\Gamma}_i + p^* \Pi_i \cdot \mathbf{1}) = \tilde{R}_i + p^* d\Pi_i \cdot \mathbf{1},$$

we have

$$\begin{aligned}
 \int_I \text{ch}(\tilde{\Gamma} + p^* \Pi \cdot \mathbf{1}) &= \int_I \text{rank}(E' \oplus F') + \sum_{m=1}^{\infty} \frac{1}{m!} \text{tr}(\tilde{R}_i + p^* d\Pi_i \cdot \mathbf{1} - p^* B'_i \cdot \mathbf{1})^m \\
 &= \int_I \text{rank}(E \oplus F) + \sum_{m=1}^{\infty} \frac{1}{m!} \text{tr}(\tilde{R}_i - p^* B_i \cdot \mathbf{1})^m \\
 &= \int_I \text{ch}(\tilde{\Gamma}).
 \end{aligned}$$

Hence the far RHS of (4) is

$$\int_I \text{ch}(\tilde{\Gamma})/\text{Im}(d + H) = \text{CS}(\Gamma \oplus \Gamma^F, \varphi^*(\overline{\Gamma} \oplus \Gamma^F)) = \omega - \bar{\omega}.$$

The map $\phi_{\hat{\alpha}}$ being bijective, a group homomorphism and natural in \mathcal{U} is straightforward.

(2) We first show that, if $(E, \Gamma, \omega) \sim (\overline{E}, \overline{\Gamma}, \bar{\omega})$, then $(E, \Gamma_{\xi}, \omega \wedge \exp(-\xi)) \sim (\overline{E}, \overline{\Gamma}_{\xi}, \bar{\omega} \wedge \exp(-\xi))$. By the premise, there exists a λ -twisted vector bundle G and a connection Γ^G on G compatible with $\hat{\lambda}$ and an isomorphism $\varphi : E \oplus G \rightarrow \overline{E} \oplus G$, such that $\omega - \bar{\omega} = \text{CS}(\Gamma \oplus \Gamma^G, \varphi^*(\overline{\Gamma} \oplus \Gamma^G)) = \int_I \text{ch}(\tilde{\Gamma}) \pmod{\text{Im}(d + H)}$, where $\tilde{\Gamma}$ is a connection on $p^*(E \oplus G)$ defined by pullback of connections on a straight line path joining $\Gamma \oplus \Gamma^G$ and $\varphi^*(\overline{\Gamma} \oplus \Gamma^G)$. Accordingly,

$$\begin{aligned}
 \text{CS}(\Gamma_{\xi} \oplus \Gamma_{\xi}^G, \varphi^*(\overline{\Gamma}_{\xi} \oplus \Gamma_{\xi}^G)) &= \int_I (\text{ch}(\tilde{\Gamma}) \wedge \exp(-p^* \xi)) \pmod{\text{Im}(d + H + d\xi)} \\
 &= \left(\int_I \text{ch}(\tilde{\Gamma}) \right) \wedge \exp(-\xi) \pmod{\text{Im}(d + H + d\xi)} \\
 &= (\omega - \bar{\omega}) \wedge \exp(-\xi) \pmod{\text{Im}(d + H + d\xi)}.
 \end{aligned}$$

From this, well-definedness of the map \mathcal{E} follows. The map \mathcal{E} being one-to one, onto, group homomorphism, and natural in \mathcal{U} are all obvious. □

4.5 The I and R map

Proposition 4.20 *Let $\widehat{\lambda} \in \text{Twist}_{\widehat{K}}^{\text{tor}}(\mathcal{U})$. The assignment*

$$I : \widehat{K}^0(\mathcal{U}, \widehat{\lambda}) \rightarrow K^0(\mathcal{U}, \lambda)$$

$$[(E, \Gamma^E, \omega)] - [(F, \Gamma^F, \eta)] \mapsto [E] - [F]$$

is a group homomorphism which is natural in \mathcal{U} .

Proposition 4.21 *Let $\widehat{\lambda} \in \text{Twist}_{\widehat{K}}^{\text{tor}}(\mathcal{U})$. The assignment*

$$R : \widehat{K}^0(\mathcal{U}, \widehat{\lambda}) \rightarrow \Omega^{\text{even}}(X; \mathbb{C})$$

$$[(E, \Gamma^E, \omega)] - [(F, \Gamma^F, \eta)] \mapsto \text{ch}(\Gamma^E) + (d + H)\omega - \text{ch}(\Gamma^F) - (d + H)\eta$$

is a group homomorphism which is natural in \mathcal{U} .

4.6 Odd twisted Chern character forms

In this subsection, we define odd twisted Chern character forms which will be used in Sects. 4.7, 4.8, and 4.9.

Definition 4.22 Let X be a manifold, \mathcal{U} a good open cover of X , $\widehat{\lambda} = (\{\lambda_{kji}\}, \{A_{ji}\}, \{B_i\})$ a $U(1)$ -gerbe with connection on \mathcal{U} whose Dixmier–Douady class is torsion. Also let E be a λ -twisted vector bundle, ϕ an automorphism on E , and Γ a connection compatible with $\widehat{\lambda}$. The **total twisted odd Chern character form** of the triple (E, ϕ, Γ) is $\text{cs}(t \mapsto (1 - t)\Gamma^E + t\phi^*\Gamma^E)$.

By Propositions 3.26 and 3.14, $\text{cs}(t \mapsto (1 - t)\Gamma^E + t\phi^*\Gamma^E)$ represents an odd twisted cohomology class.

The odd twisted Chern character form is functorial.

Proposition 4.23 *Given a map $f : (Y, \mathcal{V}) \rightarrow (X, \mathcal{U})$ with $\mathcal{V} = f^{-1}(\mathcal{U})$, the following holds:*

$$\text{Ch}(f^*E, (\phi \circ f), f^*\Gamma) = f^*\text{Ch}(E, \phi, \Gamma)$$

Proof Note that $f^*\phi^*\Gamma^E = (\phi \circ f)^{-1} \circ f^*\Gamma^E \circ (\phi \circ f) + (\phi \circ f)^{-1}d(\phi \circ f) = (\phi \circ f)^*\Gamma^E$. The proof of this statement is similar to the proof of Lemma 4.15. \square

The total odd twisted Chern character form respects change of differential twist in a manner that is similar to the even case. (Compare Propositions 3.16 and 3.19.)

Proposition 4.24 (1) *Let $\widehat{\lambda}, \widehat{\lambda}', E, E', \Gamma$, and Γ' be as in Proposition 3.16. The following holds:*

$$\text{Ch}(E', \phi, \Gamma') = \text{Ch}(E, \phi, \Gamma)$$

(2) Let $\xi \in \Omega^2(X; i\mathbb{R})$, and $\widehat{\lambda}$ and $\widehat{\lambda}_\xi$ be as in Notation 3.18. The following holds:

$$\text{Ch}(E, \phi, \Gamma_\xi) = \text{Ch}(E, \phi, \Gamma) \wedge \exp(-\xi)$$

Proof (1) $\text{Ch}(E, \phi, \Gamma') = \text{cs}(t \mapsto (1-t)\Gamma' + t\phi^*\Gamma') = \text{cs}(t \mapsto (1-t)\Gamma + t\phi^*\Gamma + \Pi \cdot \mathbf{1}) = \text{cs}(t \mapsto (1-t)\Gamma + t\phi^*\Gamma) = \text{Ch}(E, \phi, \Gamma)$, where the third equality follows from a similar calculation appearing in the proof of Proposition 4.18.

(2) Let $\widetilde{\Gamma}$ be a connection on p^*E defined by pullback of connections on the path $(1-t)\Gamma^E + t\phi^*\Gamma^E$. We have $\text{Ch}(E, \phi, \Gamma_\xi^E) = \text{cs}(t \mapsto (1-t)\Gamma_\xi^E + t\phi^*\Gamma_\xi^E) = \int_I \text{ch}(\widetilde{\Gamma}) \wedge \exp(-p^*\xi) = (\int_I \text{ch}(\widetilde{\Gamma})) \wedge \exp(-\xi) = \text{Ch}(E, \phi, \Gamma^E) \wedge \exp(-\xi)$. □

4.7 The a map and the exact sequence involving the a and I maps

Notation 4.25 We denote by $\Omega_{H,\text{Ch}}$ the abelian group $\text{Im}(\text{Ch}) + \text{Im}(d + H)$, where $\text{Im}(\text{Ch})$ is the abelian group generated by all odd twisted Chern character forms.

Definition 4.26 Let $\widehat{\lambda} \in \text{Twist}_{\widehat{K}}^{\text{tor}}(\mathcal{U})$. Define:

$$a : \Omega^{\text{odd}}(X; \mathbb{C})/\Omega_{H,\text{Ch}} \rightarrow \widehat{K}^0(\mathcal{U}, \widehat{\lambda})$$

$$\theta \mapsto [(\mathcal{O}, 0, \theta)].$$

Lemma 4.27 *The map a is well-defined, a group homomorphism and is natural in \mathcal{U} .*

Proof We prove well-definedness.

$$\begin{aligned} a(\theta + \text{Ch}(E, \phi, \Gamma^E)) &= [(\mathcal{O}, 0, \theta + \text{Ch}(E, \phi, \Gamma^E))] \\ &= [(\mathcal{O} \oplus E, 0 \oplus \Gamma^E, \theta + \text{cs}((1-t)\Gamma^E + t\phi^*\Gamma^E))] - [(E, \Gamma^E, 0)] \\ &= [(\mathcal{O}, 0, \theta)] + [(E, \Gamma^E, \text{cs}((1-t)\Gamma^E + t\phi^*\Gamma^E))] - [(E, \Gamma^E, 0)] \\ &= [(\mathcal{O}, 0, \theta)] = a(\theta). \end{aligned}$$

□

Proposition 4.28 *The following sequence is exact:*

$$0 \rightarrow \Omega^{\text{odd}}(X; \mathbb{C})/\Omega_{H,\text{Ch}} \xrightarrow{a} \widehat{K}^0(\mathcal{U}, \widehat{\lambda}) \xrightarrow{I} K^0(\mathcal{U}, \lambda) \rightarrow 0.$$

Proof It is obvious that I is surjective, and $\text{Im } a \subseteq \ker I$. We show the other inclusion.

Let $[(E, \Gamma^E, \omega)] - [(F, \Gamma^F, \eta)] \in \widehat{K}^0(\mathcal{U}, \widehat{\lambda})$ and suppose $I([(E, \Gamma^E, \omega)] - [(F, \Gamma^F, \eta)]) = 0$. Then there exists a λ -twisted vector bundle G and an isomorphism

of λ -twisted vector bundles $\varphi : E \oplus G \rightarrow F \oplus G$. Choose any connection Γ^G on G that is compatible with $\widehat{\lambda}$. Then

$$\begin{aligned}
 [(E, \Gamma^E, \omega)] - [(F, \Gamma^F, \eta)] &= [(E \oplus G, \Gamma^E \oplus \Gamma^G, \omega)] - [(F \oplus G, \Gamma^F \oplus \Gamma^G, \eta)] \\
 &= [(E \oplus G, \Gamma^E \oplus \Gamma^G, \omega)] - [(E \oplus G, \Gamma^E \oplus \Gamma^G, \mu)],
 \end{aligned}$$

where $\mu := \eta + \text{CS}(\Gamma^E \oplus \Gamma^G, \varphi^*(\Gamma^F \oplus \Gamma^G))$, and $*$ follows from the fact that $(E \oplus G, \Gamma^E \oplus \Gamma^G, \mu)$ is equivalent to $(F \oplus G, \Gamma^F \oplus \Gamma^G, \eta)$. We now add and subtract $(\mathcal{O}, 0, 0)$, and get:

$$\begin{aligned}
 &[(E \oplus G, \Gamma^E \oplus \Gamma^G, \omega)] - [(E \oplus G, \Gamma^E \oplus \Gamma^G, \mu)] \\
 &= [(E \oplus G \oplus \mathcal{O}, \Gamma^E \oplus \Gamma^G \oplus 0, \omega)] - [(E \oplus G \oplus \mathcal{O}, \Gamma^E \oplus \Gamma^G \oplus 0, \mu)] \\
 &= [(E \oplus G, \Gamma^E \oplus \Gamma^G, 0)] + [(\mathcal{O}, 0, \omega)] - [(E \oplus G, \Gamma^E \oplus \Gamma^G, 0)] - [(\mathcal{O}, 0, \mu)] \\
 &= [(\mathcal{O}, 0, \omega - \mu)] = a(\omega - \mu).
 \end{aligned}$$

We now show injectivity of the a map. Suppose $a(\theta) = 0$. Equivalently, there exists a λ -twisted vector bundle with connection (E, Γ^E) whose connection is compatible with $\widehat{\lambda}$ and an automorphism $\varphi \in \text{Aut}(E)$ satisfying that $\theta = \text{CS}(\Gamma^E, \varphi^*\Gamma^E) = \text{Ch}(E, \varphi, \Gamma^E) \pmod{\text{Im}(d + H)}$. Hence the result. \square

Notation 4.29 Let $\mathcal{U} := \{U_i\}_{i \in \Lambda}$ be an open cover of a space X . Recall that a *refinement* of \mathcal{U} is a pair (\mathcal{V}, τ) consisting of an open cover $\mathcal{V} := \{V_r\}_{r \in \mathcal{L}}$ of X and a map $\tau : \mathcal{L} \rightarrow \Lambda$ such that $V_r \subset U_{\tau(r)}$ for all $r \in \mathcal{L}$. It induces the following restriction maps on the totality of $U(1)$ -gerbes with connection, λ -twisted vector bundles, and the space of connections on a λ -twisted vector bundle E , respectively.

$$\begin{aligned}
 \widehat{\lambda} = (\lambda_{kji}, A_{ji}, B_i) &\mapsto \tau^*\widehat{\lambda} = (\tau^*\lambda_{tsr}, \tau^*A_{sr}, \tau^*B_r) := (\lambda_{\tau(t)\tau(s)\tau(r)}, A_{\tau(s)\tau(r)}, B_{\tau(r)}) \\
 E = (\mathcal{U}, \{g_{ji}\}, \{\lambda_{kji}\}) &\mapsto \tau^*E := (\mathcal{V}, \{g_{\tau(s)\tau(r)}\}, \{\tau^*\lambda_{tsr}\}) \\
 \Gamma = \{\Gamma_i\}_{i \in \Lambda} &\mapsto \tau^*\Gamma = \{(\tau^*\Gamma)_r := \Gamma_{\tau(r)}\}_{r \in \mathcal{L}}
 \end{aligned}$$

Corollary 4.30 Let $\mathcal{U} := \{U_i\}_{i \in \Lambda}$ be a good cover of X and (\mathcal{V}, τ) consisting of an open cover $\mathcal{V} := \{V_r\}_{r \in \mathcal{L}}$ of X and a map $\tau : \mathcal{L} \rightarrow \Lambda$ be a choice of refinement of \mathcal{U} . The restriction map induced by τ

$$\begin{aligned}
 \widehat{R}_\tau : \widehat{K}^0(\mathcal{U}; \widehat{\lambda}) &\rightarrow \widehat{K}^0(\mathcal{V}; \tau^*\widehat{\lambda}) \\
 [(E, \Gamma^E, \omega)] - [(F, \Gamma^F, \eta)] &\mapsto [(\tau^*F, \tau^*\Gamma^E, \omega)] - [(\tau^*F, \tau^*\Gamma^F, \eta)]
 \end{aligned}$$

is an isomorphism of abelian groups.

Proof Consider the following diagram.

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \Omega^{\text{odd}}(X; \mathbb{C})/\Omega_{H,\text{Ch}} & \xrightarrow{a} & \widehat{K}^0(\mathcal{U}, \widehat{\lambda}) & \xrightarrow{I} & K^0(\mathcal{U}; \lambda) \longrightarrow 0 \\
 \parallel & & \parallel & & \downarrow \widehat{R}_\tau & & \cong \downarrow R_\tau \\
 0 & \longrightarrow & \Omega^{\text{odd}}(X; \mathbb{C})/\Omega_{H,\text{Ch}} & \xrightarrow{a} & \widehat{K}^0(\mathcal{V}, \tau^*\widehat{\lambda}) & \xrightarrow{I} & K^0(\mathcal{V}; \tau^*\lambda) \longrightarrow 0
 \end{array}$$

The map R_τ is an isomorphism (See Karoubi [19, p. 233] Theorem 3.6), and all square diagrams commute. Hence \widehat{R}_τ is an isomorphism by the five-lemma. \square

Definition 4.31 Let $\widehat{\lambda}$ be the element in the colimit of Deligne 2-cocycles over X defined on a good open cover \mathcal{U} along a choice of refinements of \mathcal{U} represented by $\widehat{\lambda} \in \text{Twist}_K^{\text{tor}}(\mathcal{U})$. The **twisted differential K -group** of X , denoted by $\widehat{K}^0(X, \widehat{\lambda})$, is defined by the colimit of $\widehat{K}^0(\mathcal{U}, \widehat{\lambda})$ over all refinements of \mathcal{U} . The **twisted K -group** of X , denoted by $K^0(X, \underline{\lambda})$, is defined by the colimit of $K^0(\mathcal{U}, \lambda)$ over all refinements of \mathcal{U} .

4.8 The hexagon diagram

Notation 4.32 We denote by $\text{Pr} : \Omega^{\text{even}}(X; \mathbb{C})_{\text{closed}} \rightarrow H_H^{\text{even}}(X; \mathbb{C})$ the map taking twisted de Rham cohomology class, and $r : H_H^{\text{odd}}(X; \mathbb{C}) \rightarrow \Omega^{\text{odd}}(X; \mathbb{C})/\Omega_{H, \text{Ch}}$ the map that sends an odd twisted de Rham cohomology class $[\omega]$ to $\omega + \Omega_{H, \text{Ch}}$. The map r is well-defined by definition of $\omega + \Omega_{H, \text{Ch}}$ (see Notation 4.25).

Proposition 4.33 For the maps I, R , and a from or into $\widehat{K}^0(\mathcal{U}; \widehat{\lambda})$, the following holds:

- (1) $ch \circ I = \text{Pr} \circ R$.
- (2) $R \circ a = d + H$.

Definition 4.34 We define maps α and β as follows:

$$\begin{aligned} \alpha : H_H^{\text{odd}}(X; \mathbb{C}) &\rightarrow \ker R & \beta : \ker R &\rightarrow K^0(\mathcal{U}, \lambda) \\ [\omega] &\mapsto (\mathcal{O}, 0, \omega) & (E, \Gamma^E, \omega) - (F, \Gamma^F, \eta) &\mapsto [E] - [F]. \end{aligned}$$

Remark 4.35 The maps α and β are well-defined group homomorphisms.

- Proposition 4.36**
- (1) $a \circ r = \text{incl} \circ \alpha$.
 - (2) $\beta = I \circ \text{incl}$.
 - (3) The following sequences are exact:

$$\begin{aligned} H_H^{\text{odd}}(X; \mathbb{C}) &\xrightarrow{\alpha} \ker R \xrightarrow{\beta} K^0(\mathcal{U}, \lambda) \xrightarrow{ch} H_H^{\text{even}}(X; \mathbb{C}) \\ H_H^{\text{odd}}(X; \mathbb{C}) &\xrightarrow{r} \Omega^{\text{odd}}(X; \mathbb{C})/\text{Im}(d + H) \xrightarrow{d+H} \Omega^{\text{even}}(X; \mathbb{C})_{\text{closed}} \xrightarrow{\text{Pr}} H_H^{\text{even}}(X; \mathbb{C}) \end{aligned}$$

Proof All claims are obvious except that $\ker(\beta) \subseteq \text{Im}(\alpha)$, which we prove presently. Take an arbitrary element $[(E, \Gamma^E, \omega)] - [(F, \Gamma^F, \eta)] \in \ker R$ whose image under β is zero, i.e., there exists a λ -twisted vector bundle G defined on \mathcal{U} and an isomorphism $\varphi : E \oplus G \xrightarrow{\cong} F \oplus G$. Choose any connection Γ^G on G that is compatible with $\widehat{\lambda}$. At this point we introduce the following notation:

$$\zeta := \text{CS}(\Gamma^E \oplus \Gamma^G, \varphi^*(\Gamma^F \oplus \Gamma^G)).$$


We see that

$$\begin{aligned}
 [(E, \Gamma^E, \omega)] - [(F, \Gamma^F, \eta)] &= [(E, \Gamma^E, \omega)] - [(F, \Gamma^F, \omega - \zeta)] + [(\mathcal{O}, 0, \omega - \zeta - \eta)] \\
 &= a([\omega - \zeta - \eta]),
 \end{aligned}
 \tag{5}$$

where in the first equality we add and subtract $[(\mathcal{O}, 0, \omega - \zeta - \eta)]$ and in the second use Definition 4.10.

We have to verify that the differential form $\omega - \eta - \zeta$ represents an odd degree twisted cohomology class. Since $[(E, \Gamma^E, \omega)] - [(F, \Gamma^F, \eta)] \in \ker R$, we have $\text{ch}(\Gamma^E) - \text{ch}(\Gamma^F) + (d + H)(\omega - \eta) = 0$. Now $(d + H)(\omega - \eta - \zeta) = \text{ch}(\Gamma^E) - \text{ch}(\Gamma^F) - (\text{ch}(\Gamma^E \oplus \Gamma^G) - \text{ch}(\Gamma^F \oplus \Gamma^G)) = 0$. \square

Corollary 4.37 *In the following diagram for $\widehat{K}^0(X; \widehat{\lambda})$, all square and triangles are commutative and all sequences are exact.*



Remark 4.38 When the differential twist is $\widehat{\lambda} = (\{1\}, \{0\}, \{0\})$, the diagram reduces to the differential K -theory hexagon diagram of Simons and Sullivan (see [24, p. 596]).

4.9 Compatibility with change of twist map

Proposition 4.39 *Let $\widehat{\alpha}' = (\chi'_{ji}, \Pi'_i)$ be an isomorphism $\widehat{\lambda} \rightarrow \widehat{\lambda}'$ such that $\widehat{\lambda}' = \widehat{\lambda} + D\alpha'$. Then the diagram in Corollary 4.37 is natural under change of twist by $\widehat{\alpha}'$:*

- (1) $I_{\widehat{\lambda}'} \circ \phi_{\widehat{\alpha}'} = \phi_{\alpha'} \circ I_{\widehat{\lambda}}$.
- (2) $\phi_{\widehat{\alpha}'} \circ a_{\widehat{\lambda}} = a_{\widehat{\lambda}'}$.
- (3) $R_{\widehat{\lambda}'} \circ \phi_{\widehat{\alpha}'} = R_{\widehat{\lambda}}$.

Proof (1)

$$\begin{aligned}
 I_{\widehat{\lambda}'} \circ \phi_{\widehat{\alpha}'}([E, \Gamma^E, \omega] - [F, \Gamma^F, \eta]) &= I_{\widehat{\lambda}'}([\phi_{\alpha'} E, \Gamma^E + \Pi' \cdot \mathbf{1}, \omega] \\
 &\quad - [\phi_{\alpha'} F, \Gamma^F + \Pi' \cdot \mathbf{1}, \eta]) \\
 &= [\phi_{\alpha'} E] - [\phi_{\alpha'} F] \\
 &= \phi_{\alpha'} \circ I_{\widehat{\lambda}}([E, \Gamma^E, \omega] - [F, \Gamma^F, \eta]).
 \end{aligned}$$

(2) Obvious.

(3)

$$\begin{aligned}
 R_{\widehat{\lambda}'} \circ \phi_{\alpha'}([E, \Gamma^E, \omega] - [F, \Gamma^F, \eta]) &= R_{\widehat{\lambda}}([\phi_{\alpha'} E, \{\Gamma_i^E + \Pi'_i\}, \omega] - [\phi_{\alpha'} F, \{\Gamma_i^F + \Pi'_i\}, \eta]) \\
 &= (\text{ch}(\Gamma^E + \Pi' \cdot \mathbf{1}) + (d + H)\omega - \text{ch}(\Gamma^F + \Pi' \cdot \mathbf{1}) - (d + H)\eta) \\
 &= (\text{ch}(\Gamma^E) + (d + H)\omega - \text{ch}(\Gamma^F) - (d + H)\eta) \quad \text{by Lemma 3.16.} \\
 &= R_{\widehat{\lambda}}([E, \Gamma^E, \omega] - [F, \Gamma^F, \eta]).
 \end{aligned}$$

□

Proposition 4.40 *Let $\xi \in \Omega^2(X; i\mathbb{R})$ that induces an isomorphism $\widehat{\lambda} \rightarrow \widehat{\lambda}_\xi$. Then the diagram in Corollary 4.37 is natural under change of twist by ξ :*

- (1) $\mathcal{E} \circ a = a \circ \exp(-\xi)$.
- (2) $I \circ \mathcal{E} = I$.
- (3) $R \circ \mathcal{E} = \exp(-\xi) \circ R$.

Acknowledgements We would like to thank Mahmoud Zeinalian for many helpful discussions and comments as well as constant support during the entire process of this work. We also would like to thank Scott Wilson, Arthur Parzygnat, and Corbett Redden for reading the preliminary version of this paper and providing helpful suggestions and comments as well as Thomas Tradler and Cheyne Miller for useful conversations. We thank Jim Stasheff, Hisham Sati, Ping Xu, and Mathieu Stiénon for their interest and detailed comments on this work. Finally, we thank the anonymous referee for detailed comments.

References

1. Atiyah, M., Segal, G.: Twisted K -theory and cohomology. In: Griffiths, P.A. (ed.) *Inspired by S. S. Chern*, p. 543. World Sci. Publ., Hackensack (2006)
2. Bott, R., Tu, L.W.: *Differential Forms in Algebraic Topology*. Springer, New York (1982)
3. Bouwknegt, P., Carey, A.L., Mathai, V., Murray, M.K., Stevenson, D.: Twisted K -theory and K -theory of bundle gerbes. *Commun. Math. Phys.* **228**, 17–45 (2002). doi:[10.1007/s002200200646](https://doi.org/10.1007/s002200200646)
4. Brylinski, J.-L.: *Loop Spaces, Characteristic Classes and Geometric Quantization*. Birkhuser Boston Inc., Boston (2008)
5. Bunke, U., Nikolaus, T.: Twisted differential cohomology. Arxiv e-print. (2014). [arXiv:1406.3231v1](https://arxiv.org/abs/1406.3231v1) [math.AT]
6. Bunke, U., Nikolaus, T., Völkl, M.: Differential cohomology theories as sheaves of spectra. *J. Homotopy Relat. Struct.* **11**, 1–66 (2016). doi:[10.1007/s40062-014-0092-5](https://doi.org/10.1007/s40062-014-0092-5)
7. Bunke, U., Schick, T.: Smooth K -theory. *Astrisque* **328**, 45–135 (2009)
8. Bunke, U., Schick, T.: Differential K -theory: a survey. In: Bär, C., Lohkamp, J., Schwarz, M. (eds.) *Global Differential Geometry*, pp. 303–357. Springer, Heidelberg (2012)

9. Carey, A.L., Mickelsson, J., Wang, B.-L.: Differential twisted K -theory and applications. *J. Geom. Phys.* **59**, 632–653 (2009). doi:[10.1016/j.geomphys.2009.02.002](https://doi.org/10.1016/j.geomphys.2009.02.002)
10. Donovan, P., Karoubi, M.: Graded Brauer groups and K -theory with local coefficients. *Inst. Hautes Etudes Sci. Publ. Math.* **38**, 5–25 (1970)
11. Freed, D.S.: Dirac charge quantization and generalized differential cohomology. In: Yau, S.-T. (ed.) *Surveys in Differential Geometry*, pp. 129–194. Int. Press, Somerville (2000)
12. Freed, D.S., Hopkins, M.J., Teleman, C.: Loop groups and twisted K -theory I. *J. Topol.* **4**, 737–798 (2011). doi:[10.1112/jtopol/jtr019](https://doi.org/10.1112/jtopol/jtr019)
13. Freed, D.S., Lott, J.: An index theorem in differential K -theory. *Geom. Topol.* **14**, 903–966 (2010). doi:[10.2140/gt.2010.14.903](https://doi.org/10.2140/gt.2010.14.903)
14. Freed, D.S., Moore, G.W.: Twisted equivariant matter. *Ann. Henri Poincaré* **14**, 1927–2023 (2013). doi:[10.1007/s00023-013-0236-x](https://doi.org/10.1007/s00023-013-0236-x)
15. Gawędzki, K., Reis, N.: WZW branes and gerbes. *Rev. Math. Phys.* **14**, 1281–1334 (2002). doi:[10.1142/S0129055X02001557](https://doi.org/10.1142/S0129055X02001557)
16. Hopkins, M.J., Singer, I.M.: Quadratic functions in geometry, topology, and M -theory. *J. Differ. Geom.* **70**, 329–452 (2005)
17. Kahle, A., Valentino, A.: T -duality and differential K -theory. *Commun. Contemp. Math.* **16**, (2014). doi:[10.1142/S0219199713500144](https://doi.org/10.1142/S0219199713500144)
18. Karoubi, M.: Homologie cyclique et K -thorie. *Astérisque J. Lond. Math. Soc.* **149**, 1–147 (1987)
19. Karoubi, M.: Twisted bundles and twisted K -theory. In: Cortiñas, G. (ed.) *Topics in Noncommutative Geometry*, pp. 223–257. Amer. Math. Soc., Providence (2012)
20. Klonoff, K.R.: An index theorem in differential K -theory. ProQuest LLC, Ann Arbor (2008)
21. Lott, J.: R/Z index theory. *Commun. Anal. Geom.* **2**, 279–311 (1994). doi:[10.4310/CAG.1994.v2.n2.a6](https://doi.org/10.4310/CAG.1994.v2.n2.a6)
22. Murray, M.K.: Bundle gerbes. *J. Lond. Math. Soc.* **54**(2), 403–416 (1996). doi:[10.1093/acprof:oso/9780199534920.003.0012](https://doi.org/10.1093/acprof:oso/9780199534920.003.0012)
23. Rosenberg, J.: Continuous-trace algebras from the bundle theoretic point of view. *J. Aust. Math. Soc. Ser. A* **47**, 368–381 (1989)
24. Simons, J., Sullivan, D.: Structured vector bundles define differential K -theory. In: Blanchard, É., Ellwood, D., Khalkhali, M., Marcolli, M., Moscovici, H., Popa, S. (eds.) *Quanta of maths*, pp. 579–599. Amer. Math. Soc., Providence (2010)
25. Tradler, T., Wilson, S.O., Zeinalian, M.: An elementary differential extension of odd K -theory. *J. K-Theory* **12**, 331–361 (2013). doi:[10.1017/is013002018jkt218](https://doi.org/10.1017/is013002018jkt218)
26. Tradler, T., Wilson, S.O., Zeinalian, M.: Differential K -theory as equivalence classes of maps to Grassmannians and unitary groups. *N. Y. J. Math.* **22**, 527–581 (2016)
27. Witten, E.: D-branes and K -theory. *J. High Energy Phys.* **19** (1998). doi:[10.1088/1126-6708/1998/12/019](https://doi.org/10.1088/1126-6708/1998/12/019)