



Unstable splittings in Hodge filtered Brown–Peterson cohomology

Gereon Quick¹

Received: 7 December 2016 / Accepted: 7 September 2018 / Published online: 27 September 2018
© Tbilisi Centre for Mathematical Sciences 2018

Abstract

We construct Hodge filtered function spaces associated to infinite loop spaces. For Brown–Peterson cohomology, we show that the corresponding Hodge filtered spaces satisfy an analog of Wilson’s unstable splitting. As a consequence, we obtain an analog of Quillen’s theorem for Hodge filtered Brown–Peterson cohomology for complex manifolds.

Keywords Homotopy theory · Deligne cohomology · Brown-Peterson spectra · Hodge filtered spaces · Quillen theorem · Wilson splitting

1 Introduction

For a fixed prime number p , let $BP^*(-)$ denote Brown–Peterson cohomology and let BP be the Ω -spectrum representing $BP^*(-)$ (see [1, 14]). For an integer $n \geq 0$, let $BP\langle n \rangle$ be the Ω -spectrum representing the associated intermediate cohomology theory studied by Wilson in [17]. These spectra are connected via canonical maps $g_n: BP \rightarrow BP\langle n \rangle$ and $f_n: BP\langle n \rangle \rightarrow BP\langle n-1 \rangle$. The coefficient rings of BP and $BP\langle n \rangle$ are isomorphic to polynomial algebras

$$BP^* = \mathbb{Z}_{(p)}[v_1, v_2, \dots] \quad \text{and} \quad BP\langle n \rangle^* = \mathbb{Z}_{(p)}[v_1, \dots, v_n]$$

where the generator v_i has degree $-2(p^i - 1)$. The effect of g_n on coefficients is to send the generators v_i with $i \geq n + 1$ to 0, and f_n sends v_n to 0.

Let BP_k and $BP\langle n \rangle_k$ denote the k th spaces of the spectra BP and $BP\langle n \rangle$, respectively. In [17], Wilson proved that, for $k \leq 2(p^n + p^{n-1} + \dots + p + 1)$, there is a homotopy equivalence

Communicated by Stephen Wilson.

✉ Gereon Quick
gereon.quick@ntnu.no

¹ Department of Mathematical Sciences, NTNU, 7491 Trondheim, Norway

$$BP_k \xrightarrow{\approx} BP\langle n \rangle_k \times \prod_{j>n} BP\langle j \rangle_{k+2(p^j-1)}, \tag{1}$$

and the product cannot be broken down further. This result has many important consequences. For example, it yields a proof of Quillen’s theorem that, for a finite complex X , $BP^*(X)$ is generated as a BP^* -module by elements of non-negative degree [17, Theorem 5.7].

In [8], Hopkins and the author showed that, given any rationally even spectrum E , there is an associated Hodge filtered cohomology theory $E_{\mathcal{D}}^*(*)(-)$ for complex manifolds represented by a presheaf of spectra $E_{\mathcal{D}}$. These Hodge filtered cohomology theories are natural generalizations of analytic Deligne cohomology for complex manifolds. In particular, there is a Hodge filtered BP -theory represented by a presheaf of spectra $BP_{\mathcal{D}}$. The purpose of this paper is to show that Wilson’s splitting (1) of the spaces BP_k induces a splitting of the spaces, i.e. simplicial presheaves, in the spectrum $BP_{\mathcal{D}}$.

It is important to note that the splitting (1) does not exist on the level of spectra (see [2, p. 817]). Hence the first important step is to construct Hodge filtered spaces associated to BP . The presheaf of spectra $BP_{\mathcal{D}}$ is actually given by a wedge sum of presheaves of spectra $BP_{\mathcal{D}}(m)$, one summand for each integer m . For all integers m and $k \geq 0$, we construct simplicial presheaves $BP_k(m)$ on the Grothendieck site of complex manifolds together with weak equivalences $\sigma_k : BP_k(m) \rightarrow \Omega BP_{k+1}(m)$ such that the presheaf of spectra $\{BP_k(m)\}_k$ built by the $BP_k(m)$ and σ_k is equivalent to $BP_{\mathcal{D}}(m)$. In fact, we will construct spaces $E_k(m)$ for any connective rationally even spectrum E . We call $E_k(m)$ the m th Hodge filtered function space associated to E_k . For the reader who is familiar with differential cohomology theories, we remark that these constructions are similar to how Hopkins-Singer define differential function spaces for smooth manifolds in [9].

The second important step is to show that a map $E_k \rightarrow F_n$ between spaces of rationally even spectra E and F induces a map $E_k(m) \rightarrow F_n(m)$ of simplicial presheaves. The analog of Wilson’s splitting will then be a natural consequence of the constructions.

Our main results are the following.

Theorem 1.1 *Let m and n be integers with $n \geq 0$. For every $k \leq 2(p^n + \dots + 1)$, there is a weak equivalence of simplicial presheaves*

$$BP_k(m) \xrightarrow{\approx} BP\langle n \rangle_k(m) \times \prod_{j>n} BP\langle j \rangle_{k+2(p^j-1)}(m). \tag{2}$$

It follows, in particular, that, for every complex manifold M , the natural map

$$BP_{\mathcal{D}}^k(m)(M) \rightarrow BP\langle n \rangle_{\mathcal{D}}^k(m)(M)$$

is surjective for $k \leq 2(p^n + \dots + 1)$.

Furthermore, let $I^k \langle n \rangle (m)$ be the subgroup of elements in $BP_{\mathcal{D}}^k(m)(M)$ which can be written as a finite sum

$$u = \sum_{i > n} v_{i,m}(u_i)$$

with $u_i \in BP_{\mathcal{D}}^{k+2(p^i-1)}(m)(M)$, $v_i \in BP^{-2(p^i-1)}$, and $v_{i,m}$ the induced map $BP_{\mathcal{D}}^*(m)(M) \rightarrow BP_{\mathcal{D}}^{*+2(p^i-1)}(m)(M)$. Then Theorem 1.1 has the the following consequence.

Theorem 1.2 *Let M be a complex manifold and m and $n \geq 0$ be integers. The natural induced homomorphism*

$$BP_{\mathcal{D}}^k(m)(M)/I^k \langle n \rangle (m) \rightarrow BP \langle n \rangle_{\mathcal{D}}^k(m)(M)$$

is an isomorphism for $k \leq 2(p^n + \dots + 1)$ and injective for $k \leq 2(p^n + \dots + 1) + 2$.

For $n = 0$, we have $BP \langle 0 \rangle^* = \mathbb{Z}_{(p)}$ and $BP \langle 0 \rangle_{\mathcal{D}}^k(m)(M) = H_{\mathcal{D}}^k(M; \mathbb{Z}_{(p)}(m))$, where $H_{\mathcal{D}}^k(M; \mathbb{Z}_{(p)}(m))$ denotes the k th analytic Deligne cohomology of M with coefficients in $\mathbb{Z}_{(p)}(m)$. Hence Theorem 1.2 has the following special case which is an analog of [15, Theorem 2.2] for Deligne cohomology.

Corollary 1.3 *Let M be a complex manifold and m an integer. The natural map*

$$BP_{\mathcal{D}}^k(m)(M)/I^k \langle 0 \rangle (m) \rightarrow H_{\mathcal{D}}^k(M; \mathbb{Z}_{(p)}(m))$$

is an isomorphism for $k \leq 2$ and injective for $k \leq 4$. In particular, the quotient $BP_{\mathcal{D}}^k(m)(M)/I^k \langle 0 \rangle (m)$ vanishes for negative k .

The fact that $BP_{\mathcal{D}}^k(m)(M)/I^k \langle 0 \rangle (m)$ vanishes in negative degrees could also be reformulated as the following analog of Quillen’s theorem.

Theorem 1.4 *Let M be a complex manifold and m an integer. Then $BP_{\mathcal{D}}^*(m)(M)$ is generated as a BP^* -module by elements of non-negative degree.*

In [15], Totaro showed that Brown–Peterson cohomology and the map in Corollary 1.3 (for BP and H instead of $BP_{\mathcal{D}}$ and $H_{\mathcal{D}}$) are very useful tools for the study of the cycle maps for smooth projective complex algebraic varieties from Chow groups to singular cohomology and Deligne cohomology, respectively. In [8,13], Hopkins and the author used Hodge filtered complex cobordism to study cycles in algebraic cobordism. We are optimistic that Theorem 1.2 will play an important role in the study of various related cycle maps in the future.

The author would like to thank Mike Hopkins for many interesting suggestions and conversations and the anonymous referee for helpful suggestions for improving the paper.

2 Hodge filtered function spaces

We start with a brief recollection of Eilenberg–MacLane spaces, the singular functor and Hodge filtered forms in the setting of simplicial presheaves. Then we will construct Hodge filtered function spaces.

2.1 Simplicial presheaves

Let \mathbf{T} be the category $\mathbf{Man}_{\mathbb{C}}$ of complex manifolds and holomorphic maps. The Grothendieck topology defined by open coverings turns \mathbf{T} into an essentially small site with enough points. We denote by $\mathbf{sPre} = \mathbf{sPre}(\mathbf{T})$ the category of simplicial presheaves on \mathbf{T} , i.e., contravariant functors from \mathbf{T} to the category \mathbf{sS} of simplicial sets. Objects in \mathbf{sPre} will also be called *spaces*. Recall that sending an object M of \mathbf{T} to the presheaf of sets it represents defines a fully faithful embedding of \mathbf{T} into the category of presheaves of sets on \mathbf{T} . Since any presheaf of sets defines an object in \mathbf{sPre} of simplicial dimension zero, we can embed \mathbf{T} into \mathbf{sPre} . On the other hand, every simplicial set K defines a simplicial presheaf by sending every object to K . By abuse of notation, we denote this simplicial presheaf by K as well.

We will consider \mathbf{sPre} with the local projective model structure (see e.g. [5,6]). We will not discuss the specific properties of this model structure, but just recall that a map $\mathcal{F} \rightarrow \mathcal{G}$ in \mathbf{sPre} is called a (local) weak equivalence if the induced map of stalks $\mathcal{F}_x \rightarrow \mathcal{G}_x$ is a weak equivalence in \mathbf{sS} for every point x in \mathbf{T} . Furthermore, a map $\mathcal{F} \rightarrow \mathcal{G}$ is called an objectwise fibration if $\mathcal{F}(X) \rightarrow \mathcal{G}(X)$ is a fibration in \mathbf{sS} for every $X \in \mathbf{T}$. A map is a local projective fibration if it is an objectwise fibration and satisfies descent for all hypercovers in \mathbf{T} (see [6, Corollary 7.1]). We denote the corresponding homotopy category of \mathbf{sPre} by \mathbf{hosPre} .

A natural way to send a CW -complex into \mathbf{sPre} is the singular functor which is defined as follows. Let Δ^n be the standard topological n -simplex

$$\Delta^n = \left\{ (t_0, \dots, t_n) \in \mathbb{R}^{n+1} \mid 0 \leq t_j \leq 1, \sum t_j = 1 \right\}.$$

For topological spaces Y and Z , the singular function complex $\text{Sing}_*(Z, Y)$ is the simplicial set whose n -simplices are continuous maps

$$f : Z \times \Delta^n \rightarrow Y.$$

We denote the simplicial presheaf

$$M \mapsto \text{Sing}_*(M, Y) =: \text{Sing}_*Y(M)$$

on $\mathbf{Man}_{\mathbb{C}}$ by Sing_*Y . For any CW -complex Y , the simplicial presheaf Sing_*Y is objectwise fibrant and satisfies descent for hypercovers in $\mathbf{Man}_{\mathbb{C}}$ by [7, Theorem 1.3] (see also [8, Lemma 2.3]). By [6, Corollary 7.1], this implies that Sing_*Y is a fibrant object in the local projective model structure on \mathbf{sPre} .

Furthermore, for a simplicial set K , let $|K|$ be its geometric realization in the category of CW -complexes. By [8, Proposition 2.4], the natural map

$$K \rightarrow \text{Sing}_*|K|$$

is a weak equivalence of simplicial presheaves. Hence we can use the assignment $K \mapsto \text{Sing}_*|K|$ as a natural fibrant replacement in \mathbf{sPre} for simplicial presheaves coming from simplicial sets.

An important class of simplicial presheaves are Eilenberg–MacLane spaces. Let \mathcal{C}^* be a cochain complex of presheaves of abelian groups on \mathbf{T} . For any integer n , we denote by $\mathcal{C}^*[n]$ the cochain complex given in degree q by $\mathcal{C}^q[n] := \mathcal{C}^{q+n}$. The differential on $\mathcal{C}^*[n]$ is the one of \mathcal{C}^* multiplied by $(-1)^n$. The hypercohomology $H^*(M; \mathcal{C}^*)$ of an object M of \mathbf{T} with coefficients in \mathcal{C}^* is the graded group of morphisms $\text{Hom}(\mathbb{Z}_M, a\mathcal{C}^*)$ in the derived category of cochain complexes of sheaves on \mathbf{T} , where $a\mathcal{C}^*$ denotes the complex of associated sheaves of \mathcal{C}^* . We will denote by $K(\mathcal{C}^*, n)$ the Eilenberg–MacLane spaces, i.e., simplicial presheaf, associated to $\mathcal{C}^*[-n]$. The following result is a version of Verdier’s hypercovering theorem due to Ken Brown.

Proposition 2.1 [3, Theorem 2] *Let \mathcal{C}^* be a cochain complex of presheaves of abelian groups on \mathbf{T} . Then for any integer n and any object M of \mathbf{T} , one has a canonical isomorphism*

$$H^n(M; \mathcal{C}^*) \cong \text{Hom}_{\text{hosPre}}(M, K(\mathcal{C}^*, n)).$$

Example 2.2 Let Ω_M^n denote the sheaf of holomorphic n -forms on a complex manifold M and let Ω^* be the complex of presheaves on $\mathbf{Man}_{\mathbb{C}}$ that sends a complex manifold M to the complex $\Omega_M^*(M)$. The inclusion of complexes $\mathbb{C} \hookrightarrow \Omega^*$ is a quasi-isomorphism and induces a weak equivalence of simplicial presheaves

$$K(\mathbb{C}, k) \rightarrow K(\Omega^*, k).$$

This implies that there is a natural isomorphism

$$H^k(M; \mathbb{C}) \cong \text{Hom}_{\text{hosPre}}(M, K(\Omega^*, k))$$

for every k and $M \in \mathbf{Man}_{\mathbb{C}}$.

2.2 Hodge filtration on forms

For a complex manifold M , let Ω_M^* again denote the complex of sheaves of holomorphic forms on M . Let \mathcal{V}_* be an evenly graded \mathbb{C} -vector space such that each \mathcal{V}_{2i} is a finite dimensional complex vector space. For an integer n , we denote by $H^n(M; \mathcal{V}_*)$ the graded cohomology group

$$H^n(M; \mathcal{V}_*) := \bigoplus_i H^{n+2i}(M; \mathcal{V}_{2i})$$

where $H^{n+2i}(M; \mathcal{V}_{2i}) = H^{n+2i}(M; \mathbb{C}) \otimes_{\mathbb{C}} \mathcal{V}_{2i}$.

The groups $H^n(M; \mathcal{V}_*)$ are equipped with a Hodge filtration which can be defined as follows (see e.g. [16, Def. 8.2]). Let Ω_M^* be the complex of sheaves of holomorphic forms on M and let $\Omega_M^{*\geq p}$ be the subcomplex of forms of degree at least p . The Hodge filtration on $H^n(M; \mathbb{C})$ can be defined by

$$F^p H^n(M; \mathbb{C}) := \text{Im}(\mathbb{H}^n(M; \Omega_M^{*\geq p}) \rightarrow \mathbb{H}^n(M; \Omega_M^*) = H^n(M; \mathbb{C}))$$

where $\mathbb{H}^n(M; -)$ denotes the n th hypercohomology. If M is a compact Kähler manifold, then $F^p H^n(M; \mathbb{C})$ is just $\mathbb{H}^n(M; \Omega_M^{*\geq p})$.

For a general complex manifold, we extend the Hodge filtration to the graded groups $H^n(M; \mathcal{V}_*)$ as follows. For integers m and n , let $F^{m+i} H^{n+2i}(M; \mathcal{V}_{2i})$ be the $(m+i)$ th step in the Hodge filtration of $H^{n+2i}(M; \mathcal{V}_{2i})$. We will write

$$F^m H^n(M; \mathcal{V}_*) := \bigoplus_i F^{m+i} H^{n+2i}(M; \mathcal{V}_{2i})$$

for the graded Hodge filtered cohomology groups of M .

Let

$$\Omega_M^* \rightarrow A_M^* \text{ and } \Omega_M^{*\geq m} \rightarrow F^m A_M^*$$

be resolutions by cohomologically trivial sheaves which are functorial in M . We can choose these resolutions in such a way as to fit into a commutative diagram

$$\begin{CD} \Omega_M^* @>>> \cdots @>>> \Omega_M^{*\geq m-1} @>>> \Omega_M^{*\geq m} @>>> \cdots \\ @VVV @. @VVV @VVV @. \\ A_M^* @>>> \cdots @>>> F^{m-1} A_M^* @>>> F^m A_M^* @>>> \cdots \end{CD}$$

For example, we could use the Godement resolution [4, §3.2.3]. Let A^* and $F^m A^*$ be the associated presheaves of complexes on $\mathbf{Man}_{\mathbb{C}}$ defined by

$$A^*: M \mapsto A_M^*(M) \text{ and } F^m A^*: M \mapsto F^m A_M^*(M).$$

Note that even though A^* and $F^m A^*$ are double complexes, we will only consider their total complexes.

We denote by $A^*(\mathcal{V}_{2i})[-2i]$ the presheaf of forms with coefficients in \mathcal{V}_{2i} shifted by degree $2i$ and we will write

$$A^*(\mathcal{V}_*) := \bigoplus_i A^*(\mathcal{V}_{2i})[-2i].$$

For an integer m , we define the complex of presheaves $F^m A^*(\mathcal{V}_*)$ to be

$$F^m A^*(\mathcal{V}_*) := \bigoplus_i F^{m+i} A^*(\mathcal{V}_{2i})[-2i]. \tag{3}$$

For an integer n , let $K(A^*(\mathcal{V}_*), n)$ and $K(F^m A^*(\mathcal{V}_*), n)$ denote the associated Eilenberg–MacLane spaces. Note that (3) induces isomorphisms

$$\begin{aligned} K(A^*(\mathcal{V}_*), n) &\cong \bigvee_i K(A^*(\mathcal{V}_{2i}), n + 2i) \\ K(F^m A^*(\mathcal{V}_*), n) &\cong \bigvee_i K(F^{m+i} A^*(\mathcal{V}_{2i}), n + 2i). \end{aligned}$$

Recall that $|\cdot|$ denotes the geometric realization functor which sends simplicial sets to CW-complexes. The simplicial presheaf $\text{Sing}_*|K(\mathcal{V}_*, n)|$ represents the functor of cocycles with coefficients in \mathcal{V}_* , i.e., for every $M \in \mathbf{Man}_{\mathbb{C}}$, there is a natural isomorphism of abelian groups

$$Z^n(M; \mathcal{V}_*) \cong \text{Hom}_{\mathbf{sPre}}(M, \text{Sing}_*|K(\mathcal{V}_*, n)|),$$

where we write $Z^n(M; \mathcal{V}_*) := \bigoplus_i Z^{n+2i}(M; \mathcal{V}_{2i})$ for the graded group of cocycles on M . Since M is a representable presheaf, we have a natural bijection of sets

$$\text{Hom}_{\mathbf{sPre}}(M, \text{Sing}_*|K(\mathcal{V}_*, n)|) \cong \text{Sing}_0|K(\mathcal{V}_*, n)|(M).$$

Moreover, since M is a cofibrant object in the local projective model structure on \mathbf{sPre} , there are natural bijections

$$\text{Hom}_{\text{hosPre}}(M, \text{Sing}_*|K(\mathcal{V}_*, n)|) \cong \pi_0(\text{Sing}_*|K(\mathcal{V}_*, n)|(M)) \cong H^n(M; \mathcal{V}_*). \tag{4}$$

Since the canonical inclusion $\mathcal{V}_* \hookrightarrow A^*_M(\mathcal{V}_*)$ is a quasi-isomorphism of complexes of sheaves for every M , it induces a weak equivalence of simplicial presheaves $K(\mathcal{V}_*, n) \rightarrow K(A^*(\mathcal{V}_*), n)$. Together with (4), this implies that there are natural bijections

$$\begin{aligned} \text{Hom}_{\text{hosPre}}(M, \text{Sing}_*|K(A^*(\mathcal{V}_*), n)|) &\cong \pi_0(\text{Sing}_*|K(A^*(\mathcal{V}_*), n)|(M)) \\ &\cong H^n(A^*(M; \mathcal{V}_*)) \\ &\cong H^n(M; \mathcal{V}_*). \end{aligned}$$

If M is a compact Kähler manifold, we even have natural bijections

$$\begin{aligned} \text{Hom}_{\text{hosPre}}(M, \text{Sing}_*|K(F^m A^*(\mathcal{V}_*), n)|) &\cong \pi_0(\text{Sing}_*|K(F^m A^*(\mathcal{V}_*), n)|(M)) \\ &\cong H^n(F^m A^*(M; \mathcal{V}_*)) \\ &\cong F^m H^n(M; \mathcal{V}_*). \end{aligned} \tag{5}$$

The sequence of isomorphisms in (5) expresses that, at least for compact Kähler manifolds, the simplicial presheaves $\text{Sing}_*|K(F^m A^*(\mathcal{V}_*), n)|$, for $n \geq 0$, represent Hodge filtered cohomology groups in hosPre . We remark that there is an analogous result for smooth complex varieties (see [12]).

2.3 Hodge filtered function spaces

We will now define Hodge filtered function spaces. The idea is similar to the way that differential function spaces were defined for smooth manifolds in [9].

Let m and n be integers and \mathcal{V}_* an evenly graded complex vector space. Let Y be a CW-complex and let $\iota \in Z^n(Y; \mathcal{V}_*)$ by a cocycle on Y . A cocycle corresponds to a map of CW-complexes

$$Y \rightarrow |K(\mathcal{V}_*, n)|$$

and induces a map of simplicial presheaves on $\mathbf{Man}_{\mathbb{C}}$

$$\text{Sing}_* Y \rightarrow \text{Sing}_*|K(\mathcal{V}_*, n)|.$$

The canonical inclusion $\mathcal{V}_* \hookrightarrow A^*(\mathcal{V}_*)$ induces a map $K(\mathcal{V}_*, n) \rightarrow K(A^*(\mathcal{V}_*), n)$, and we can form the following diagram of simplicial presheaves

$$\begin{array}{ccc} & \text{Sing}_* Y & (6) \\ & \downarrow \iota^* & \\ \text{Sing}_*|K(F^m A^*(\mathcal{V}_*), n)| & \longrightarrow & \text{Sing}_*|K(A^*(\mathcal{V}_*), n)|. \end{array}$$

Definition 2.3 We define the *Hodge filtered function space* $(Y(m), \iota, n)$ to be the homotopy pullback of (6) in \mathbf{sPre} .

Remark 2.4 Note that $(Y(m), \iota, n)$ depends on ι only up to homotopy, i.e., if ι' is another cocycle which represents the same cohomology class as ι , then $(Y(m), \iota, n)$ and $(Y(m), \iota', n)$ are equivalent.

Remark 2.5 For $Y = K(\mathbb{Z}, n)$ and $\mathcal{V}_* = \mathbb{C}$ concentrated in degree 0, we recover analytic Deligne cohomology for complex manifolds. Since this observation is one of the key motivations for the construction of Hodge filtered function spaces, we are now going to explain this fact in more detail.

Recall that, for given integers $m \geq 0$ and $n \geq 0$, the n th Deligne cohomology group $H_{\mathcal{D}}^n(M; \mathbb{Z}(m))$ of a complex manifold M is defined as the n th hypercohomology of the complex of sheaves

$$0 \rightarrow \mathbb{Z} \xrightarrow{\cdot(2\pi i)^m} \Omega_M^0 \rightarrow \Omega_M^1 \rightarrow \dots \rightarrow \Omega_M^{m-1} \rightarrow 0 \tag{7}$$

where \mathbb{Z} is placed in degree 0 and the map from the locally constant sheaf \mathbb{Z} to the sheaf of holomorphic functions Ω_M^0 is the natural inclusion multiplied by $(2\pi i)^m$ (i

being a square root of -1). Complex (7) is equivalent to the homotopy pullback of complexes of sheaves of the diagram

$$\begin{array}{ccc} & \mathbb{Z} & \\ & \downarrow (2\pi i)^m & \\ \Omega_M^{*\geq m} & \longrightarrow & \Omega_M^* \end{array} \tag{8}$$

After replacing Ω_M^* with the complex of presheaves A^* and after applying the Eilenberg–MacLane functor $K(-, n)$ we obtain the diagram in **sPre**

$$\begin{array}{ccc} & \text{Sing}_* K(\mathbb{Z}, n) & \\ & \downarrow \text{Sing}_* \iota & \\ \text{Sing}_* |K(F^m A^*, n)| & \longrightarrow & \text{Sing}_* |K(A^*, n)| \end{array} \tag{9}$$

where $\iota: K(\mathbb{Z}, n) \rightarrow K(A^*, n)$ is the map that is induced by the $(2\pi i)^m$ -multiple of $\mathbb{Z} \hookrightarrow \Omega^0 \rightarrow A^0$. Since the Eilenberg–MacLane functor is an equivalence between the category of presheaves of complexes of abelian groups and the category of presheaves of simplicial abelian groups, there is an isomorphism between the n th hypercohomology of the homotopy pullback of (8) and π_0 of the homotopy pullback of (9). In other words, the simplicial presheaf

$$K(\mathbb{Z}, n)(m) := (K(\mathbb{Z}, n)(m), \iota, n)$$

(which is the homotopy pullback of (9)) represents analytic Deligne cohomology in **sPre**, i.e., for every $M \in \mathbf{Man}_{\mathbb{C}}$, there is a natural isomorphism

$$H_{\mathcal{D}}^n(M; \mathbb{Z}(m)) \cong \text{Hom}_{\text{hosPre}}(M, K(\mathbb{Z}, n)(m)).$$

2.4 An alternative definition

An equivalent way to construct $(Y(m), \iota, n)$ is the following. For a complex manifold M , let $Z^n(M \times \Delta^\bullet; \mathcal{V}_*)$ be the simplicial abelian group whose group of k -simplices is given by C^∞ - n -cocycles on $M \times \Delta^k$ with coefficients in \mathcal{V}_* . We denote the corresponding simplicial presheaf

$$M \mapsto Z^n(M \times \Delta^\bullet; \mathcal{V}_*)$$

on $\mathbf{Man}_{\mathbb{C}}$ by $Z^n(- \times \Delta^\bullet; \mathcal{V}_*)$. A cocycle ι determines a map of simplicial presheaves

$$\text{Sing}_* Y \rightarrow Z^n(- \times \Delta^\bullet; \mathcal{V}_*), \quad f \mapsto \iota^* f,$$

given by taking the pullback along ι . Let I denote the map given by integration of forms

$$I: F^{m+i} A^{n+2i}(M; \mathcal{V}_{2i}) \rightarrow C^{n+2i}(M; \mathcal{V}_{2i}), \eta \mapsto \left(\sigma \mapsto \int_{\Delta^{n+2i}} \sigma^* \eta \right).$$

We can form a diagram of simplicial presheaves

$$\begin{array}{ccc} & \text{Sing}_* Y & (10) \\ & \downarrow \iota^* & \\ \text{Sing}_* |K(F^m A^*(\mathcal{V}_*), n)| & \xrightarrow{I} & Z^n(- \times \Delta^\bullet; \mathcal{V}_*). \end{array}$$

The map $\text{Sing}_* K(\mathcal{V}_*, n)(M) \rightarrow Z^n(M \times \Delta^\bullet; \mathcal{V}_*)$ given by pulling back a fundamental cocycle in $Z^n(K(\mathcal{V}_*, n); \mathcal{V}_*)$ is a simplicial homotopy equivalence (see e.g. [9, Proposition A.12]). Hence the homotopy pullback of (6) is homotopy equivalent to the homotopy pullback of (10).

For given m, n and ι , let us write $Y(m)$ for the homotopy pullback of (10). To make the construction more concrete, we describe the 0-simplices of $Y(m)(M)$ for a complex manifold M . Since one can calculate homotopy pullbacks in **sPre** objectwise (see e.g. [13, Proposition 2.7]), we can assume that $Y(m)(M)$ is the homotopy pullback of the diagram of simplicial sets

$$\begin{array}{ccc} & \text{Sing}_* Y(M) & \\ & \downarrow \iota^* & \\ \text{Sing}_* |K(F^m A^*(\mathcal{V}_*), n)|(M) & \xrightarrow{I} & Z^n(M \times \Delta^\bullet; \mathcal{V}_*). \end{array}$$

A 0-simplex of $Y(m)(M)$ is given by a triple

$$q: M \rightarrow Y, \eta \in F^m A^n(M; \mathcal{V}_*)_{\text{cl}}, h \in C^{n-1}(M; \mathcal{V}_*)$$

where q is a continuous map and η is a closed form such that $\delta h = \iota^* q - I(\eta)$, where δ denotes the differential in $C^*(M; \mathcal{V}_*)$.

3 Hodge filtered function spaces and spectra

Our main case of interest is the construction of spaces in the Hodge filtered spectra defined in [8, §4]. We will first define such Hodge filtered function spaces and then explain how maps between loop spaces induce maps between Hodge filtered function spaces.

3.1 Spaces in Hodge filtered Ω -spectra

Let m be an integer. Let E be a connective Ω -spectrum built by CW -complexes. We assume that E is rationally even, i.e., $\pi_* E \otimes \mathbb{Q}$ is concentrated in even degrees, and finitely generated in each degree. For example, E could be either BP or $BP\langle n \rangle$.

Let E_k be the k th space of E . By our assumption on E , we have $\pi_{*+k} E_k = \pi_* E$. We set $\pi_* E_{\mathbb{C}} = \pi_* E \otimes_{\mathbb{Z}} \mathbb{C}$ and let

$$\mu_{E_k} : \pi_{*+k} E_k \rightarrow \pi_* E_{\mathbb{C}}$$

be the graded homomorphism defined in degree $2j$ by

$$\pi_{2j+k} E_k \rightarrow \pi_{2j} E \otimes_{\mathbb{Z}} \mathbb{C}, \quad x \mapsto x \otimes (2\pi i)^{j+m}.$$

The homomorphism μ_{E_k} corresponds to a cohomology class $c_{E_k} \in H^k(E_k; \pi_* E_{\mathbb{C}})$ under the Hurewicz isomorphism

$$H^k(E_k; \pi_* E_{\mathbb{C}}) \cong \text{Hom}(\pi_{*+k} E_k, \pi_* E_{\mathbb{C}}).$$

Let

$$\iota_{E_k} : E_k \rightarrow |K(\pi_* E_{\mathbb{C}}, k)| \tag{11}$$

be the map which represents a cocycle in $Z^k(E_k; \pi_* E_{\mathbb{C}})$ whose cohomology class is c_{E_k} . The choice of such a map is unique up to homotopy.

Definition 3.1 We call ι_{E_k} an m -twisted fundamental cocycle of E_k .

We will also need the following variation of such fundamental cocycles. The inclusion $\pi_* E_{\mathbb{C}} \hookrightarrow A^*(\pi_* E_{\mathbb{C}})$ induces a map of simplicial presheaves

$$K(\pi_* E_{\mathbb{C}}, k) \rightarrow K(A^*(\pi_* E_{\mathbb{C}}), k).$$

Composition with (11) defines a map in **sPre**

$$\iota_{E_k} : E_k \rightarrow |K(A^*(\pi_* E_{\mathbb{C}}), k)|$$

which we also denote by ι_{E_k} .

We can form the diagram in **sPre**

$$\begin{array}{ccc} & & \text{Sing}_* E_k \\ & & \downarrow \iota_{E_k} \\ \text{Sing}_* |K(F^m A^*(\pi_* E_{\mathbb{C}}), k)| & \longrightarrow & \text{Sing}_* |K(A^*(\pi_* E_{\mathbb{C}}), k)|. \end{array} \tag{12}$$

We write $(E_k(m), \iota_{E_k})$ for the homotopy pullback of (12) in **sPre**. Note that a different choice ι'_{E_k} of an m -twisted fundamental cocycle of E yields a homotopy equivalent simplicial presheaf $(E_k(m), \iota'_{E_k})$. Therefore, we will usually drop ι_{E_k} from the notation and write $E_k(m)$ for $(E_k(m), \iota_{E_k})$.

Definition 3.2 We call $E_k(m)$ the m th Hodge filtered function space of E_k .

The relationship between the spaces $E_k(m)$ and the spectra $E_{\mathcal{D}}(m)$ defined in [8, §4] is summarized in the following theorem.

Theorem 3.3 For each m , we can choose the cocycles ι_{E_k} such that $\{E_k(m)\}_k$ forms an Ω -spectrum in the category of presheaves of spectra which is equivalent to the spectrum $E_{\mathcal{D}}(m)$ of [8, §4]. For each k , the simplicial presheaf $E_k(m)$ represents Hodge filtered E -cohomology groups of degree k and twist m in \mathbf{hosPre} , i.e., for any $M \in \mathbf{Man}_{\mathbb{C}}$, there is a natural isomorphism

$$\mathrm{Hom}_{\mathbf{hosPre}}(M, E_k(m)) \cong E_{\mathcal{D}}^k(m)(M).$$

Proof First, we need to show that we can choose the cocycles ι_{E_k} such that we obtain structure maps

$$\sigma_k(m) : E_k(m) \rightarrow \Omega E_{k+1}(m)$$

which are weak equivalences for every k .

Let

$$\mu_E : \pi_* E \rightarrow \pi_* E_{\mathbb{C}}$$

be the graded homomorphism defined by multiplication by $(2\pi i)^{j+m}$ in degree $2j$. The homomorphism μ_E corresponds to a cohomology class $c_E \in H^0(E; \pi_* E_{\mathbb{C}})$ under the Hurewicz isomorphism

$$H^0(E; \pi_* E_{\mathbb{C}}) \cong \mathrm{Hom}(\pi_* E, \pi_* E_{\mathbb{C}}).$$

The class c_E in $H^0(E; \pi_* E_{\mathbb{C}})$ can be represented by a map of spectra

$$\iota_E : E \rightarrow H(\pi_* E_{\mathbb{C}})$$

where $H(\pi_* E_{\mathbb{C}})$ denotes the Eilenberg–MacLane spectrum associated to the graded \mathbb{C} -vector space $\pi_* E_{\mathbb{C}}$. This map consists of a family of maps

$$\iota_{E_k} : E_k \rightarrow |K(\pi_* E_{\mathbb{C}}, k)|$$

which are compatible with the structure maps of the spectra E and $H(\pi_* E_{\mathbb{C}})$, at least up to homotopy. More precisely, if $\sigma_k : E_k \rightarrow \Omega E_{k+1}$ denotes the k th structure map of E , these cocycles induce homotopy commutative diagrams of the form

$$\begin{array}{ccc} E_k & \xrightarrow{\iota_{E_k}} & |K(\pi_* E_{\mathbb{C}}, k)| \\ \sigma_k \downarrow & & \downarrow \\ \Omega E_{k+1} & \xrightarrow{\Omega \iota_{E_{k+1}}} & \Omega |K(\pi_* E_{\mathbb{C}}, k+1)| \end{array}$$

where the right hand vertical map is the k th structure map of $H(\pi_* E_{\mathbb{C}})$.

Hence, since σ_k and ι_{E_k} are compatible, we obtain a homotopy commutative diagram

$$\begin{array}{ccccc}
 \text{Sing}_* E_k & \xrightarrow{\iota_{E_k}} & \text{Sing}_* |K(A^*(\pi_* E_{\mathbb{C}}), k)| & \longleftarrow & \text{Sing}_* |K(F^m A^*(\pi_* E_{\mathbb{C}}), k)| \\
 \sigma_k \downarrow & & \downarrow & & \downarrow \\
 \text{Sing}_* \Omega E_{k+1} & \xrightarrow{\Omega \iota_{E_{k+1}}} & \text{Sing}_* \Omega |K(A^*(\pi_* E_{\mathbb{C}}), k+1)| & \longleftarrow & \text{Sing}_* \Omega |K(F^m A^*(\pi_* E_{\mathbb{C}}), k+1)|.
 \end{array}$$

Since E is an Ω -spectrum, the vertical maps are weak equivalences in **sPre**. Hence the map from the homotopy pullback of the upper row to the homotopy pullback of the lower row, which is induced by the vertical maps, is a weak equivalence in **sPre**. Since taking loop spaces commutes with the singular functor and with taking homotopy pullbacks, this shows that the diagram above induces a map of simplicial presheaves

$$\sigma_k(m) : E_k(m) \rightarrow \Omega E_{k+1}(m)$$

which is a weak equivalence in **sPre**. This proves that the family of simplicial presheaves $E_k(m)$ together with the maps $\sigma_k(m)$, indexed by k , forms a presheaf of spectra which we denote by $E(m)$. It follows from the construction of $E(m)$ and the presheaf of spectra $E_{\mathcal{D}}(m)$ in [8, §4.1] that there is a canonical map of presheaves of spectra

$$E(m) \rightarrow E_{\mathcal{D}}(m). \tag{13}$$

Now let M be a complex manifold. It follows from the definition of $E_k(m)$ as a homotopy pullback that the group $\text{Hom}_{\text{hosPre}}(M, E_k(m))$ sits in a long exact sequence analog to the one of [8, Proposition 4.5]. Moreover, the map (13) induces a natural morphism of long exact sequences

$$\begin{array}{ccccccc}
 \dots & \longrightarrow & E_{\mathbb{C}}^{k-1}(M) & \longrightarrow & \text{Hom}_{\text{hosPre}}(M, E_k(m)) & \longrightarrow & E^k(M) \oplus H^k(M; F^m A^*(\pi_* E_{\mathbb{C}})) \longrightarrow \dots \\
 & & \downarrow & & \downarrow & & \downarrow \\
 \dots & \longrightarrow & E_{\mathbb{C}}^{k-1}(M) & \longrightarrow & E_{\mathcal{D}}^k(m)(M) & \longrightarrow & E^k(M) \oplus H^k(M; \Omega^{*\geq m}(\pi_* E_{\mathbb{C}})) \longrightarrow \dots
 \end{array} \tag{14}$$

Since the outer vertical maps are isomorphisms, the induced map

$$\text{Hom}_{\text{hosPre}}(M, E_k(m)) \xrightarrow{\cong} E_{\mathcal{D}}^k(m)(M)$$

is an isomorphism as well. Since we can assume that $E_{\mathcal{D}}(m)$ is an Ω -spectrum, this also implies that the map (13) is an objectwise and hence also stalkwise weak equivalence of spectra. □

Remark 3.4 In order to indicate how the groups $E_{\mathcal{D}}^k(m)(M)$ look like, let us assume $k = 2m$ and that M is a compact Kähler manifold. In this case, one can show using Hodge theory as in [8, §4.3] that the (upper) long exact sequence in (14) can be split into short exact sequences of the form

$$0 \rightarrow E^{2m-1}(M) \otimes_{\mathbb{Z}} \mathbb{R}/\mathbb{Z} \rightarrow E_D^{2m}(m)(M) \rightarrow \text{Hdg}_E^{2m}(M) \rightarrow 0$$

where $\text{Hdg}_E^{2m}(M)$ is the subgroup of $E^{2m}(M)$ that is defined by the cartesian square

$$\begin{array}{ccc} \text{Hdg}_E^{2m}(M) & \longrightarrow & E^{2m}(M) \\ \downarrow & & \downarrow \\ F^m H^{2m}(M; \pi_* E_{\mathbb{C}}) & \longrightarrow & H^{2m}(M; \pi_* E_{\mathbb{C}}). \end{array}$$

3.2 Induced maps and products

The construction of Hodge filtered function spaces is functorial in the following way.

Proposition 3.5 *Let m be an integer. Let E and F be connective rationally even Ω -spectra and $f: E_k \rightarrow F_n$ be a map from the k th space of E to the n th space of F .*

(a) *Then f induces a map of Hodge filtered function spaces*

$$f(m): E_k(m) \rightarrow F_n(m).$$

(b) *If f is a weak equivalence, then $f(m)$ is a weak equivalence of simplicial presheaves.*

Proof a) We set $\pi_* E_{\mathbb{C}} := \pi_* E \otimes_{\mathbb{Z}} \mathbb{C}$ and $\pi_* F_{\mathbb{C}} := \pi_* F \otimes_{\mathbb{Z}} \mathbb{C}$. We define graded homomorphisms $\mu_{E_k}: \pi_{*+k} E_k \rightarrow \pi_* E_{\mathbb{C}}$ and $\mu_{F_n}: \pi_{*+n} F_n \rightarrow \pi_* F_{\mathbb{C}}$ by multiplication by $(2\pi i)^{j+m}$ for $* = 2j$.

The given map f induces graded homomorphisms

$$\pi_* E_k \rightarrow \pi_* F_n \text{ and } \pi_* E \rightarrow \pi_{*+(k-n)} F.$$

On the level of Eilenberg–MacLane spaces, f induces a map

$$f_K: K(\pi_* E_{\mathbb{C}}, k) \rightarrow K(\pi_* F_{\mathbb{C}}, n).$$

Now let c_{E_k} be the cohomology class in $H^k(E_k; \pi_* E_{\mathbb{C}})$ corresponding to μ_{E_k} under the Hurewicz isomorphism, and let

$$\iota_{E_k}: E_k \rightarrow |K(\pi_* E_{\mathbb{C}}, k)|$$

be a cocycle in $Z^k(E_k; \pi_* E_{\mathbb{C}})$ whose cohomology class is c_{E_k} . Similarly, let c_{F_n} be the cohomology class in $H^n(F_n; \pi_* F_{\mathbb{C}})$ corresponding to μ_{F_n} under the Hurewicz isomorphism, and let

$$\iota_{F_n}: F_n \rightarrow |K(\pi_* F_{\mathbb{C}}, n)|$$

be a cocycle in $Z^n(F_n; \pi_* F_{\mathbb{C}})$ with cohomology class c_{F_n} . The images of c_{E_k} and c_{F_n} in $H^n(E_k; \pi_* F_{\mathbb{C}})$ under the maps

$$\begin{array}{ccc}
 & H^n(F_n; \pi_* F_{\mathbb{C}}) & \\
 & \downarrow f^* & \\
 H^k(E_k; \pi_* E_{\mathbb{C}}) & \xrightarrow{f_{K*}} & H^n(E_k; \pi_* F_{\mathbb{C}})
 \end{array}$$

induced by f and f_K agree, since they are both equal to the class in $H^n(E_k; \pi_* F_{\mathbb{C}})$ which corresponds to the composed graded homomorphism $\pi_{*+k} E_k \rightarrow \pi_{*+(k-n)} F_{\mathbb{C}}$ under $H^n(E_k; \pi_* F_{\mathbb{C}}) \cong \text{Hom}(\pi_{*+k} E_k, \pi_{*+(k-n)} F_{\mathbb{C}})$. This implies that the diagram

$$\begin{array}{ccc}
 E_k & \xrightarrow{f} & F_n & (15) \\
 \downarrow \iota_{E_k} & & \downarrow \iota_{F_n} & \\
 |K(\pi_* E_{\mathbb{C}}, k)| & \xrightarrow{f_K} & |K(\pi_* F_{\mathbb{C}}, n)| &
 \end{array}$$

commutes up to homotopy.

Hence the map f induces a homotopy commutative diagram of simplicial presheaves

$$\begin{array}{ccccc}
 \text{Sing}_* E_k & \xrightarrow{\iota_{E_k}} & \text{Sing}_* |K(A^*(\pi_* E_{\mathbb{C}}), k)| & \longleftarrow & \text{Sing}_* |K(F^m A^*(\pi_* E_{\mathbb{C}}), k)| \\
 \text{Sing}_* f \downarrow & & \text{Sing}_* f_K \downarrow & & \text{Sing}_* f_K \downarrow \\
 \text{Sing}_* F_n & \xrightarrow{\iota_{F_n}} & \text{Sing}_* |K(A^*(\pi_* F_{\mathbb{C}}), n)| & \longleftarrow & \text{Sing}_* |K(F^m A^*(\pi_* F_{\mathbb{C}}), n)|.
 \end{array} \tag{16}$$

Taking the homotopy pullback of the top row, which is $E_k(m)$, and the homotopy pullback of the bottom row, which is $F_n(m)$, we obtain that f induces a map of simplicial presheaves

$$f(m): E_k(m) \rightarrow F_n(m).$$

b) If $f: E_k \rightarrow F_n$ is a weak equivalence, then the vertical maps in diagram (16) are all weak equivalences. Hence the induced map of homotopy pullbacks $f(m)$ is also a weak equivalence. □

Remark 3.6 Let $f: E_k \rightarrow F_n$ be a map as in Proposition 3.5 and let M be a complex manifold M . One should note that even though $f(m)$ may depend on the chosen cocycles, the induced map on Hodge filtered cohomology groups

$$f(m)_* : \text{Hom}_{\text{hosPre}}(M, E_k(m)) \rightarrow \text{Hom}_{\text{hosPre}}(M, F_n(m))$$

only depends on the homotopy type of f . More precisely, let ι'_{E_k} and ι'_{F_n} be a another choice of cocycles also representing the cohomology classes c_{E_k} and c_{F_n} , respectively, and let f' be a map homotopic to f . This data fits into a homotopy commutative diagram

$$\begin{array}{ccc}
 & E_k & \xrightarrow{f} & F_n & \\
 & \swarrow & & \searrow & \\
 E_k & \xrightarrow{f'} & F_n & & \\
 \downarrow \iota'_{E_k} & & \downarrow \iota'_{F_n} & & \\
 |K(\pi_* E_{\mathbb{C}}, k)| & \xrightarrow{f_K} & |K(\pi_* F_{\mathbb{C}}, n)| & & \\
 \downarrow \iota'_{E_k} & & \downarrow \iota'_{F_n} & & \\
 |K(\pi_* E_{\mathbb{C}}, k)| & \xrightarrow{f'_K} & |K(\pi_* F_{\mathbb{C}}, n)| & &
 \end{array} \tag{17}$$

in which the maps from the back to the front square are homotopy equivalences. This diagram then induces up to coherent homotopy a map of diagrams from (16) to the corresponding diagram built using ι'_{E_k} , ι'_{F_n} and f' . Hence (17) induces a commutative diagram

$$\begin{array}{ccc}
 \text{Hom}_{\text{hosPre}}(M, E_k(m)) & \xrightarrow{f^{(m)*}} & \text{Hom}_{\text{hosPre}}(M, F_n(m)) \\
 \cong \downarrow & & \downarrow \cong \\
 \text{Hom}_{\text{hosPre}}(M, E_k(m)) & \xrightarrow{f'^{(m)*}} & \text{Hom}_{\text{hosPre}}(M, F_n(m))
 \end{array}$$

in which the vertical maps are isomorphisms induced by the homotopies $\iota_{E_k} \sim \iota'_{E_k}$ and $\iota_{F_n} \sim \iota'_{F_n}$

Now we show that Hodge filtered function spaces behave well under taking products. Let E and F be connective rationally even Ω -spectra, and let E_k and F_n be their k th and n th spaces, respectively. We can choose, independently, cocycles ι_{E_k} and ι_{F_n} which represent the homomorphisms μ_E and μ_F as in the beginning of the proof of Proposition 3.5. Since we have

$$\pi_*(E \times F) = \pi_*E \oplus \pi_*F,$$

there is a canonical homotopy equivalence

$$|K(\pi_*(E \times F)_{\mathbb{C}}, k + n)| \xrightarrow{\approx} |K(\pi_*E_{\mathbb{C}}, k)| \times |K(\pi_*F_{\mathbb{C}}, n)|.$$

Hence we can use ι_{E_k} and ι_{F_n} to obtain a cocycle

$$\iota_{E_k \times F_n} : E_k \times F_n \rightarrow |K(A^*(\pi_*(E \times F)_{\mathbb{C}}), k + n)|.$$

We can then form the diagram in **sPre**

$$\text{Sing}_*|K(F^m A^*(\pi_*(E \times F)_{\mathbb{C}}), k + n)| \longrightarrow \text{Sing}_*|K(A^*(\pi_*(E \times F)_{\mathbb{C}}), k + n)|$$

$$\begin{array}{c} \text{Sing}_*(E_k \times F_n) \\ \downarrow \iota_{E_k \times F_n} \end{array}$$

The homotopy pullback of this diagram is the simplicial presheaf $(E_k \times F_n)(m)$.

Lemma 3.7 *Let E_k and F_n be as above. Then, for any integer m , there is a canonical equivalence of Hodge filtered function spaces*

$$(E_k \times F_n)(m) \xrightarrow{\approx} E_k(m) \times F_n(m).$$

Proof This follows from the fact that the singular functor and taking homotopy pullbacks commute with products and preserve weak equivalences. \square

Proposition 3.8 *Let E, F and G be connective rationally even Ω -spectra, and, for integers k, n and j , let $f: E_k \rightarrow F_n \times G_j$ be a continuous map. Then, for any integer m , there is an induced commutative diagram*

$$\begin{array}{ccc} E_k(m) & \longrightarrow & (F_n \times G_j)(m) \\ & \searrow & \downarrow \approx \\ & & F_n(m) \times G_j(m). \end{array}$$

Proof The map f consists of maps $E_k \rightarrow F_n$ and $E_k \rightarrow G_j$ which by Proposition 3.5 induce maps $E_k(m) \rightarrow F_n(m)$ and $E_k(m) \rightarrow G_j(m)$. Together they give the map $E_k(m) \rightarrow F_n(m) \times G_j(m)$. That the diagram commutes follows from the fact that homotopy pullbacks respect products. \square

4 Unstable splitting for Hodge filtered BP-spaces

Let p be a fixed prime number and n a non-negative integer. Let BP denote the Ω -spectrum representing BP -cohomology at p and let $BP\langle n \rangle$ be the Ω -spectrum representing the n th intermediate theory defined in [17]. We write BP_k and $BP\langle n \rangle_k$ for the k th spaces of these spectra. For a given integer m , let $BP_k(m)$ and $BP\langle n \rangle_k(m)$ be the Hodge filtered function spaces associated to BP_k and $BP\langle n \rangle_k$, respectively. Our main result is the following analog of Wilson’s theorem [17, Theorem 5.4].

Theorem 4.1 *Let m and n be integers with $n \geq 0$.*

(a) *For every $k \leq 2(p^n + \dots + 1)$, there is a weak equivalence of simplicial presheaves*

$$BP_k(m) \xrightarrow{\approx} BP\langle n \rangle_k(m) \times \prod_{j>n} BP\langle j \rangle_{k+2(p^j-1)}(m). \tag{18}$$

(b) For every $k \leq 2(p^{n-1} + \dots + 1)$, there is a weak equivalence of simplicial presheaves

$$BP_k \langle n \rangle (m) \xrightarrow{\approx} BP \langle n-1 \rangle_k (m) \times BP \langle n \rangle_{k+2(p^{n-1})} (m). \tag{19}$$

Proof (a) In [17, Corollary 3.6], Wilson shows that, for $k \leq 2(p^n + \dots + 1)$, there is a map

$$BP_k \rightarrow BP \langle n \rangle_k \times \prod_{j>n} BP \langle j \rangle_{k+2(p^j-1)}. \tag{20}$$

By Proposition 3.8, we obtain the map (18) as the induced map of Hodge filtered function spaces. By [17, Corollary 5.4], the map (20) is actually a homotopy equivalence. By the two-out-of-three property for weak equivalences, it follows from the commutative diagram of Proposition 3.8 that the map (18) is a weak equivalence.

(b) The second equivalence is induced in the same way by the equivalence of [17, Corollary 5.5]. □

Together with Theorem 3.3, Theorem 4.1 implies the following result.

Corollary 4.2 *Let M be a complex manifold and m and $n \geq 0$ be integers. Then the natural map*

$$BP_{\mathcal{D}}^k(m)(M) \rightarrow BP \langle n \rangle_{\mathcal{D}}^k(m)(M)$$

is surjective for $k \leq 2(p^n + \dots + 1)$.

As in [17, Theorem 5.7], we can use Corollary 4.2 to deduce an analog of Quillen’s theorem. Recall [17, p.105] that, for given integers k, n and $j := 2(p^n - 1)$, there is a fiber sequence

$$BP \langle n \rangle_{k+j} \xrightarrow{v_n} BP \langle n \rangle_k \xrightarrow{f_n} BP \langle n-1 \rangle_k$$

where v_n denotes the n th generator in $BP^* = \mathbb{Z}_{(p)}[v_1, \dots, v_n, \dots]$. For every integer m , this sequence induces a homotopy commutative diagram in **sPre**

$$\begin{array}{ccccc}
 \text{Sing}_* BP \langle n \rangle_{k+j} & \longrightarrow & \text{Sing}_* BP \langle n \rangle_k & \longrightarrow & \text{Sing}_* BP \langle n-1 \rangle_k \\
 \downarrow & & \downarrow & & \downarrow \\
 \text{Sing}_* |K(A^*(\pi_* BP \langle n \rangle_{\mathbb{C}}, k+j))| & \longrightarrow & \text{Sing}_* |K(A^*(\pi_* BP \langle n \rangle_{\mathbb{C}}, k))| & \longrightarrow & \text{Sing}_* |K(A^*(\pi_* BP \langle n-1 \rangle_{\mathbb{C}}, k))| \\
 \uparrow & & \uparrow & & \uparrow \\
 \text{Sing}_* |K(F^m A^*(\pi_* BP \langle n \rangle_{\mathbb{C}}, k+j))| & \longrightarrow & \text{Sing}_* |K(F^m A^*(\pi_* BP \langle n \rangle_{\mathbb{C}}, k))| & \longrightarrow & \text{Sing}_* |K(F^m A^*(\pi_* BP \langle n-1 \rangle_{\mathbb{C}}, k))|
 \end{array} \tag{21}$$

in which each row is a fiber sequence (the functor Sing_* is right Quillen adjoint and preserves fiber sequences). Since homotopy pullbacks preserve fiber sequences, diagram (21) induces a fiber sequence in **sPre**

$$BP \langle n \rangle_{k+j}(m) \xrightarrow{v_n(m)} BP \langle n \rangle_k(m) \xrightarrow{f_n(m)} BP \langle n-1 \rangle_k(m). \tag{22}$$

Furthermore, there are the maps $g_n : BP_k \rightarrow BP\langle n \rangle_k$ which are compatible with f_n in the sense that $f_n \circ g_n = g_{n-1}$. Each g_n induces a map of simplicial presheaves

$$g_n(m) : BP_k(m) \rightarrow BP\langle n \rangle_k(m).$$

For every complex manifold M , the above maps induce a commutative diagram

$$\begin{CD}
 BP_{\mathcal{D}}^{k+2(p^n-1)}(m)(M) @>v_{n,m}>> BP_{\mathcal{D}}^k(m)(M) \\
 @Vg_{n,m}VV @VVg_{n,m}V \\
 BP\langle n \rangle_{\mathcal{D}}^{k+2(p^n-1)}(m)(M) @>v_{n,m}>> BP\langle n \rangle_{\mathcal{D}}^k(m)(M) @>f_{n,m}>> BP\langle n-1 \rangle_{\mathcal{D}}^k(m)(M)
 \end{CD}
 \tag{23}$$

where the lower row is exact, since (22) is a fiber sequence.

Let $I^k\langle n \rangle(m)$ be the subgroup of elements in $BP_{\mathcal{D}}^k(m)(M)$ which can be written as a finite sum

$$u = \sum_{i>n} v_{i,m}(u_i) \tag{24}$$

with $u_i \in BP_{\mathcal{D}}^{k+2(p^i-1)}(m)(M)$ and $v_i \in BP^{-2(p^i-1)}$. Since the lower row of diagram (23) is exact, the subgroup $I^k\langle n \rangle(m)$ is contained in the kernel of the natural homomorphism

$$BP_{\mathcal{D}}^k(m)(M) \rightarrow BP\langle n \rangle_{\mathcal{D}}^k(m)(M).$$

Theorem 4.3 *Let M be a complex manifold and m and $n \geq 0$ be integers. Then the induced homomorphism*

$$BP_{\mathcal{D}}^k(m)(M)/I^k\langle n \rangle(m) \rightarrow BP\langle n \rangle_{\mathcal{D}}^k(m)(M)$$

is an isomorphism for $k \leq 2(p^n + \dots + 1)$ and injective for $k \leq 2(p^n + \dots + 1) + 2$.

Proof The surjectivity in dimensions $k \leq 2(p^n + \dots + 1)$ follows directly from Corollary 4.2. It remains to check injectivity the proof of which will follow as in [17, Proof of Theorem 5.7].

Let $k \leq 2(p^n + \dots + 1) + 2$ and $u \in BP_{\mathcal{D}}^k(m)(M)$ be an element which maps to 0 in $BP\langle n \rangle_{\mathcal{D}}^k(m)(M)$. Since M has the homotopy type of a finite complex, we know that the natural map is an isomorphism $BP^k(M) \cong BP\langle n \rangle^k(M)$ if n is large enough. The map g_n induces a morphism of long exact sequences

$$\begin{CD}
 \cdots @>>> BP_{\mathbb{C}}^{k-1}(M) @>>> BP_{\mathcal{D}}^k(m)(M) @>>> BP^k(M) \oplus H^k(M; F^m A^*(\pi_* BP_{\mathbb{C}})) @>>> \cdots \\
 @. @VVV @VVV @VVV \\
 \cdots @>>> BP\langle n \rangle_{\mathbb{C}}^{k-1}(M) @>>> BP\langle n \rangle_{\mathcal{D}}^k(m)(M) @>>> BP\langle n \rangle^k(M) \oplus H^k(M; F^m A^*(\pi_* BP\langle n \rangle_{\mathbb{C}})) @>>> \cdots
 \end{CD}
 \tag{25}$$

This shows that the associated Hodge filtered theories satisfy

$$BP_{\mathcal{D}}^k(m)(M) \cong BP\langle n \rangle_{\mathcal{D}}^k(m)(M) \text{ for } n \text{ large enough.}$$

Hence we can find and fix the integer $q > n$ such that

$$g_{q,m}(u) \neq 0 \text{ and } f_{q,m}(g_{q,m}(u)) = g_{q-1,m}(u) = 0.$$

By the exactness of the lower row in diagram (23), there is then an element

$$u' \in BP\langle q \rangle_{\mathcal{D}}^{k+2(p^q-1)}(m)(M) \text{ with } v_{q,m}(u') = g_{q,m}(u).$$

Since $k \leq 2(p^n + \dots + 1) + 2$ and $q \geq n + 1$, we have

$$k + 2(p^q - 1) \leq 2(p^n + \dots + 1) + 2 + 2(p^q - 1) \leq 2(p^q + \dots + 1).$$

Hence, by Corollary 4.2, there is an element u_q in $BP_{\mathcal{D}}^{k+2(p^q-1)}(m)(M)$ such that $g_{q,m}(u_q) = u'$ and $g_{q,m}(v_{q,m}(u_q)) = g_{q,m}(u)$. Since M has the homotopy type of a finite complex, it follows again from the upper long exact sequence in (25) that $BP_{\mathcal{D}}^{k+2(p^j-1)}(m)(M)$ will be zero for j large enough. Hence repeating this process with u replaced by $u - v_{q,m}(u_q)$ shows that u can be written as a finite sum of the form (24) and lies in $I^k\langle n \rangle(m)$. \square

Remark 4.4 As mentioned in the introduction, we hope that Theorem 4.3 will help to find new interesting examples of algebraic cobordism cycles on smooth projective complex algebraic varieties. The idea is to study analogs of the Abel-Jacobi map of [13] for $BP\langle n \rangle$ for various n in order to find families of cycles which are topologically trivial but non-trivial algebraically. Eventually, this should lead to new insights into the map from algebraic to complex cobordism for smooth complex varieties.

References

1. Brown, E.H., Jr., Peterson, F.P.: A spectrum whose \mathbb{Z}_p -cohomology is the algebra of reduced p th powers. *Topology* **5**, 149–154 (1966)
2. Boardman, J.M., Johnson, D.C., Wilson, W.S.: Unstable operations in generalized cohomology. In: James, I.M. (ed.) *Handbook of Algebraic Topology*, pp. 687–828. North-Holland, Amsterdam (1995)
3. Brown, K.S.: Abstract homotopy theory and generalized sheaf cohomology. *Trans. Am. Math. Soc.* **186**, 419–458 (1973)
4. Deligne, P.: Théorie de Hodge II. *Publ. Math. IHES* **40**, 5–57 (1971)
5. Dugger, D.: Universal homotopy theories. *Adv. Math.* **164**, 144–176 (2001)
6. Dugger, D., Hollander, S., Isaksen, D.C.: Hypercovers and simplicial presheaves. *Math. Proc. Camb. Philos. Soc.* **136**(1), 9–51 (2004)
7. Dugger, D., Isaksen, D.C.: Topological hypercovers and \mathbb{A}^1 -realizations. *Math. Z.* **246**, 667–689 (2004)
8. Hopkins, M.J., Quick, G.: Hodge filtered complex bordism. *J. Topol.* **8**, 147–183 (2015)
9. Hopkins, M.J., Singer, I.M.: Quadratic functions in geometry, topology, and M -theory. *J. Differ. Geom.* **70**, 329–452 (2005)
10. Jardine, J.F.: Presheaves of symmetric spectra. *J. Pure Appl. Algebra* **150**, 137–154 (2000)

11. Johnson, D.C., Wilson, W.S.: Projective dimension and Brown–Peterson homology. *Topology* **12**, 327–353 (1973)
12. Quick, G.: Homotopy theory of smooth compactifications of algebraic varieties. *N. Y. J. Math.* **19**, 533–544 (2013)
13. Quick, G.: An Abel–Jacobi invariant for cobordant cycles. *Doc. Math.* **21**, 1645–1668 (2016)
14. Quillen, D.G.: On the formal group laws of unoriented and complex cobordism theory. *Bull. Am. Math. Soc.* **75**, 1293–1298 (1969)
15. Totaro, B.: Torsion algebraic cycles and complex cobordism. *J. Am. Math. Soc.* **10**, 467–493 (1997)
16. Voisin, C.: *Hodge Theory and Complex Algebraic Geometry I*. Cambridge Studies in Advanced Mathematics, vol. 76. Cambridge University Press, Cambridge (2002)
17. Wilson, W.S.: The Ω -spectrum for Brown–Peterson cohomology, part II. *Am. J. Math.* **97**, 101–123 (1975)